

**G0.0+0.0**

**RA:** 17<sup>h</sup>45<sup>m</sup>44<sup>s</sup>  
**Dec:** −29°00'

**1-GHz flux/Jy:** 100?  
**Spectral index:** 0.8?

**Sgr A East**  
**Size/arcmin:** 3.5 × 2.5  
**Type:** S

**Radio:** Non-thermal shell, in complex region, interacting with molecular material to the west.

**References:**

- Ekers *et al.* 1983, A&A, 122, 143. VLA at 1.4 GHz and 5 GHz (both 5'' × 8''),  $S_{1.4 \text{ GHz}} = 77 \text{ Jy}$ ,  $S_{5.0 \text{ GHz}} = 31 \text{ Jy}$ .  
Pedlar *et al.* 1989, ApJ, 342, 769. VLA at 332 MHz (12''), 1.4 GHz (1''.3 × 2''.5) and 5 GHz (1''.3 × 2''.5).  
Mezger *et al.* 1989, A&A, 209, 337. Nearby molecular material.  
Ho *et al.* 1991, Natur, 350, 309. VLA of NH<sub>3</sub> emission from surroundings.  
Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (17'' × 33'').  
Serabyn *et al.* 1992, ApJ, 395, 166. Nearby molecular material.  
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 88'').  
Yusef-Zadeh & Mehringer 1995, ApJ, 452, L37. VLA of nearby H<sub>2</sub>O masers.  
Yusef-Zadeh *et al.* 1999, ApJ, 512, 230. OH maser observations.  
Coil & Ho 2000, ApJ, 533, 245. NH<sub>3</sub> observations of surroundings.  
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').  
Fattuzzo *et al.* 2001, ApJ, 549, 293. Electron–positron lines from the vicinity.  
Yusef-Zadeh *et al.* 2001, ApJ, 560, 749. Observations of nearby molecular hydrogen.

**G0.3+0.0**

**RA:** 17<sup>h</sup>46<sup>m</sup>15<sup>s</sup>  
**Dec:** −28°38'

**1-GHz flux/Jy:** 22  
**Spectral index:** 0.6

**Size/arcmin:** 15 × 8  
**Type:** S

Has been called G0.33+0.04 and G0.4+0.1.

**Radio:** Bilateral shell, near Galactic Centre.

**References:**

- Kassim & Frail 1996, MNRAS, 283, L51. VLA at 333 MHz (23'' × 42''), plus review of flux densities and other observations.  
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').

**G0.9+0.1**

**RA:** 17<sup>h</sup>47<sup>m</sup>21<sup>s</sup>  
**Dec:** −28°09'

**1-GHz flux/Jy:** 18?  
**Spectral index:** varies

**Size/arcmin:** 8  
**Type:** C

**Radio:** Flat spectrum core within steep spectrum shell.

**X-ray:** Central core, with non-thermal spectrum.

**References:**

- Helfand & Becker 1987, ApJ, 314, 203. VLA at 1.4 and 5 GHz, and Einstein observations.  
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 91'').  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
Mereghetti *et al.* 1998, A&A, 331, L77. X-ray detection.  
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').  
Sridoli *et al.* 2000, A&A, 361, 719. BeppoSAX observations.  
Gaensler *et al.* 2001, ApJ, 556, L107. Chandra observations.

**G1.0–0.1**

**RA:** 17<sup>h</sup>48<sup>m</sup>30<sup>s</sup>  
**Dec:** −28°09'

**1-GHz flux/Jy:** 15  
**Spectral index:** 0.6?

**Size/arcmin:** 8  
**Type:** S

Has been called G1.05–0.1 and G1.05–0.15.

**Radio:** Incomplete shell, to the S of Sgr D.

**X-ray:** Possibly detected.

**References:**

- Downes *et al.* 1979, A&AS, 35, 1. Review of flux densities.  
 Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz ( $64'' \times 100''$ :  $S=12.3$  Jy).  
 Liszt 1992, ApJS, 82, 495. VLA at 1.6 GHz ( $13'' \times 23''$ ).  
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ( $43'' \times 91''$ ).  
 Mehringer *et al.* 1998, ApJ, 493, 274. VLA at 1.6 GHz ( $15'' \times 28''$ ) and 5 GHz, including masers observations.  
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.  
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz ( $43'' \times 24''$ ).  
 Sidoli *et al.* 2001, A&A, 372, 651. BeppoSAX possible detection.

**G1.4–0.1**

**RA:** 17<sup>h</sup>49<sup>m</sup>39<sup>s</sup>  
**Dec:** −27°46'

**1-GHz flux/Jy:** 2?  
**Spectral index:** ?

**Size/arcmin:** 10  
**Type:** S

**Radio:** Shell, brightest in E.

**References:**

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ( $43'' \times 92''$ :  $S=2$  Jy).  
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.

**G1.9+0.3**

**RA:** 17<sup>h</sup>48<sup>m</sup>45<sup>s</sup>  
**Dec:** −27°10'

**1-GHz flux/Jy:** 0.6  
**Spectral index:** 0.7

**Size/arcmin:** 1.2  
**Type:** S

**Radio:** Shell, brighter to the N.

**References:**

- Green & Gull 1984, Natur, 312, 527. VLA at 5 GHz ( $4''.4 \times 2''$ ).  
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ( $43'' \times 94''$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G3.7–0.2**

**RA:** 17<sup>h</sup>55<sup>m</sup>26<sup>s</sup>  
**Dec:** −25°50'

**1-GHz flux/Jy:** 2.3  
**Spectral index:** 0.65

**Size/arcmin:** 14 × 11  
**Type:** S

Has been called G003.8–00.3.

**Radio:** Double arc.

**References:**

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ( $43'' \times 99''$ :  $S=2.4$  Jy).  
 Gaensler 1998, ApJ, 493, 781. VLA at 1.4 GHz ( $9'' \times 15''$ :  $S=1.7 \pm 0.1$  Jy).

**G3.8+0.3**

**RA:** 17<sup>h</sup>52<sup>m</sup>55<sup>s</sup>  
**Dec:** −25°28'

**1-GHz flux/Jy:** 4?  
**Spectral index:** ?

**Size/arcmin:** 18  
**Type:** S?

**Radio:** Incomplete shell.

**References:**

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ( $43'' \times 100''$ :  $S=3.5$  Jy).

**G4.2–3.5**

**RA:** 18<sup>h</sup>08<sup>m</sup>55<sup>s</sup>  
**Dec:** −27°03'

**1-GHz flux/Jy:** 3.2?  
**Spectral index:** 0.6?

**Size/arcmin:** 28  
**Type:** S

**Radio:** Elongated shell.

**References:**

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

**G4.5+6.8**

**RA:** 17<sup>h</sup>30<sup>m</sup>42<sup>s</sup>  
**Dec:** −21°29'

**1-GHz flux/Jy:** 19  
**Spectral index:** 0.64

Kepler, SN1604, 3C358

**Size/arcmin:** 3  
**Type:** S

This is the remnant of Kepler's SN of AD1604.

**Radio:** Incomplete shell, brighter to the N.

**Optical:** Faint filaments.

**X-ray:** Shell, brighter to the N.

**Distance:** Optical expansion and proper motion indicates about 2.9 kpc, HI observations suggest 3.4 to 6.4 kpc.

**References:**

van den Bergh & Kamper 1977, ApJ, 218, 617. Optical proper motions.  
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.  
 White & Long 1983, ApJ, 264, 196. Einstein observations.  
 Matsui *et al.* 1984, ApJ, 287, 295. VLA at 1.4 (2''.5 × 3''.2) and 5 GHz (3''.2 × 4''.8) and Einstein image (5'').  
 Dickey *et al.* 1988, ApJ, 330, 254. VLA at 1.4 (1''.2 × 2''.3) and 5 GHz (0''.6 × 1''.0) at two epochs.  
 Smith *et al.* 1989, ApJ, 347, 925. EXOSAT observations.  
 Hatsukade *et al.* 1990, PASJ, 42, 279. X-ray spectrum.  
 Blair *et al.* 1991, ApJ, 366, 484. Optical imaging and spectroscopy.  
 Bandiera & van den Bergh 1991, ApJ, 374, 186. Optical changes.  
 van den Bergh 1991, PASP, 103, 194. Optical imaging.  
 Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Hughes 1999, ApJ, 527, 298. ROSAT and Einstein image comparison for expansion studies.  
 Reynoso & Goss 1999, AJ, 118, 926. VLA at 1.4 GHz (23'' × 13'') for HI studies.  
 Kinugasa & Tsunemi 1999, PASJ, 51, 239. ASCA observations.  
 Gerardy & Fesen 2001, AJ, 121, 2781. IR spectroscopy and imaging.

**G4.8+6.2**

**RA:** 17<sup>h</sup>33<sup>m</sup>25<sup>s</sup>  
**Dec:** −21°34'

**1-GHz flux/Jy:** 3  
**Spectral index:** 0.6

**Size/arcmin:** 18  
**Type:** S

Has been called G4.5+6.2.

**Radio:** Faint shell.

**References:**

Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (2''.2 × 1''.3 :  $S = 5.5 \pm 1.2$  Jy), and NVSS at 1.4 GHz.

**G5.2–2.6**

**RA:** 18<sup>h</sup>07<sup>m</sup>30<sup>s</sup>  
**Dec:** −25°45'

**1-GHz flux/Jy:** 2.6?  
**Spectral index:** 0.6?

**Size/arcmin:** 18  
**Type:** S

**Radio:** Poorly resolved shell.

**References:**

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

<b>G5.4–1.2</b>		<b>Milne 56</b>
<b>RA:</b> 18 <sup>h</sup> 02 <sup>m</sup> 10 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 35?	<b>Size/arcmin:</b> 35
<b>Dec:</b> −24°54'	<b>Spectral index:</b> 0.2?	<b>Type:</b> C?
Part been called G5.3–1.0. Has been suggested that this is not a SNR.		
<b>Radio:</b> Incomplete shell, including wide ‘v’ of emission to east with small flat-spectrum source at apex.		
<b>Optical:</b> Detected.		
<b>X-ray:</b> Pulsar detected, with faint extension.		
<b>Distance:</b> HI absorption suggests >4.3 kpc.		
<b>Point sources:</b> Pulsar associated with flat spectrum source.		
<b>References:</b>		
Clark <i>et al.</i> 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3': S=38 Jy).		
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').		
Dickey & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).		
Angerhofer <i>et al.</i> 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 : S=21.9±2.4 Jy).		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Zealey <i>et al.</i> 1979, A&AS, 38, 39. Optical detection.		
Becker & Helfand 1985, Natur, 313, 115. VLA at 1.4 and 5 GHz. Suggest it is not SNR, see Helfand & Becker 1985, Natur, 313, 118.		
Manchester <i>et al.</i> 1985, MNRAS, 212, 975. Pulsar detection.		
Caswell <i>et al.</i> 1987, MNRAS, 225, 329. MOST at 843 MHz (42'' × 110'').		
Frail & Kulkarni 1991, Natur, 352, 785. Pulsar and remnant association.		
Manchester <i>et al.</i> 1991, MNRAS, 253, 7P. Pulsar and remnant association.		
Milne <i>et al.</i> 1992, MNRAS, 255, 707. Parkes 64-m at 4.75 (4'.5 : S=30.8±2.1 Jy) and 8.4 GHz (3' : S=24±3 Jy), including polarization.		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.0 × 4'.9 : S=38 Jy).		
Frail <i>et al.</i> 1994, AJ, 107, 1120. VLA at 327 MHz (68'' × 73''), plus HI absorption.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Gaensler & Frail 2000, Natur, 406, 158. Pulsar observations, including proper motion.		
Kaspi <i>et al.</i> 2001, ApJ, 562, L163. X-ray detection of pulsar, and upper limit for remnant.		

<b>G5.9+3.1</b>		<b>Size/arcmin:</b> 20
<b>RA:</b> 17 <sup>h</sup> 47 <sup>m</sup> 20 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 3.3?	<b>Type:</b> S
<b>Dec:</b> −22°16'	<b>Spectral index:</b> 0.4?	
<b>Radio:</b> Asymmetric shell.		

**References:**  
Reich *et al.* 1988, in SNRISM, p293. Effelsberg 100-m at 2.7 GHz (4'.3).

<b>G6.1+1.2</b>		<b>Size/arcmin:</b> 30 × 26
<b>RA:</b> 17 <sup>h</sup> 54 <sup>m</sup> 55 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 4.0?	<b>Type:</b> F
<b>Dec:</b> −23°05'	<b>Spectral index:</b> 0.3?	
Has been called G6.1+1.15.		

**Radio:** Faint, diffuse emission.

**References:**  
Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

G6.4–0.1	W28
<b>RA:</b> 18 <sup>h</sup> 00 <sup>m</sup> 30 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 310
<b>Dec:</b> −23°26'	<b>Spectral index:</b> varies
Has been called G6.6–0.2.	
<b>Radio:</b> Several non-thermal sources in a ring, with flat spectrum core.	
<b>Optical:</b> Filaments.	
<b>X-ray:</b> Diffuse emission from most of the remnant.	
<b>Point sources:</b> Young pulsar near edge of remnant.	
<b>Distance:</b> Uncertain mean optical velocity indicates 3.5–4 kpc.	
<b>References:</b>	
Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10 GHz (3').	
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical.	
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).	
Dopita <i>et al.</i> 1977, ApJ, 214, 179. Some optical line ratios.	
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).	
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.	
Andrews <i>et al.</i> 1983, ApJ, 266, 684. VLA at 1.4 GHz (8'') 4.9 GHz (3'') and 15 GHz (1'') of central region only, plus Einstein image of central region.	
Bohigas <i>et al.</i> 1983, RMxAA, 8, 155. Optical spectra.	
Odenwald <i>et al.</i> 1984, ApJ, 279, 162. Nearby IR source.	
Andrews <i>et al.</i> 1985, AJ, 90, 310. VLA of central component.	
Long <i>et al.</i> 1991, ApJ, 373, 567. Einstein and optical observations.	
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.1 × 4'.4 : S = 660 Jy).	
Kaspi <i>et al.</i> 1993, ApJ, 409, L57. Pulsar association.	
Frail <i>et al.</i> 1993, Natur, 365, 136. VLA at 327 MHz (smoothed to 65''), plus pulsar association.	
Frail <i>et al.</i> 1994, ApJ, 424, L111. VLA of associated OH masers.	
Esposito <i>et al.</i> 1996, ApJ, 461, 820. Possible associated γ-ray emission.	
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH maser emission.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	
Claussen <i>et al.</i> 1997, ApJ, 489, 143. VLA of OH masers.	
Claussen <i>et al.</i> 1999, ApJ, 522, 349. High resolution observations of OH masers.	
Yusef-Zadeh <i>et al.</i> 2000, ApJ, 540, 842. VLA at 327 MHz (2'.6 × 5'.5) and 1.48 GHz (40'' × 65'').	
Arikawa <i>et al.</i> 1999, PASJ, 51, L7. Observations of shocked CO.	
Dubner <i>et al.</i> 2000, AJ, 120, 1933. VLA at 328 MHz (97'' × 52'' : S = 425 ± 40 Jy) and 1415 MHz (88'' × 48'' : S = 246 ± 20 Jy), and comparison with other observations.	
Reach & Rho 2000, ApJ, 544, 843. ISO observations on interactions with surroundings.	
Rowell <i>et al.</i> 2000, A&A, 359, 337. Upper limit on high energy γ-rays.	
Douvion <i>et al.</i> 2001, A&A, 373, 281. ISO observations.	
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.	

**G6.4+4.0**

<b>RA:</b> 17 <sup>h</sup> 45 <sup>m</sup> 10 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 1.3?	<b>Size/arcmin:</b> 31
<b>Dec:</b> −21°22'	<b>Spectral index:</b> 0.4?	<b>Type:</b> S

**Radio:** Faint asymmetric shell.**References:**Reich *et al.* 1988, in SNRISM, p293. Effelsberg 100-m at 2.7 GHz (4'.3).**G7.0–0.1**

<b>RA:</b> 18 <sup>h</sup> 01 <sup>m</sup> 50 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 2.5?	<b>Size/arcmin:</b> 15
<b>Dec:</b> −22°54'	<b>Spectral index:</b> 0.5?	<b>Type:</b> S

Has been called G7.06–0.12.

**Radio:** Double rim, brightest in W, confused by bright HII region M20 in SE.**References:**Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz (2'.6 × 5'.5) and 1.48 GHz (40'' × 65'').  
Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz (97'' × 52'') and 1415 MHz (88'' × 48'').

**G7.7–3.7** 1814–24

**RA:** 18<sup>h</sup>17<sup>m</sup>25<sup>s</sup>  
**Dec:** −24°04'

**1-GHz flux/Jy:** 11  
**Spectral index:** 0.32

**Size/arcmin:** 22  
**Type:** S

**Radio:** Shell, with high polarization.

**References:**

- Dickel & Milne 1976, *AuJPh*, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).  
Milne *et al.* 1986, *MNRAS*, 223, 487. MOST at 843 MHz (44'' × 108'') and Parkes 64-m at 8.4 GHz (3':  $S=4.6 \pm 0.5$  Jy), with polarization, plus review of flux densities.  
Dubner *et al.* 1996, *AJ*, 111, 1304. VLA at 1.4 GHz (70'' × 35''):  $S=9.9 \pm 0.1$  Jy), including polarization.

**G8.7–5.0**

**RA:** 18<sup>h</sup>24<sup>m</sup>10<sup>s</sup>  
**Dec:** −23°48'

**1-GHz flux/Jy:** 4.4  
**Spectral index:** 0.3

**Size/arcmin:** 26  
**Type:** S

**Radio:** Asymmetric shell.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
Reich *et al.* 1990, *A&AS*, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

**G8.7–0.1**

(W30)

**RA:** 18<sup>h</sup>05<sup>m</sup>30<sup>s</sup>  
**Dec:** −21°26'

**1-GHz flux/Jy:** 80  
**Spectral index:** 0.5

**Size/arcmin:** 45  
**Type:** S?

**Radio:** Clumpy non-thermal shell, with low-frequency turnover.

**X-ray:** Northern edge detected.

**Point sources:** Pulsar inside western edge.

**References:**

- Odegard 1986, *AJ*, 92, 1372. TPT at 57.5 MHz (7'.2 × 9'.7):  $S=190 \pm 50$  Jy.  
Kassim & Weiler 1990, *Natur*, 343, 146. VLA at 327 MHz (3'.0 × 3'.7).  
Kassim & Weiler 1990, *ApJ*, 360, 184. VLA at 327 MHz (2'.8 × 4'.1):  $S=129 \pm 11$  Jy), and part at 1.4 GHz (0'.9 × 1'.8), plus review of flux densities.  
Frail *et al.* 1994, *AJ*, 107, 1120. VLA at 327 MHz (37'' × 55'').  
Finley & Ögelman 1994, *ApJ*, 434, L25. ROSAT observations, including pulsar.

**G9.8+0.6**

**RA:** 18<sup>h</sup>05<sup>m</sup>08<sup>s</sup>  
**Dec:** −20°14'

**1-GHz flux/Jy:** 3.9  
**Spectral index:** 0.5

**Size/arcmin:** 12  
**Type:** S

**Radio:** Asymmetric shell.

**References:**

- Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
Caswell 1983, *MNRAS*, 204, 833. Molonglo at 408 MHz (3':  $S=5.8 \pm 0.6$  Jy).  
Dubner *et al.* 1993, *AJ*, 105, 2251. VLA at 1.47 GHz (41'' × 63''):  $S=3.5 \pm 0.4$  Jy).  
Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

**G10.0–0.3**

**RA:** 18<sup>h</sup>08<sup>m</sup>39<sup>s</sup>  
**Dec:** −20°25'

**1-GHz flux/Jy:** 2.9  
**Spectral index:** 0.8

**Size/arcmin:** 8?  
**Type:** ?

The identification as a SNR has been questioned.

**Radio:** Not well resolved.

**X-ray:** X-ray pulsar, and extended emission from remnant.

**Point sources:** X-ray pulsar associated with soft gamma repeater.

**Distance:** Association with W31 gives 14.5 kpc.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
Kulkarni & Frail 1993, Natur, 365, 33. VLA at 330 MHz (smoothed to 2') and 1.49 GHz (40'' × 65''), including association with soft gamma repeater.  
Sonobe *et al.* 1994, ApJ, 436, L23. X-ray observations.  
Kulkarni *et al.* 1994, Natur, 368, 129. VLA at 1.4 GHz (31'') and 8.4 GHz (5''.5)  
Vasisht *et al.* 1995, ApJ, 440, L65. Radio monitoring of core.  
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.  
Corbel *et al.* 1997, ApJ, 478, 624. CO observations of surroundings.  
Frail *et al.* 1997, ApJ, 480, L129. VLA monitoring of changing core.  
Kouveliotou *et al.* 1998, Natur, 393, 235. X-ray pulsar detection.  
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.  
Gaensler *et al.* 2001, ApJ, 559, 963. Questioning of remnant identification.

**G11.2–0.3**

**RA:** 18<sup>h</sup>11<sup>m</sup>27<sup>s</sup>  
**Dec:** −19°25'

**1-GHz flux/Jy:** 22  
**Spectral index:** 0.50

**Size/arcmin:** 4  
**Type:** C

Probably associated with the SN of AD386.

**Radio:** Symmetrical clumpy shell, possibly with flatter spectrum core.

**X-ray:** Shell, with hard spectrum centrally brightened region around pulsar.

**Point sources:** Central pulsar.

**Distance:** HI absorption indicates 5 kpc.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
Radhakrishnan *et al.* 1972, ApJS, 24, 49. HI absorption.  
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').  
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20'' × 25'') and Einstein observations, with review of flux densities.  
Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz, plus HI absorption, Einstein observations.  
Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26''.5 : S = 4.04 ± 0.24 Jy).  
Green *et al.* 1988, MNRAS, 231, 735. VLA at 1.4 and 5 GHz.  
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.2 × 4'.1 : S = 39 Jy).  
Reynolds *et al.* 1994, MNRAS, 271, L1. ROSAT image and spectra.  
Vasisht *et al.* 1996, ApJ, 456, L59. ASCA observations.  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
Torii *et al.* 1997, ApJ, 489, L145. ASCA detection of pulsar.  
Torii *et al.* 1999, ApJ, 523, L69. X-ray timing observations of pulsar.  
Kothes & Reich 2001, A&A, 372, 627. Effelsberg 100-m at 4.25, 10.45, 14.9 and 32 GHz (2'.5, 1'.1, 0'.86, 0'.45 : S = 9.6 ± 0.5, 6.3 ± 0.4, 5.7 ± 0.4, 3.8 ± 0.4).  
Kaspi *et al.* 2001, ApJ, 560, 371. Chandra observations.

**G11.4–0.1**

**RA:** 18<sup>h</sup>10<sup>m</sup>47<sup>s</sup>  
**Dec:** −19°05'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.5

**Size/arcmin:** 8  
**Type:** S?

**Radio:** Incomplete shell, possibly with central core.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' :  $S=9.4$  Jy) contaminated by sidelobes of a nearby source, and Parkes 64-m at 5 GHz (4' :  $S=2.8$  Jy).  
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 :  $S=2.0 \pm 0.4$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.2 × 4'.1 :  $S=18$  Jy).  
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (44'' × 63'' :  $S=5.1 \pm 0.6$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G12.0–0.1**

**RA:** 18<sup>h</sup>12<sup>m</sup>11<sup>s</sup>  
**Dec:** −18°37'

**1-GHz flux/Jy:** 3.5  
**Spectral index:** 0.7

**Size/arcmin:** 7?  
**Type:** ?

**Radio:** Incomplete shell, defined in E only.

**X-ray:** Detected.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' :  $S=6.6$  Jy) and Parkes 64-m at 5 GHz (4' :  $S=1.1$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.2 × 4'.1).  
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41'' × 61'' :  $S=0.7$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G13.3–1.3**

**RA:** 18<sup>h</sup>19<sup>m</sup>20<sup>s</sup>  
**Dec:** −18°00'

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:** 70 × 40  
**Type:** S?

**Radio:** Amorphous emission.

**X-ray:** Elongated emission.

**Optical:** Filaments in S.

**Distance:** Absorption indicates 2–4 kpc.

**References:**

- Seward *et al.* 1995, ApJ, 449, 681. ROSAT detection, optical studies and observations of CO.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G13.5+0.2**

**RA:** 18<sup>h</sup>14<sup>m</sup>14<sup>s</sup>  
**Dec:** −17°12'

**1-GHz flux/Jy:** 3.5?  
**Spectral index:** 1.0?

**Size/arcmin:** 5 × 4  
**Type:** S

Has been called G13.46+0.16.

**Radio:** Elongated, incomplete shell.

**References:**

- Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ( $S=0.65 \pm 0.05$  Jy) and 1.4 GHz (15'' :  $S=2.67 \pm 0.5$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G15.1–1.6**

**RA:** 18<sup>h</sup>24<sup>m</sup>00<sup>s</sup>  
**Dec:** −16°34'

**1-GHz flux/Jy:** 5.5?  
**Spectral index:** 0.8?

**Size/arcmin:** 30 × 24  
**Type:** S

**Radio:** Elongated, incomplete shell.

**References:**

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

**G15.9+0.2**

**RA:** 18<sup>h</sup>18<sup>m</sup>52<sup>s</sup>  
**Dec:** −15°02'

**1-GHz flux/Jy:** 5  
**Spectral index:** 0.6?

**Size/arcmin:** 7 × 5  
**Type:** S?

**Radio:** Incomplete shell, with bright concentration to the E.

**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=7.7$  Jy) and Parkes 64-m at 5 GHz (4': $S=1.9$  Jy).  
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8: $S=1.1\pm0.2$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz (58'' × 44'').  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (77'' × 61'': $S=11.2\pm1.0$  Jy), 1.4 GHz (23'' × 14'': $S=3.9\pm0.1$  Jy) and 4.9 GHz (13'' × 16'').  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G16.2–2.7**

**RA:** 18<sup>h</sup>28<sup>m</sup>50<sup>s</sup>  
**Dec:** −16°11'

**1-GHz flux/Jy:** 2  
**Spectral index:** 0.5

**Size/arcmin:** 17  
**Type:** S

**Radio:** Double rim.

**References:**

Trushkin 1999, A&A, 352, L103. Review of radio observations.

**G16.7+0.1**

**RA:** 18<sup>h</sup>20<sup>m</sup>56<sup>s</sup>  
**Dec:** −14°20'

**1-GHz flux/Jy:** 3.0  
**Spectral index:** 0.6

**Size/arcmin:** 4  
**Type:** C

Has been called G16.73+0.08.

**Radio:** Asymmetric shell with flat-spectrum core.

**X-ray:** Detected.

**References:**

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz (5: $S=0.95$  Jy) and 1.4 GHz (15'': $S=2.43$  Jy) and Ooty at 327 MHz ( $S=5.13$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G16.8–1.1**

**RA:** 18<sup>h</sup>25<sup>m</sup>20<sup>s</sup>  
**Dec:** −14°46'

**1-GHz flux/Jy:** 2?  
**Spectral index:** ?

**Size/arcmin:** 30 × 24?  
**Type:** ?

Has been called G16.85–1.05.

**Radio:** Overlapping thermal and non-thermal emission, parameters uncertain.

**Point sources:** Pulsar within boundary of non-thermal emission.

**References:**

Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'.4), plus other observations.

**G17.4–2.3**

**RA:** 18<sup>h</sup>30<sup>m</sup>55<sup>s</sup>  
**Dec:** −14°52'

**1-GHz flux/Jy:** 4.8?  
**Spectral index:** 0.8?

**Size/arcmin:** 24?  
**Type:** S

**Radio:** Incomplete, poorly defined shell.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G17.8–2.6**

**RA:** 18<sup>h</sup>32<sup>m</sup>50<sup>s</sup>  
**Dec:** −14°39'

**1-GHz flux/Jy:** 4.0?  
**Spectral index:** 0.3?

**Size/arcmin:** 24  
**Type:** S

**Radio:** Well defined shell.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G18.8+0.3**

**RA:** 18<sup>h</sup>23<sup>m</sup>58<sup>s</sup>  
**Dec:** −12°23'

**1-GHz flux/Jy:** 33  
**Spectral index:** 0.4

**Size/arcmin:** 17 × 11  
**Type:** S

Kes 67

Has been called G18.9+0.3.

**Radio:** Incomplete shell, in complex region near the HII region W39.

**Distance:** HI absorption indicates > 9.5 kpc, and possibly < 19 kpc.

**References:**

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5': S = 17 ± 7 Jy).  
 Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3': S = 38 Jy).  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Milne *et al.* 1989, PASAu, 8, 187. Parkes 64-m at 8.4 GHz (3'.0: S = 12.9 ± 1.0 Jy), including polarization.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (2'.9 × 3'.5: S = 55 Jy).  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (75" × 55": S = 29.9 ± 0.3 Jy).  
 Dubner *et al.* 1999, AJ, 118, 930. Parkes 64-m at 1.6 GHz (15") for HI, VLA at 1.6 GHz (17" × 12") for OH, plus CO observations.

**G18.9–1.1**

**RA:** 18<sup>h</sup>29<sup>m</sup>50<sup>s</sup>  
**Dec:** −12°58'

**1-GHz flux/Jy:** 37  
**Spectral index:** varies

**Size/arcmin:** 33  
**Type:** C?

Has been called G18.95–1.1 and G18.94–1.04.

**Radio:** Non-thermal, diffuse partially limb-brightened, with central ridge.

**X-ray:** Partial shell.

**References:**

- Fürst *et al.* 1985, Natur, 314, 720. Effelsberg 100-m at 4.75 GHz (2'.4: S = 23.8 Jy), plus other flux densities.  
 Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7'.2 × 8': S = 82 ± 15 Jy), plus review of flux densities.  
 Barnes & Turtle 1988, in SNRISM, p347. Molonglo at 408 MHz (2'.9 × 3'.1: S = 58 ± 9 Jy) and Parkes 64-m at 5 GHz (4'.4 × 4'.1: S = 23 ± 6 Jy).  
 Patnaik *et al.* 1988, Natur, 332, 136. Ooty at 327 MHz (0'.6 × 1'.6).  
 Fürst *et al.* 1989, A&A, 209, 361. Effelsberg 100-m at 4.75 GHz (2'.45: S = 23.8 Jy) and VLA at 1.5 and 4.9 GHz (19" × 14"), and Effelsberg 100-m at 1.4 GHz (9') for HI.  
 Aschenbach *et al.* 1991, A&A, 246, L32. ROSAT observations.  
 Fürst *et al.* 1997, A&A, 319, 655. ROSAT observations, and Effelsberg 100-m at 10.6 GHz (1'.1).

**G20.0–0.2**

**RA:** 18<sup>h</sup>28<sup>m</sup>07<sup>s</sup>  
**Dec:** −11°35'

**1-GHz flux/Jy:** 10  
**Spectral index:** 0.0

**Size/arcmin:** 10  
**Type:** F

**Radio:** Faint, filled-centre, polarized.

**Point sources:** OH source 20.1–0.1 is nearby.

**References:**

Becker & Helfand 1985, ApJ, 297, L25. VLA at 1.4 and 5 GHz (12'').  
 Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz ( $7'.2 \times 8' : S = 8.5 \pm 2$  Jy), plus review of flux densities.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

**G21.5–0.9**

**RA:** 18<sup>h</sup>33<sup>m</sup>33<sup>s</sup>  
**Dec:** −10°35'

**1-GHz flux/Jy:** 6?  
**Spectral index:** 0.0

**Size/arcmin:** 4  
**Type:** C

Early observations relate to the central core only.

**Radio:** Filled-centre, with high frequency turnover.

**X-ray:** Central core, with extended, faint halo.

**Distance:** HI absorption indicates 5.5 kpc or more.

**References:**

Wilson & Weiler 1976, A&A, 53, 89. WSRT at 5 GHz (6''  $\times$  35'').  
 Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20''  $\times$  5'') and 8 GHz (7''  $\times$  2''), plus review of flux densities.  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Becker & Szymkowiak 1981, ApJ, 248, L23. VLA at 5 GHz (8''), and Einstein observations.  
 Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT X-ray spectrum, and VLA HI observations.  
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (26''.5 :  $S = 5.64 \pm 0.29$  Jy).  
 Fürst *et al.* 1988, PASJ, 40, 347. NRO array at 22.3 GHz (4''.4  $\times$  7''.3).  
 Salter *et al.* 1989, A&A, 225, 167. Observations at 90.7 (29''.5 :  $S = 3.8 \pm 0.4$  Jy) and 141.9 GHz ( $S = 2.5 \pm 1.2$  Jy).  
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ( $S = 3.94 \pm 0.70$  Jy), plus review of flux densities.  
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3  $\times$  3'.4 :  $S = 9$  Jy).  
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Slane *et al.* 2000, ApJ, 533, L29. Chandra observations identifying X-ray halo.  
 Warwick *et al.* 2001, A&A, 365, L248. XMM-Newton observations of X-ray halo.  
 Bock *et al.* 2001, ApJ, 561, L203. BIMA at 94 GHz (8''.6  $\times$  4''.6).  
 Safi-Harb *et al.* 2001, ApJ, 561, 308. Chandra and other X-ray observations.

**G21.8–0.6**

Kes 69

**RA:** 18<sup>h</sup>32<sup>m</sup>45<sup>s</sup>  
**Dec:** −10°08'

**1-GHz flux/Jy:** 69  
**Spectral index:** 0.5

**Size/arcmin:** 20  
**Type:** S

**Radio:** Incomplete shell.

**X-ray:** Detected.

**Distance:** H<sub>2</sub>CO absorption indicates > 6.3 kpc.

**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Wilson 1972, A&A, 19, 354. H<sub>2</sub>CO absorption.  
 Kundu *et al.* 1974, AJ, 79, 1332. NRAO 140-ft at 5 GHz (6') and 10 GHz (3').  
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft et 2.7 GHz (5' :  $S = 42.3 \pm 4.6$  Jy).  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Seward 1990, ApJS, 73, 781. Einstein observations.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3  $\times$  3'.5 :  $S = 132$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.

**G22.7–0.2**

**RA:** 18<sup>h</sup>33<sup>m</sup>15<sup>s</sup>  
**Dec:** −09°13'

**1-GHz flux/Jy:** 33  
**Spectral index:** 0.6

**Size/arcmin:** 26  
**Type:** S?

**Radio:** Non-thermal ring in complex region, overlapping G23.3–0.3.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3 × 3'.4 :  $S=82$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G23.3–0.3**

**RA:** 18<sup>h</sup>34<sup>m</sup>45<sup>s</sup>  
**Dec:** −08°48'

**1-GHz flux/Jy:** 70  
**Spectral index:** 0.5

**W41**  
**Size/arcmin:** 27  
**Type:** S

**Radio:** Incomplete ring, in complex region, overlapping G22.7–0.2.

**Point sources:** Pulsar association suggested.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3 × 3'.4 :  $S=138$  Jy).  
 Gaensler & Johnston 1995, MNRAS, 275, L73. Possible pulsar association.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G23.6+0.3**

**RA:** 18<sup>h</sup>33<sup>m</sup>03<sup>s</sup>  
**Dec:** −08°13'

**1-GHz flux/Jy:** 8?  
**Spectral index:** 0.3

**Size/arcmin:** 10?  
**Type:** ?

**Radio:** Not well resolved, in complex region.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G24.7–0.6**

**RA:** 18<sup>h</sup>38<sup>m</sup>43<sup>s</sup>  
**Dec:** −07°32'

**1-GHz flux/Jy:** 8  
**Spectral index:** 0.5

**Size/arcmin:** 15?  
**Type:** S?

**Radio:** Incomplete shell, defined in SW.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' :  $S=12.3$  Jy) and Parkes 64-m at 5 GHz (4' :  $S=3.6$  Jy).  
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 :  $S=2.2 \pm 0.5$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (44'' × 56'' :  $S=1.9$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

**G24.7+0.6**

**RA:** 18<sup>h</sup>34<sup>m</sup>10<sup>s</sup>  
**Dec:** −07°05'

**1-GHz flux/Jy:** 20?  
**Spectral index:** 0.2?

**Size/arcmin:** 30 × 15  
**Type:** C?

**Radio:** Filled-centre, with faint shell, and a compact HII region to the S.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2').  
 Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4.3 :  $S = 19 \pm 3$  Jy) and 4.75 GHz (2'.4 :  $S = 17 \pm 4$  Jy) and NRO 45-m at 10.2 GHz (2'.7 :  $S = 15 \pm 3$  Jy).  
 Becker & Helfand 1987, ApJ, 316, 660. VLA at 1.4 GHz (12''), and X-ray upper limit.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

**G27.4+0.0**

**RA:** 18<sup>h</sup>41<sup>m</sup>19<sup>s</sup>  
**Dec:** −04°56'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.68

**Size/arcmin:** 4  
**Type:** S

4C−04.71

Early references refer to G27.3−0.1 (Kes 73), a supposed larger remnant.

**Radio:** Incomplete shell.

**X-ray:** Diffuse emission, with central low period pulsar.

**Point sources:** Central AXp.

**Distance:** HI absorption indicates 6 to 7.5 kpc.

**References:**

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5').  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).  
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 :  $S = 2.0 \pm 0.5$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3'.5 :  $S = 4.4$  Jy).  
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz (60'' × 45'' :  $S = 3.5$  Jy). Revise  $S_{408\text{ MHz}} = 10.4$  Jy, and  $S_{5\text{ GHz}} = 1.9 \pm 0.2$  Jy.  
 Kriss *et al.* 1985, ApJ, 288, 703. Einstein observations, plus VLA at 1.4 and 5 GHz (12'').  
 Sanbonmatsu & Helfand 1992, AJ, 104, 2189. VLA at 1.4 GHz for HI absorption.  
 Helfand *et al.* 1994, ApJ, 434, 627. ROSAT observations, particularly of central source.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including nearby masers.  
 Vasish & Gotthelf 1997, ApJ, 486, L129. ASCA detection of pulsar.  
 Gotthelf & Vasish 1997, ApJ, 486, L133. ASCA observations.  
 Gotthelf *et al.* 1999, ApJ, 522, L49. X-ray timing observations of pulsar.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.  
 Mereghetti *et al.* 2001, MNRAS, 321, 143. Search for optical/IR counterpart to pulsar.

**G27.8+0.6**

**RA:** 18<sup>h</sup>39<sup>m</sup>50<sup>s</sup>  
**Dec:** −04°24'

**1-GHz flux/Jy:** 30  
**Spectral index:** varies

**Size/arcmin:** 50 × 30  
**Type:** F

**Radio:** Filled-centre, with spectral turnover.

**References:**

- Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4'.3 :  $S = 23 \pm 2$  Jy), and 4.75 GHz (2'.4 :  $S = 18 \pm 2$  Jy) and NRO 45-m at 10.2 GHz (smoothed to 4'.3 :  $S = 8.5 \pm 2$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

**G28.6–0.1**

**RA:** 18<sup>h</sup>43<sup>m</sup>55<sup>s</sup>  
**Dec:** −03°53'

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:** 13 × 9  
**Type:** S

**Radio:** Poorly defined regions of non-thermal emission.

**X-ray:** Diffuse shell.

**References:**

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 1.4 GHz (15'') and 5 GHz (15'').  
Bamba *et al.* 2001, PASJ, 53, L21. ASCA observations.

**G28.8+1.5**

**RA:** 18<sup>h</sup>39<sup>m</sup>00<sup>s</sup>  
**Dec:** −02°55'

**1-GHz flux/Jy:** ?  
**Spectral index:** 0.4?

**Size/arcmin:** 100?  
**Type:** S?

**Radio:** Part of rim detected.

**X-ray:** Diffuse, Centrally brightened.

**References:**

Schwentker 1994, A&A, 286, L47. ROSAT observations.  
Song *et al.* 2000, PASJ, 52, 181. ASCA observations.

**G29.6+0.1**

**RA:** 18<sup>h</sup>44<sup>m</sup>52<sup>s</sup>  
**Dec:** −02°57'

**1-GHz flux/Jy:** 1.5?  
**Spectral index:** 0.5?

**Size/arcmin:** 5  
**Type:** S

**Radio:** Diffuse shell.

**Point sources:** AXP associated.

**References:**

Gaensler *et al.* 1999, ApJ, 526, L37. VLA at 5 GHz (13'') and 8 GHz (8'').  
Vasisht *et al.* 2000, ApJ, 542, L49. X-ray observations of AXP.

**G29.7–0.3**

Kes 75

**RA:** 18<sup>h</sup>46<sup>m</sup>25<sup>s</sup>  
**Dec:** −02°59'

**1-GHz flux/Jy:** 10  
**Spectral index:** 0.7

**Size/arcmin:** 3  
**Type:** C

**Radio:** Shell with flatter spectrum emission from centre.

**X-ray:** Thermal shell and non-thermal core, and central pulsar.

**Point sources:** X-ray pulsar.

**Distance:** HI absorption indicates > 9 kpc and possibly at 21 kpc.

**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (8'.4:S=5±20% Jy).  
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').  
Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20'' × 7'') and 8 GHz (25'' × 8''), plus review of flux densities.  
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
Becker *et al.* 1983, ApJ, 268, L93. VLA at 1.4 GHz (3''), plus Einstein observations.  
Becker & Helfand 1984, ApJ, 283, 154. VLA at 5 GHz (2''.6), plus HI.  
Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26''.5:S=1.02±0.07 Jy).  
Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.  
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.4×3'.7:S=27.4 Jy).  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Blanton & Helfand 1996, ApJ, 470, 961. ASCA observations.  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
Gotthelf *et al.* 2000, ApJ, 542, L37. X-ray pulsar detection.  
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G30.7–2.0**

**RA:** 18<sup>h</sup>54<sup>m</sup>25<sup>s</sup>  
**Dec:** −02°54'

**1-GHz flux/Jy:** 0.5?  
**Spectral index:** 0.7?

**Size/arcmin:** 16  
**Type:** ?

**Radio:** Poorly defined.

**References:**

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G30.7+1.0**

**RA:** 18<sup>h</sup>44<sup>m</sup>00<sup>s</sup>  
**Dec:** −01°32'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.4

**Size/arcmin:** 24 × 18  
**Type:** S?

**Radio:** Non-thermal, highly polarized part shell?

**Point sources:** Compact source near centre.

**References:**

Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'.4 :  $S = 3.4 \pm 0.4$  Jy), plus other flux densities.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.6 × 3'.9 :  $S = 8.6$  Jy).  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

**G31.5–0.6**

**RA:** 18<sup>h</sup>51<sup>m</sup>10<sup>s</sup>  
**Dec:** −01°31'

**1-GHz flux/Jy:** 2?  
**Spectral index:** ?

**Size/arcmin:** 18?  
**Type:** S?

Has been called G31.55–0.65.

**Radio:** Distorted shell? near HII region.

**Optical:** Diffuse, incomplete shell.

**References:**

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4), plus other flux densities.  
 Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

<b>G31.9+0.0</b>		<b>3C391</b>
<b>RA:</b> 18 <sup>h</sup> 49 <sup>m</sup> 25 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 24	<b>Size/arcmin:</b> 7 × 5
<b>Dec:</b> −00°55'	<b>Spectral index:</b> 0.55	<b>Type:</b> S

**Radio:** Shell, brightest in NW.

**X-ray:** Diffuse with central core.

**Distance:** HI absorption is seen to the tangent point (8.5 kpc).

**References:**

- Radhakrishnan *et al.* 1972, ApJS, 24, 49. HI absorption.  
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').  
 Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15 GHz (58'':  $S > 1.6$ ).  
 Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20'' × 23'') and 8 GHz (24'' × 9''), plus review of flux densities.  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Goss *et al.* 1979, A&A, 78, 75. FIRST at 1.4 GHz (54'' × 66'':  $S = 20 \pm 2$  Jy) and Effelsberg 100-m at 10.7 GHz (77'':  $S = 7.5 \pm 0.8$  Jy).  
 Wang & Seward 1984, ApJ, 279, 705. Einstein observations.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.4 × 3'.7:  $S = 44.8$  Jy).  
 Reynolds & Moffett 1993, AJ, 105, 2226. VLA at 1.4 GHz (6''), including possible associated CO.  
 Moffett & Reynolds 1994, ApJ, 425, 668. VLA at 330 MHz (smoothed to 30'':  $S = 38.5 \pm 0.5$  Jy) 1.46 GHz (6''.7) and 4.85 GHz (6''.2 × 6''.4), including spectral index and polarization studies.  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
 Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.  
 Rho & Petre 1996, ApJ, 467, 698. ROSAT observations.  
 Wilner *et al.* 1998, AJ, 115, 247. CO observations of surroundings.  
 Reach & Rho 1998, ApJ, 507, L93. ISO observations.  
 Reach & Rho 1999, ApJ, 511, 836. CO, HCO<sup>+</sup> and CS observations of surroundings.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey survey observations.

<b>G32.0−4.9</b>		<b>3C396.1</b>
<b>RA:</b> 19 <sup>h</sup> 06 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 22?	<b>Size/arcmin:</b> 60?
<b>Dec:</b> −03°00'	<b>Spectral index:</b> 0.5?	<b>Type:</b> S?

**Radio:** Possible large shell?

**References:**

- Milne & Hill 1969, AuJPh, 22, 211. Parkes 64-m at 635 MHz (31'':  $S = 25 \pm 30\%$  Jy), 1410 MHz (15'':  $S = 19 \pm 15\%$  Jy) and 2650 MHz (8'.4:  $S = 8.6 \pm 30$  Jy). Fluxes if size is 60', plus review of flux densities.  
 Caswell 1970, AuJPh, 23, 105. Revision of low frequency flux densities.  
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 111 MHz ( $S = 105 \pm 30$  Jy).

<b>G32.1−0.9</b>		
<b>RA:</b> 18 <sup>h</sup> 53 <sup>m</sup> 10 <sup>s</sup>	<b>1-GHz flux/Jy:</b> ?	<b>Size/arcmin:</b> 40?
<b>Dec:</b> −01°08'	<b>Spectral index:</b> ?	<b>Type:</b> C?

**Radio:** Possible faint shell, not well defined.

**X-ray:** Diffuse, with clumps.

**References:**

- Folgheraiter *et al.* 1997, MNRAS, 292, 365. ROSAT and ASCA observations.

<b>G32.8–0.1</b>		Kes 78
<b>RA:</b> 18 <sup>h</sup> 51 <sup>m</sup> 25 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 11?	<b>Size/arcmin:</b> 17
<b>Dec:</b> −00°08'	<b>Spectral index:</b> 0.2?	<b>Type:</b> S?

Part has been called G33.1–0.1.

**Radio:** Elongated shell?

**References:**

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 7.2 \pm 0.5$  Jy).  
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 430 MHz ( $S = 19.0 \pm 15.5$  Jy).  
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz ( $3'$ ).  
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 12.8$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 7.7$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ( $2'.6$ ).  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ( $3'.6 \times 3'.8 : S = 31.3$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.

<b>G33.2–0.6</b>		
<b>RA:</b> 18 <sup>h</sup> 53 <sup>m</sup> 50 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 3.5	<b>Size/arcmin:</b> 18
<b>Dec:</b> −00°02'	<b>Spectral index:</b> varies	<b>Type:</b> S

**Radio:** Incomplete shell.

**References:**

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ( $2'.6$ ).  
 Reich 1982, A&A, 106, 314. Effelsberg 100-m at 2.7 GHz ( $4'.4 : S = 2.6 \pm 0.3$  Jy) and 4.75 GHz ( $2'.5 : S = 1.75 \pm 0.2$  Jy).  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ( $68'' \times 52'' : S = 2.7 \pm 0.3$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

<b>G33.6+0.1</b>		Kes 79, 4C00.70, HC13
<b>RA:</b> 18 <sup>h</sup> 52 <sup>m</sup> 48 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 22	<b>Size/arcmin:</b> 10
<b>Dec:</b> +00°41'	<b>Spectral index:</b> 0.5	<b>Type:</b> S

Has been called G33.7+0.0.

**Radio:** Shell, with bright central region, in complex region.

**X-ray:** Detected.

**Distance:** HI absorption gives about 10 kpc.

**References:**

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 35.5$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 7.8$  Jy).  
 Caswell *et al.* 1975, A&A, 45, 239. HI absorption.  
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo  $S_{430\text{MHz}} = 69 \pm 33$  Jy.  
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz ( $3' : S = 6.8 \pm 1.5$  Jy).  
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ( $6'.8 : S = 11.4 \pm 1.1$  Jy).  
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz ( $1'$ ), plus observations of the nearby point source.  
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source, possibly extragalactic.  
 Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source, plus Einstein observations.  
 Green 1989, MNRAS, 238, 737. OH absorption.  
 Frail & Clifton 1989, ApJ, 336, 854. VLA at 1.4 GHz ( $1' \times 2'.9$ ), including HI absorption.  
 Velusamy *et al.* 1991, AJ, 102, 676. VLA at 327 MHz ( $1'$ ),  $1.5(7'' \times 14'')$  and 5 GHz ( $7''$ ), including spectral comparison.  
 Green & Dewdney 1992, MNRAS, 254, 686. Observations of adjacent molecular material.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ( $3'.6 \times 3'.8 : S = 34.8$  Jy).  
 Seward & Velusamy 1995, ApJ, 439, 715. ROSAT observations.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey survey observations.

**G34.7–0.4**

**RA:** 18<sup>h</sup>56<sup>m</sup>00<sup>s</sup>  
**Dec:** +01°22'

**1-GHz flux/Jy:** 230  
**Spectral index:** 0.30

**W44, 3C392**  
**Size/arcmin:** 35 × 27  
**Type:** C

Has been called G34.6–0.5.

**Radio:** Distorted shell, brighter to the E, with pulsar and associated nebula.

**X-ray:** Centrally concentrated, thermal spectrum, plus pulsar nebula.

**Point sources:** Pulsar within the boundary of the remnant.

**Distance:** HI absorption indicates 2.5 kpc.

**References:**

- Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10.7 GHz ( $3' : S = 105 \pm 7$  Jy).  
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 177 \pm 10$  Jy).  
 Caswell *et al.* 1975, A&A, 45, 239. HI absorption.  
 Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ( $3' : S = 299$  Jy which is less than the 390 Jy found by Kesteven 1968, AuJPh, 21, 739 with the same telescope).  
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo  $S_{430\text{ MHz}} = 540 \pm 187$  Jy.  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ( $8'.4$ ) and 5 GHz ( $4'.4$ ).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ( $2'.6$ ).  
 Smith *et al.* 1985, MNRAS, 217, 99. Einstein observations.  
 Wolszczan *et al.* 1991, ApJ, 372, L99. Pulsar detection.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ( $3'.6 \times 3'.8 : S = 469$  Jy).  
 Jones *et al.* 1993, MNRAS, 265, 631. VLA at 1.4 GHz ( $15''$ ), plus X-ray spectra.  
 Rho *et al.* 1994, ApJ, 430, 757. Optical and ROSAT observations.  
 Koo & Heiles 1995, ApJ, 442, 679. HI of surrounding shell.  
 Esposito *et al.* 1996, ApJ, 461, 820. Possible associated  $\gamma$ -ray emission.  
 Harrus *et al.* 1996, ApJ, 464, L161. ASCA observations.  
 Frail *et al.* 1996, ApJ, 464, L165. VLA at 1.5 and 8.4 GHz ( $8''.9 \times 7''.8$ ) of pulsar nebula.  
 Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
 Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.  
 Harrus *et al.* 1997, ApJ, 488, 781. X-ray observations.  
 Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.  
 Giacani *et al.* 1997, AJ, 113, 1379. VLA at 1.4 GHz ( $15''$ ), plus optical images.  
 Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.  
 Cox *et al.* 1999, ApJ, 524, 179. Revision of distance.  
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.

**G36.6–0.7**

**RA:** 19<sup>h</sup>00<sup>m</sup>35<sup>s</sup>  
**Dec:** +02°56'

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:** 25?  
**Type:** S?

**Radio:** Polarized arc, possibly part of a larger shell?

**References:**

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz ( $2'.4$ ), plus other flux densities.  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ( $3'.4 \times 3'.7 : S = 6.7$  Jy).  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

**G36.6+2.6**

**RA:** 18<sup>h</sup>48<sup>m</sup>49<sup>s</sup>  
**Dec:** +04°26'

**1-GHz flux/Jy:** 0.7?  
**Spectral index:** 0.5?

**Size/arcmin:** 17 × 13?  
**Type:** S

**Radio:** Poorly resolved shell.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ( $4'.3$ ).

G39.2–0.3	3C396, HC24, NRAO 593
<b>RA:</b> 19 <sup>h</sup> 04 <sup>m</sup> 08 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 18
<b>Dec:</b> +05°28'	<b>Spectral index:</b> 0.6
<b>Radio:</b> Shell, brightest to the W, with faint ‘tail’ to E.	<b>Size/arcmin:</b> 8 × 6
<b>X-ray:</b> Centrally brightened.	<b>Type:</b> S
<b>Distance:</b> HI absorption suggests at least 7.7 kpc.	
<b>References:</b>	
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3').	
Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430 \text{ MHz}} = 54 \pm 38 \text{ Jy}$ .	
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3' : $S = 4.1 \pm 1.0 \text{ Jy}$ ).	
Caswell <i>et al.</i> 1975, A&A, 45, 239. HI absorption.	
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).	
Caswell <i>et al.</i> 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (65'' × 48'').	
Becker & Helfand 1987, AJ, 94, 1629. VLA at 1.4 GHz (12' : $S = 14 \text{ Jy}$ ) and 5 GHz, plus Einstein observations.	
Patnaik <i>et al.</i> 1990, A&A, 232, 467. VLA at 1.5 GHz (25'') and 1.4 GHz (7''.8 × 7''.5) and 5 GHz (25'') including polarization, plus Ooty at 327 MHz (100'' × 31''), including review of flux densities.	
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.5 × 3'.6 : $S = 42.5 \text{ Jy}$ ).	
Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz for spectral index studies.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	
Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.	
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.	
Koralesky <i>et al.</i> 1998, AJ, 116, 1323. VLA search for OH emission.	
Harrus & Slane 1999, ApJ, 516, 811. ASCA observations.	
Aharonian <i>et al.</i> 2001, A&A, 375, 1008. Limit on high energy $\gamma$ -rays.	

G39.7–2.0	W50, SS433
<b>RA:</b> 19 <sup>h</sup> 12 <sup>m</sup> 20 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 85?
<b>Dec:</b> +04°55'	<b>Spectral index:</b> 0.7?
Eastern part has been called G40.0–3.1. Is this a SNR?	<b>Size/arcmin:</b> 120 × 60
<b>Radio:</b> Elongated shell, containing SS433, adjacent to the HII region S74.	<b>Type:</b> ?
<b>Optical:</b> Faint filaments at the edge of the radio emission.	
<b>X-ray:</b> Emission from SS433 and two lobes.	
<b>Point sources:</b> SS433 is the compact source in the centre of the W50.	
<b>Distance:</b> Distance to SS433 is 5 kpc.	
<b>References:</b>	
van den Bergh 1980, ApJ, 236, L23. Optical in H $\alpha$ and [SII].	
Zealey <i>et al.</i> 1980, MNRAS, 192, 731. Optical spectra.	
van Gorkom <i>et al.</i> 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source (not SS433).	
Sequaist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.	
Watson <i>et al.</i> 1983, ApJ, 273, 688. X-ray observations.	
Downes <i>et al.</i> 1986, MNRAS, 218, 393. Effelsberg 100-m at 4.75 GHz (2'.4 : $S = 34 \pm 4 \text{ Jy}$ ), plus previous 1.7 and 2.7 GHz data.	
Romney <i>et al.</i> 1987, ApJ, 321, 822. VLBI of SS433, including distance.	
Elston & Baum 1987, AJ, 94, 1633. Mosaic with VLA at 1.4 GHz (30'') of fine structure only.	
Kawai <i>et al.</i> 1989, PASJ, 41, 491. X-ray observations of SS433.	
Band 1989, ApJ, 336, 937. Einstein and EXOSAT observations.	
Yamauchi <i>et al.</i> 1994, PASJ, 46, L109. X-ray spectral observations.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	
Brinkmann <i>et al.</i> 1996, A&A, 312, 306. ROSAT observations.	
Kotami <i>et al.</i> 1996, PASJ, 48, 619. X-ray line emission from SS433.	
Safi-Harb & Ögelman 1997, ApJ, 483, 868. ROSAT observations.	
Dubner <i>et al.</i> 1998, AJ, 116, 1842. VLA at 328 MHz (64'' × 60' : $S = 160 \pm 20 \text{ Jy}$ ), and 1.4 GHz (56'' × 54''), plus NRAO 140-ft at 1.4 GHz (21') for HI observations.	
Safi-Harb & Petre 1999, ApJ, 512, 784. X-ray observations.	
Aharonian <i>et al.</i> 2001, A&A, 375, 1008. Limit on high energy $\gamma$ -rays.	

**G40.5–0.5**

**RA:** 19<sup>h</sup>07<sup>m</sup>10<sup>s</sup>  
**Dec:** +06°31'

**1-GHz flux/Jy:** 11  
**Spectral index:** 0.5

**Size/arcmin:** 22  
**Type:** S

**Radio:** Shell, brightest to the NE.

**References:**

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Downes *et al.* 1980, A&A, 92, 47. Effelsberg 100-m at 1.7 GHz (7'.6 :  $S = 9.3 \pm 1.3$  Jy), and 2.7 GHz (4'.4 :  $S = 7.2 \pm 0.5$  Jy), plus review of flux densities.  
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy  $\gamma$ -rays.

**G41.1–0.3**

**RA:** 19<sup>h</sup>07<sup>m</sup>34<sup>s</sup>  
**Dec:** +07°08'

**1-GHz flux/Jy:** 22  
**Spectral index:** 0.48

**Size/arcmin:** 4.5  $\times$  2.5  
**Type:** S

3C397

**Radio:** 3C397 is two sources: the E is the SNR, the W is a HII region.

**X-ray:** Brighter to the E and W, with possible central component.

**Distance:** Possible limit of  $> 7.5$  kpc for non-thermal component from HI absorption.

**References:**

- Kundu *et al.* 1974, AJ, 79, 1332. NRAO 140-ft at 10 GHz (3') and 5 GHz (6').  
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5' :  $S = 21.3 \pm 1.2$  Jy for both components).  
 Caswell *et al.* 1975, A&A, 45, 239. HI absorption.  
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo  $S_{430 \text{ MHz}} = 82 \pm 51$  Jy, also Algonquin 46-m at 10.6 GHz (3' :  $S = 12 \pm 2$  Jy), and Haystack 36-m at 15.5 GHz (2'.3 :  $S = 8.5 \pm 3.0$  Jy).  
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' :  $S = 29.8$  Jy) and Parkes 64-m at 5 GHz (4' :  $S = 8.7$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (58''  $\times$  52'').  
 Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz (8''), plus Einstein observations.  
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30' :  $S = 1.10 \pm 0.19$  Jy).  
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.5  $\times$  3'.6 :  $S = 46.3$  Jy).  
 Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz, for spectral index studies.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Chen *et al.* 1999, ApJ, 520, 737. ASCA and ROSAT observations.  
 Dyer & Reynolds 1999, ApJ, 526, 365. VLA at 1.5 GHz (6''.9  $\times$  6''.6) and 4.8 GHz (46''.4  $\times$  5''.6), including polarization and comparison with ROSAT image.  
 Safi-Harb *et al.* 2000, ApJ, 545, 922. ROSAT, ASCA and other X-ray observations.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey survey observations.  
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy  $\gamma$ -rays.

**G42.8+0.6**

**RA:** 19<sup>h</sup>07<sup>m</sup>20<sup>s</sup>  
**Dec:** +09°05'

**1-GHz flux/Jy:** 3?  
**Spectral index:** 0.5?

**Size/arcmin:** 24  
**Type:** S

Has been called G42.8+0.65.

**Radio:** Faint shell.

**Point sources:** Near soft gamma repeater, and young pulsar.

**References:**

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4 :  $S = 1.5 \pm 0.2$  Jy), plus other flux densities.  
 Vasish特 *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3'.2  $\times$  3'.4).  
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.  
 Lorimer & Xilouris 2000, ApJ, 545, 385. Pulsar detection.  
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy  $\gamma$ -rays.

**G43.3–0.2**

W49B

**RA:** 19<sup>h</sup>11<sup>m</sup>08<sup>s</sup>  
**Dec:** +09°06'

**1-GHz flux/Jy:** 38  
**Spectral index:** 0.48

**Size/arcmin:** 4 × 3  
**Type:** S

**Radio:** Shell, brightest to the SE and W, near the HII region W49A.

**X-ray:** Filled-centre.

**Distance:** HI absorption indicates 12.5 to 14 kpc.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 2.7 GHz (4').  
 Downes & Wilson 1974, A&A, 34, 133. Effelsberg 100-m at 10.7 GHz (1'.3).  
 Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz (58'' :  $S = 9.0 \pm 0.7$  Jy).  
 Lockhart & Goss 1978, A&A, 67, 355. HI absorption.  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Pye *et al.* 1984, MNRAS, 207, 649. Einstein observations, and VLA at 1.4 and 5 GHz (both 12'').  
 Smith *et al.* 1985, ApJ, 296, 469. EXOSAT spectrum.  
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30'' :  $S = 6.90 \pm 0.38$  Jy).  
 Moffett & Reynolds 1994, ApJ, 437, 705. VLA at 330 MHz (6''.7 × 7''.7 :  $S = 64.4$  Jy), 1.48 GHz (4''.8 × 5''.2 :  $S = 31.8$  Jy) and 4.85 GHz (4''.0 × 4''.1), including polarization.  
 Fujimoto *et al.* 1995, PASJ, 47, L31. ASCA observations.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Hwang *et al.* 2000, ApJ, 532, 970. ROSAT image and ASCA spectroscopy.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey survey observations.  
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy  $\gamma$ -rays.  
 Brogan & Troland 2001, ApJ, 550, 799. VLA at 1.4 GHz (27'' × 24'' and 5'') for HI Zeeman splitting.  
 Lacey *et al.* 2001, ApJ, 559, 954. VLA at 74 MHz (26'' × 23'' :  $S = 55.6$  Jy) and 326 MHz (6''.6 × 6''.2 :  $S = 56.0$  Jy).

**G43.9+1.6**

**RA:** 19<sup>h</sup>05<sup>m</sup>50<sup>s</sup>  
**Dec:** +10°30'

**1-GHz flux/Jy:** 8.6?  
**Spectral index:** 0.2?

**Size/arcmin:** 60?  
**Type:** S?

**Radio:** Large, poorly defined faint shell.

**Point sources:** Soft gamma repeater nearby.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
 Vasish特 *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3'.2 × 3'.4).  
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.

**G45.7–0.4**

**RA:** 19<sup>h</sup>16<sup>m</sup>25<sup>s</sup>  
**Dec:** +11°09'

**1-GHz flux/Jy:** 4.2?  
**Spectral index:** 0.4?

**Size/arcmin:** 22  
**Type:** S

**Radio:** Shell, brightest to the SE, poorly defined to NW.

**References:**

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4 :  $S = 2.6 \pm 0.3$  Jy), plus other flux densities.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

**G46.8–0.3**

(HC30)

**RA:** 19<sup>h</sup>18<sup>m</sup>10<sup>s</sup>**1-GHz flux/Jy:** 14**Dec:** +12°09'**Spectral index:** 0.5**Size/arcmin:** 17 × 13**Type:** S

Has been called G46.6–0.2.

**Radio:** Shell, two bright arcs to NNW and SSE.

**Distance:** HI absorption suggests 6.8–8.8 kpc.

**References:**

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ( $5' : S = 9.8 \pm 0.9$  Jy), and VRO 37-m at 1.7 GHz ( $S = 14.5 \pm 5.5$  Jy).

Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 20.3$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 7.1$  Jy).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S<sub>430 MHz</sub> = 46 ± 21 Jy.

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ( $6'.8 : S = 5.9 \pm 0.6$  Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ( $2'.6$ ).

Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ( $58'' \times 53'' : S = 13.3 \pm 0.1$  Jy).

**G49.2–0.7**

(W51)

**RA:** 19<sup>h</sup>23<sup>m</sup>50<sup>s</sup>**1-GHz flux/Jy:** 160?**Dec:** +14°06'**Spectral index:** 0.3?**Size/arcmin:** 30**Type:** S?

**Radio:** In complex region, parameters uncertain.

**X-ray:** Elongated east–west.

**Optical:** Some diffuse emission possibly associated.

**Distance:** HI absorption suggests 4.1 kpc.

**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Parkes 64-m at 5 GHz ( $4'$ ).

Sato 1973, PASJ, 25, 135. HI absorption.

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5'$ ),  $S = 51.5 \pm 3.2$  Jy, for the non-thermal component, but probably confused.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ( $2'.6$ ).

Seward 1990, ApJS, 73, 781. Einstein observations.

Copetti & Schmidt 1991, MNRAS, 250, 127. CLFST at 151 MHz.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz ( $3'.1 \times 3'.5$ ).

Subrahmanyam & Goss 1995, MNRAS, 275, 755. VLA at 330 MHz ( $1'.1$ ).

Koo *et al.* 1995, ApJ, 447, 211. ROSAT observations.

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Green *et al.* 1997, AJ, 114, 2058. OH masers.

Koo & Moon 1997, ApJ, 475, 194. Arecibo ( $3'$ ) and VLA ( $40'' \times 42''$ ) at 1.4 GHz for HI.

Koo & Moon 1997, ApJ, 485, 263. NRAO 12-m CO and HCO<sup>+</sup> observations.

Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

**G53.6–2.2**

**RA:** 19<sup>h</sup>38<sup>m</sup>50<sup>s</sup>  
**Dec:** +17°14'

**1-GHz flux/Jy:** 8  
**Spectral index:** 0.75

3C400.2, NRAO 611

**Size/arcmin:** 33 × 28  
**Type:** S

Has been called G53.7–2.2.

**Radio:** Ring of emission, with extension to NW.

**Optical:** Filaments and diffuse emission.

**X-ray:** Centrally brightened, offset to NW.

**Distance:** Mean optical velocity indicates 6.7 kpc, HI absorption indicates 2.3 kpc.

**References:**

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ( $5' : S = 5.3 \pm 0.6$  Jy).  
van den Bergh 1978, ApJS, 38, 119. Optical.  
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 4.8 \pm 0.3$  Jy).  
Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ( $3' : S = 11.7$  Jy).  
Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo  $S_{430 \text{ MHz}} = 20 \pm 10$  Jy,  $S_{318 \text{ MHz}} = 20 \pm 3.6$  Jy.  
Goss *et al.* 1975, A&A, 43, 459. WSRT at 610 MHz ( $1' \times 3' : S = 13.2 \pm 1.6$  Jy).  
Sabaddin & d'Odorico 1976, A&A, 49, 119. Optical spectra.  
Rosado 1983, RMxAA, 8, 59. Optical spectra.  
Blair & Long 1988, PASP, 100, 461. Optical imaging and spectroscopy.  
Long *et al.* 1991, ApJ, 373, 567. Einstein and optical observations.  
Winkler *et al.* 1993, ApJ, 405, 608. Optical imaging.  
Dubner *et al.* 1994, AJ, 108, 207. VLA at 327 MHz (59'') and 1.49 GHz (52''), plus X-rays.  
Saken *et al.* 1995, ApJ, 443, 231. ROSAT observations.  
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
Giacani *et al.* 1998, A&AS, 133, 61. DRAO at 1.4 GHz for HI studies, including distance.  
Yoshita *et al.* 2001, PASJ, 53, 93. ASCA observations, and spectral comparison with ROSAT.

**G54.1+0.3**

**RA:** 19<sup>h</sup>30<sup>m</sup>31<sup>s</sup>  
**Dec:** +18°52'

**1-GHz flux/Jy:** 0.5  
**Spectral index:** 0.1

**Size/arcmin:** 1.5  
**Type:** F?

**Radio:** Filled-centre.

**X-ray:** Centrally concentrated, with extension to NE.

**References:**

- Green 1985, MNRAS, 216, 691. 5km at 2.7 GHz (7'' × 20'').  
Reich *et al.* 1985, A&A, 151, L10. Effelsberg 100-m at 4.75 GHz (2'.4 :  $S = 0.37 \pm 0.04$  Jy)  
Velusamy & Becker 1988, AJ, 95, 1162. VLA at 1.4 (14'':  $S = 0.48 \pm 0.03$  Jy), 1.6 (14'':  $S = 0.42 \pm 0.03$  Jy) and 5 GHz (5'':  $S = 0.33 \pm 0.02$  Jy), Ooty at 327 MHz ( $S = 0.50 \pm 0.08$  Jy), plus review of flux densities.  
Seward 1989, AJ, 97, 481. Einstein observations.  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
Lu *et al.* 2001, A&A, 370, 570. ROSAT and ASCA observations.

**G54.4–0.3**

(HC40)

**RA:** 19<sup>h</sup>33<sup>m</sup>20<sup>s</sup>  
**Dec:** +18°56'

**1-GHz flux/Jy:** 28  
**Spectral index:** 0.5

**Size/arcmin:** 40  
**Type:** S

Has been called G54.5–0.3.

**Radio:** Shell, in complex region.

**References:**

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 34.4 \pm 5.0$  Jy).  
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
Caswell 1985, AJ, 90, 1224. DRAO at 1.4 GHz (1'.3 × 2'.6 :  $S = 18 \pm 4$  Jy).  
Velusamy *et al.* 1986, JApA, 7, 105. WSRT at 609 MHz (50'' × 191'' smoothed to 100'' × 200'').  
Junkes *et al.* 1992, A&AS, 96, 1. Surrounding CO.  
Junkes *et al.* 1992, A&A, 261, 289. Nearby IRAS sources.

**G55.0+0.3**

**RA:** 19<sup>h</sup>32<sup>m</sup>00<sup>s</sup>  
**Dec:** +19°50'

**1-GHz flux/Jy:** 0.5?  
**Spectral index:** 0.5?

**Size/arcmin:** 20 × 15?  
**Type:** S

Has been called G55.2+0.5.

**Radio:** Faint, partial shell.

**Distance:** Association with HI features implies 14 kpc.

**Point sources:** Old pulsar nearby.

**References:**

Matthews *et al.* 1998, ApJ, 493, 312. WSRT at 327 MHz ( $1'.0 \times 2'.9 : S = 0.98 \pm 0.15$  Jy), DRAO at 1.4 GHz ( $1'.0 \times 2'.9 : S = 0.25 \pm 0.12$  Jy), plus HI observations.

**G55.7+3.4**

**RA:** 19<sup>h</sup>21<sup>m</sup>20<sup>s</sup>  
**Dec:** +21°44'

**1-GHz flux/Jy:** 1.4  
**Spectral index:** 0.6

**Size/arcmin:** 23  
**Type:** S

**Radio:** Incomplete shell.

**Point sources:** Old pulsar within the boundary of the remnant.

**References:**

Goss *et al.* 1977, A&A, 61, 93. WSRT observations at 610 MHz ( $57'' \times 156'' : S = 1.9 \pm 0.2$  Jy) and 1415 MHz ( $27'' \times 72'' : S = 1.0 \pm 0.1$  Jy).

**G57.2+0.8**

(4C21.53)

**RA:** 19<sup>h</sup>34<sup>m</sup>59<sup>s</sup>  
**Dec:** +21°57'

**1-GHz flux/Jy:** 1.8?  
**Spectral index:** ?

**Size/arcmin:** 12?  
**Type:** S?

**Radio:** Extended non-thermal arc.

**Point sources:** Near the millisecond pulsar, but not thought to be related.

**References:**

Sieber & Seiradakis 1984, A&A, 130, 257. Effelsberg 100-m at 1.4 GHz ( $8'.8 : 1.34 \pm 0.1$ ), 2.7 GHz ( $4'.3 : 0.86 \pm 0.1$ ), plus other surveys of the area.

Caswell *et al.* 1985, AJ, 90, 488. DRAO at 1.4 GHz ( $1' \times 3'$ ).

**G59.5+0.1**

**RA:** 19<sup>h</sup>42<sup>m</sup>33<sup>s</sup>  
**Dec:** +23°35'

**1-GHz flux/Jy:** 3?  
**Spectral index:** ?

**Size/arcmin:** 5  
**Type:** S

Has been called G59.6+0.1.

**Radio:** Incomplete shell.

**References:**

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz ( $1'.0 \times 2'.5 : S = 5.1 \pm 0.2$  Jy), and northern sky survey at 4.9 GHz.  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G59.8+1.2**

**RA:** 19<sup>h</sup>38<sup>m</sup>55<sup>s</sup>  
**Dec:** +24°19'

**1-GHz flux/Jy:** 1.6  
**Spectral index:** 0.5

**Size/arcmin:** 20 × 16?  
**Type:** ?

Has been called G59.7+1.2.

**Radio:** Poorly defined source.

**References:**

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ( $4'.3$ ).

**G63.7+1.1**

**RA:** 19<sup>h</sup>47<sup>m</sup>52<sup>s</sup>  
**Dec:** +27°45'

**1-GHz flux/Jy:** 1.8  
**Spectral index:** 0.3

**Size/arcmin:** 8  
**Type:** F

**Radio:** Centrally brightened, with core.

**References:**

Wallace *et al.* 1997, AJ, 114, 2068. WSRT at 1.4 GHz (14'' × 26'' :  $S = 1.63$  Jy), DRAO at 1.4 GHz (smoothed to 2'), plus review of flux densities and other observations.

**G65.1+0.6**

**RA:** 19<sup>h</sup>54<sup>m</sup>40<sup>s</sup>  
**Dec:** +28°35'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.6

**Size/arcmin:** 90 × 50  
**Type:** S

**Radio:** Large, faint shell.

**Point sources:** Pulsar nearby.

**References:**

Landecker *et al.* 1990, A&A, 232, 207. DRAO at 408 MHz (3'.5 × 7'.0 :  $S = 9.5 \pm 0.1$  Jy), and 1.4 GHz (1'.0 × 2'.0 :  $S = 5.4 \pm 1.0$  Jy).  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G65.3+5.7**

**RA:** 19<sup>h</sup>33<sup>m</sup>00<sup>s</sup>  
**Dec:** +31°10'

**1-GHz flux/Jy:** 52?  
**Spectral index:** 0.6?

**Size/arcmin:** 310 × 240  
**Type:** S?

Has been called G65.2+5.7.

**Radio:** Large, faint ring? near S91 and S94.

**Optical:** Strong in Hα+[NII] and [OIII].

**X-ray:** Detected.

**Distance:** Mean optical velocity suggests 0.8 kpc.

**References:**

Gull *et al.* 1977, ApJ, 215, L69. Optical plates.  
Reich *et al.* 1979, A&A, 72, 270. Effelsberg 100-m observations at 1.42 GHz (smoothed to 11' :  $S = 42.4 \pm 1.6$  Jy), estimate  $S_{408\text{ MHz}} = 91 \pm 5$  Jy from previous sky survey.  
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.  
Rosado 1981, ApJ, 250, 222. Optical interferometry.  
Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.  
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.  
Seward 1990, ApJS, 73, 781. Einstein observations.  
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

**G65.7+1.2**

DA 495

**RA:** 19<sup>h</sup>52<sup>m</sup>10<sup>s</sup>  
**Dec:** +29°26'

**1-GHz flux/Jy:** 5.1  
**Spectral index:** 0.6

**Size/arcmin:** 18  
**Type:** ?

Has mistakenly been called G55.7+1.2.

**Radio:** Filled-centre or thick shell?

**References:**

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' :  $S = 2.8 \pm 0.4$  Jy), and VRO 37-m at 1.7 GHz ( $S = 4.4 \pm 0.5$  Jy), plus review of flux densities.  
Dickey & DeNoyer 1975, AJ, 80, 437. Arecibo  $S_{430\text{ MHz}} = 8.7 \pm 4.9$  Jy,  $S_{318\text{ MHz}} = 9.7 \pm 2.2$  Jy.  
Landecker & Caswell 1983, AJ, 88, 1810. DRAO at 1.4 GHz (0'.9 × 1'.5 :  $S = 4.4 \pm 0.2$  Jy).  
Velusamy *et al.* 1989, JApA, 10, 161. Ooty at 327 MHz (36'' × 64''), WSRT at 610 MHz (62'') and VLA at 1.4 GHz (36'' and 12''), including IRAS imaging.  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G67.7+1.8**

**RA:** 19<sup>h</sup>54<sup>m</sup>32<sup>s</sup>  
**Dec:** +31°29'

**1-GHz flux/Jy:** 1.4  
**Spectral index:** 0.3

**Size/arcmin:** 9  
**Type:** S

**Radio:** Double arc shell.

**Optical:** Filaments in N.

**References:**

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz ( $1'.9 \times 1'.0 : S = 1.9 \pm 0.1$  Jy), and northern sky survey at 4.9 GHz ( $S = 0.42 \pm 0.05$  Jy).  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

**G68.6–1.2**

**RA:** 20<sup>h</sup>08<sup>m</sup>40<sup>s</sup>  
**Dec:** +30°37'

**1-GHz flux/Jy:** 0.7?  
**Spectral index:** 0.0?

**Size/arcmin:** 28 × 25?  
**Type:** ?

**Radio:** Faint, poorly defined source.

**References:**

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G69.0+2.7**

**RA:** 19<sup>h</sup>53<sup>m</sup>20<sup>s</sup>  
**Dec:** +32°55'

**1-GHz flux/Jy:** 120?  
**Spectral index:** varies

**CTB 80**  
**Size/arcmin:** 80?  
**Type:** ?

An association with a SN in AD1408 has been suggested. Has been called G68.8+2.8. Is it a SNR?

**Radio:** Compact core, flat spectrum plateau, and steeper spectrum extensions, with spectral break?

**Optical:** Expanding nebulosity near centre, with filaments to the SW and far NE.

**X-ray:** Diffuse emission with compact source.

**Point sources:** Pulsar at western edge of core.

**References:**

Angerhofer *et al.* 1981, A&A, 94, 313. WSRT at 610 MHz ( $56'' \times 103''$ ) 1.4 GHz ( $24'' \times 44''$ ) and 5 GHz ( $7'' \times 13''$ ), plus optical.  
Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.  
Sofue *et al.* 1983, PASJ, 35, 437. NRO 45-m at 10.2 GHz (2'.7).  
Velusamy & Kundu 1983, JApA, 3, 253. VLA of compact sources.  
Blair *et al.* 1984, ApJ, 282, 161. Optical images and spectra.  
Wang & Seward 1984, ApJ, 285, 607. Einstein observations.  
Strom *et al.* 1984, A&A, 139, 43. Radio observations of flat spectrum component, VLA 5 GHz (1''.7) and 1.4 GHz.  
Mantovani *et al.* 1985, A&A, 145, 50. Bologna at 408 MHz ( $2'.6 \times 4'.9 : S = 67.5 \pm 10.5$  Jy), Effelsberg 100-m at 1.41 (9':  $S = 62 \pm 9$  Jy), 1.72 (7'.6:  $S = 66 \pm 5$  Jy), 2.7 (4'.5:  $S = 52 \pm 4$  Jy) and 4.75 GHz (2'.4:  $S = 44 \pm 3.3$  Jy), plus review of flux densities.  
Kulkarni *et al.* 1988, Natur, 331, 50. Pulsar detection.  
Angelini *et al.* 1988, ApJ, 330, L43. EXOSAT spectra.  
Fesen *et al.* 1988, Natur, 334, 229. IRAS of surrounding shell.  
Whitehead *et al.* 1989, MNRAS, 237, 1109. Optical of core.  
Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.  
Hester & Kulkarni 1989, ApJ, 340, 362. Optical imaging and spectroscopy.  
Koo *et al.* 1990, ApJ, 364, 178. Large, expanding HI shell.  
Greidanus & Strom 1990, A&A, 240, 376. Optical kinematics of core.  
Koo *et al.* 1993, ApJ, 417, 196. VLA at 1.4 GHz of surrounding HI (1'), plus IRAS.  
Safi-Harb *et al.* 1995, ApJ, 439, 722. ROSAT observations.  
Srinivasan 1997, ApJ, 489, 170.  $\gamma$ -ray observations of pulsar.  
Mavromatakis *et al.* 2001, A&A, 371, 300. Optical observations.

**G69.7+1.0**

**RA:** 20<sup>h</sup>02<sup>m</sup>40<sup>s</sup>  
**Dec:** +32°43'

**1-GHz flux/Jy:** 1.6  
**Spectral index:** 0.8

**Size/arcmin:** 16  
**Type:** S

**Radio:** Poorly resolved source.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G73.9+0.9**

**RA:** 20<sup>h</sup>14<sup>m</sup>15<sup>s</sup>  
**Dec:** +36°12'

**1-GHz flux/Jy:** 9?  
**Spectral index:** 0.3?

**Size/arcmin:** 22?  
**Type:** S?

**Radio:** Diffuse, centrally brightened to SW.

**Optical:** Weak shell.

**References:**

- Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'.4 :  $S = 6.7 \pm 0.5$  Jy), plus other flux densities.  
 Chastenay & Pineault 1988, in SNRISM, p297. DRAO at 408 MHz (3'.5  $\times$  5'.9) and 1.4 GHz (1'.0  $\times$  1'.7).  
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz (3'.4  $\times$  5'.8 :  $S = 12.7 \pm 1.2$  Jy) and 1.4 GHz (1'.0  $\times$  1'.7 :  $S = 7.4 \pm 1.0$  Jy).  
 Lozinskaya *et al.* 1993, ARep, 37, 240. Optical observations.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
 Pineault *et al.* 1996, AJ, 112, 201. DRAO at 1.4 GHz (smoothed to 2') for H<sub>I</sub>.  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

<b>G74.0–8.5</b>		<b>Cygnus Loop</b>
<b>RA:</b> 20 <sup>h</sup> 51 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 210	<b>Size/arcmin:</b> 230 × 160
<b>Dec:</b> +30°40'	<b>Spectral index:</b> varies	<b>Type:</b> S
<b>Radio:</b> Shell, brightest to the NE, with fainter breakout region to S, with spectral variations.		
<b>Optical:</b> Large filamentary loop, brightest to the NE, not well defined to the S or W.		
<b>X-ray:</b> Shell in soft X-rays.		
<b>Point sources:</b> Several compact radio sources within the boundary of the remnant, including CL4, plus X-ray sources in S.		
<b>Distance:</b> Optical proper motion and shock velocity gives 0.44 kpc.		
<b>References:</b>		
Sequoia & Gilmore 1982, AJ, 87, 378. VLA observations of CL4.		
Green 1984, MNRAS, 211, 433. Most of remnant at 408 MHz (80'' × 160'').		
Charles <i>et al.</i> 1985, ApJ, 295, 456. Einstein observations.		
Braun & Strom 1986, A&A, 164, 208. IRAS observations.		
Straka <i>et al.</i> 1986, ApJ, 306, 266. Radio and optical comparison of NE. VLA at 1.6 GHz (4''.7) plus optical.		
Raymond <i>et al.</i> 1988, ApJ, 324, 869. Detailed optical study of filaments.		
Ballet <i>et al.</i> 1989, A&A, 211, 217. [FeX] and [FeXIV] spectroscopy.		
Ballet & Rothenflug 1989, A&A, 218, 277. EXOSAT observations.		
Green 1990, AJ, 100, 1927. DRAO at 408 MHz (3'.3 × 6'.7) for spectral index study, plus X-ray and optical.		
Graham <i>et al.</i> 1991, AJ, 101, 175. Shocked molecular H outside rim in NE.		
Shull & Clarke 1991, PASP, 103, 811. Optical spectroscopy of nearside filaments.		
Blair <i>et al.</i> 1991, ApJ, 374, 202. Far UV mapping.		
Blair <i>et al.</i> 1991, ApJ, 379, L33. UV spectroscopy.		
Gredanis & Strom 1992, A&A, 257, 265. Optical kinematics.		
Shull & Hippelain 1992, ApJ, 383, 714. Optical kinematics and proper motion.		
Fesen <i>et al.</i> 1992, AJ, 104, 719. H $\alpha$ imagery.		
Cornett <i>et al.</i> 1992, ApJ, 395, L9. UV imagery.		
Long <i>et al.</i> 1992, ApJ, 400, 214. Optical of Balmer dominated filament.		
Arendt <i>et al.</i> 1992, ApJ, 400, 562. IRAS observations.		
Vancura <i>et al.</i> 1993, ApJ, 417, 663. Far UV observations.		
Hester <i>et al.</i> 1994, ApJ, 420, 721. H $\alpha$ , [OIII] and other optical observations of Balmer dominated filaments in NE.		
Miyata <i>et al.</i> 1994, PASJ, 46, L101. X-ray observations of NE.		
Graham <i>et al.</i> 1995, ApJ, 444, 787. X-ray and optical studies of interaction with surroundings.		
Sauvageot & Decourchelle 1995, A&A, 296, 201. [FeX] and [FeXIV] observations.		
Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Levenson <i>et al.</i> 1996, ApJ, 468, 323. ROSAT and optical observations of W.		
Decourchelle <i>et al.</i> 1997, A&A, 326, 811. ROSAT observations of N.		
Levenson <i>et al.</i> 1997, ApJ, 484, 304. ROSAT observations.		
Leahy <i>et al.</i> 1997, AJ, 114, 2081. DRAO at 1.4 GHz (1' × 2'), including polarisation.		
Miyata <i>et al.</i> 1998, PASJ, 50, 257. ASCA observations.		
Miyata <i>et al.</i> 1998, PASJ, 50, 475. ASCA observations of compact X-ray source in S.		
Leahy & Roger 1998, ApJ, 505, 784. DRAO at 1.4 GHz (1'.0 × 1'.9) and 408 MHz (3'.4 × 6'.9), for spectral index studies in comparison with other radio observations.		
Levenson <i>et al.</i> 1998, ApJS, 118, 541. Optical images.		
Roger <i>et al.</i> 1999, A&AS, 137, 7. 22 MHz flux density ( $S = 1400 \pm 400$ Jy).		
Bohigas <i>et al.</i> 1999, ApJ, 518, 324. Optical spectroscopy of surroundings.		
Levenson <i>et al.</i> 1999, ApJ, 526, 874. ROSAT images.		
Blair <i>et al.</i> 1999, AJ, 118, 942. HST observations, for distance.		
Aschenbach & Leahy 1999, A&A, 341, 602. ROSAT image, and comparison with radio.		
Sauvageot <i>et al.</i> 1999, A&A, 351, 669. [Nev] observations.		
Szentygyorgyi <i>et al.</i> 2000, ApJ, 529, 279. [Nev] images.		
Danforth <i>et al.</i> 2000, AJ, 119, 2319. UV, optical and X-ray comparison of selected regions.		
Sankrit <i>et al.</i> 2000, AJ, 120, 1925. HST far-UV spectra on non-radiative shock.		
Miyata <i>et al.</i> 2001, ApJ, 550, 1023. ASCA observations of compact X-ray sources.		
Miyata & Tsunemi 2001, ApJ, 552, 624. ASCA spectroscopy of regions in N and E.		
Ghavamian <i>et al.</i> 2001, ApJ, 547, 995. Optical spectroscopy.		
Danforth <i>et al.</i> 2001, AJ, 122, 938. Far-UV spectroscopy, H $\alpha$ and other optical observations of NE region.		
Levenson & Graham 2001, ApJ, 559, 948. HST of SE region.		

**G74.9+1.2**

CTB 87

**RA:** 20<sup>h</sup>16<sup>m</sup>02<sup>s</sup>**1-GHz flux/Jy:** 9**Size/arcmin:** 8 × 6**Dec:** +37°12'**Spectral index:** varies**Type:** F**Radio:** Filled-centre, with high polarization and high frequency turnover.**X-ray:** Centrally brightened.**Distance:** HI absorption indicates 12 kpc.**Point sources:** Extragalactic compact source is nearby.**References:**

- Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo  $S_{430 \text{ MHz}} = 12.2 \pm 9.7 \text{ Jy}$ ,  $S_{318 \text{ MHz}} = 17.7 \pm 5.0 \text{ Jy}$ .  
 Weiler & Shaver 1978, A&A, 70, 389. WSRT at 610 MHz ( $57'' \times 94''$ :  $S = 9.1 \pm 1.2 \text{ Jy}$ ), 1.4 ( $24'' \times 40''$ :  $S = 8.7 \pm 1.2 \text{ Jy}$ ) and 5 GHz ( $24'' \times 40''$ :  $S = 5.6 \pm 1.3 \text{ Jy}$ ).  
 Geldzahler *et al.* 1980, A&A, 84, 237. Effelsberg 100-m at 2.7 GHz ( $4'.4$ :  $S = 7.6 \pm 0.5 \text{ Jy}$ ).  
 Wilson 1980, ApJ, 241, L19. Einstein observations.  
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption of nearby compact source.  
 Sequaist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.  
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (smoothed to  $40''$ :  $S = 1.47 \pm 0.19 \text{ Jy}$ ).  
 Green & Gull 1989, MNRAS, 237, 555. VLA at 1.4 GHz ( $1''.2 \times 1''.4$ ) including HI.  
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz, plus review of flux densities.  
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ( $3''.4 \times 5''.8$ :  $S = 11.6 \pm 0.4 \text{ Jy}$ ) and 1.4 GHz ( $1''.0 \times 1''.7$ :  $S = 7.2 \pm 0.3 \text{ Jy}$ ).  
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.  
 Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz ( $3''.5 \times 5''.2$ :  $S = 13.3 \pm 0.8 \text{ Jy}$ ) and Effelsberg 100-m at 4.8 GHz ( $S = 7.5 \pm 0.7 \text{ Jy}$ ).  
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.  
 Cho *et al.* 1994, AJ, 108, 634. CO of adjacent molecular clouds.  
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
 Wallace *et al.* 1997, A&A, 317, 212. DRAO at 408 MHz ( $3''.4 \times 5''.5$ ) and 1.4 GHz ( $1''.0 \times 1''.6$ ) including HI (smoothed to  $2''$ ).  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G76.9+1.0****Size/arcmin:** 12 × 9**RA:** 20<sup>h</sup>22<sup>m</sup>20<sup>s</sup>**1-GHz flux/Jy:** 2**Type:** ?**Dec:** +38°43'**Spectral index:** 0.6**Radio:** Diffuse, non-thermal, with low frequency turnover.**References:**

- Landecker *et al.* 1993, A&A, 276, 522. VLA at 1.49 GHz ( $14''$ ), 4.86 GHz ( $13'' \times 16''$ ) and 8.55 GHz ( $11'' \times 12''$ ), including polarization and review of flux densities.  
 Landecker *et al.* 1997, A&AS, 123, 199. Miyun at 232-MHz ( $3''.8 \times 5''.4$ ).  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G78.2+2.1**

**RA:** 20<sup>h</sup>20<sup>m</sup>50<sup>s</sup>  
**Dec:** +40°26'

**1-GHz flux/Jy:** 340  
**Spectral index:** 0.5

**DR4, γ Cygni SNR**  
**Size/arcmin:** 60  
**Type:** S

Has been called G78.1+1.8.

**Radio:** In complex region (early catalogues refer to other proposed remnants in this region).

**Optical:** Faint filaments, spectra indicate a SNR superposed on a HII region.

**X-ray:** Weak emission from the SE of the remnant.

**Point sources:** γ-ray and X-ray point source in remnant.

**References:**

- Higgs *et al.* 1977, AJ, 82, 718. DRAO at 1.4 GHz ( $2' \times 3'$ :  $S = 270 \pm 40$  Jy) plus some 10 GHz ( $4'$ ) survey data, reveals true extent of remnant.  
d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.  
van den Bergh 1978, ApJS, 38, 119. Optical.  
Landecker *et al.* 1980, A&AS, 39, 133. DRAO HI observations ( $2' \times 3'.1$ ).  
Higgs *et al.* 1983, AJ, 88, 97. CO of surroundings.  
Bohigas *et al.* 1983, RMxAA, 8, 155. Optical spectra, find thermal only.  
Braun & Strom 1986, A&AS, 63, 345. WSRT HI observations.  
Fukui & Tatematsu 1988, in SNRISM, p261. CO observations of the vicinity ( $2'.7$ ).  
Green 1989, MNRAS, 238, 737. OH observations.  
Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ( $3'.4 \times 5'.8$ :  $S = 480 \pm 60$  Jy) and 1.4 GHz ( $1'.0 \times 1'.7$ :  $S = 270 \pm 40$  Jy).  
Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz ( $3'.5 \times 5'.2$ :  $S = 540 \pm 40$  Jy) and Effelsberg 100-m at 4.8 GHz ( $S = 150 \pm 15$  Jy).  
Esposito *et al.* 1996, ApJ, 461, 820. Associated γ-ray emission.  
Brazier *et al.* 1996, MNRAS, 281, 1033. γ-ray and X-ray point source.  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Zhang *et al.* 1997, A&A, 324, 641. Multi-frequency radio comparison.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
Lozinskaya *et al.* 2000, AstL, 26, 77. Hα and X-ray observations.  
Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.

**G82.2+5.3**

**RA:** 20<sup>h</sup>19<sup>m</sup>00<sup>s</sup>  
**Dec:** +45°30'

**1-GHz flux/Jy:** 120?  
**Spectral index:** 0.5?

W63

**Size/arcmin:** 95 × 65  
**Type:** S

**Radio:** Shell in the Cygnus X complex.

**Optical:** In complex region, but spectra indicate SNR filaments.

**X-ray:** Detected.

**References:**

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5'$ :  $S = 59.0 \pm 3.5$  Jy).  
Sabbadin 1976, A&A, 51, 159. Optical spectra.  
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ( $6'.8$ ). Incomplete mapping.  
Rosado & González 1981, RMxAA, 5, 93. Optical spectra.  
Seward 1990, ApJS, 73, 781. Einstein observations.  
Higgs *et al.* 1991, JRASC, 85, 24. DRAO at 408 MHz ( $3'.4 \times 5'.0$ :  $S = 165 \pm 21$  Jy), plus review of flux densities.  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G84.2-0.8**

**RA:** 20<sup>h</sup>53<sup>m</sup>20<sup>s</sup>  
**Dec:** +43°27'

**1-GHz flux/Jy:** 11  
**Spectral index:** 0.5

**Size/arcmin:** 20 × 16  
**Type:** S

**Radio:** Elongated shell, with a filament aligned with the major axis.

**References:**

- Matthews *et al.* 1977, A&A, 55, 1. WSRT at 610 MHz ( $56'' \times 81''$ :  $S = 12.4 \pm 1.5$  Jy) and Effelsberg 100-m at 2.7 GHz ( $4'.4$ :  $S = 6.8 \pm 1.3$  Jy).  
Matthews & Shaver 1980, A&A, 87, 255. WSRT at 1415 MHz ( $23'' \times 32''$ ), and Effelsberg 100-m at 2.7 GHz ( $4'.4$ :  $S = 5.6 \pm 0.5$  Jy).  
Feldt & Green 1993, A&A, 274, 421. DRAO at 1.4 GHz ( $1' \times 1'.5$ ), including HI, plus CO observations.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G84.9+0.5**

**RA:** 20<sup>h</sup>50<sup>m</sup>30<sup>s</sup>  
**Dec:** +44°53'

**1-GHz flux/Jy:** 0.8  
**Spectral index:** 0.4

**Size/arcmin:** 6  
**Type:** S

**Radio:** Incomplete shell.

**References:**

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz ( $1'.4 \times 1'.0 : S = 1.2 \pm 0.1$  Jy), and northern sky survey at 4.9 GHz ( $S = 0.40 \pm 0.01$  Jy).  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G85.4+0.7**

**RA:** 20<sup>h</sup>50<sup>m</sup>40<sup>s</sup>  
**Dec:** +45°22'

**1-GHz flux/Jy:** ?  
**Spectral index:** 0.5?

**Size/arcmin:** 24  
**Type:** S

**Radio:** Faint, incomplete shell, within larger thermal shell.

**X-ray:** Detected.

**References:**

Kothes *et al.* 2001, A&A, 376, 641. DRAO at 408 MHz ( $4'.4 \times 2'.8 : S < 0.45$  Jy) and 1.4 GHz ( $1'.1 \times 0'.8$ ), plus H<sub>i</sub>, X-ray and optical data.

**G85.9-0.6**

**RA:** 20<sup>h</sup>58<sup>m</sup>40<sup>s</sup>  
**Dec:** +44°53'

**1-GHz flux/Jy:** ?  
**Spectral index:** 0.5?

**Size/arcmin:** 24  
**Type:** S

**Radio:** Faint, incomplete shell.

**X-ray:** Detected.

**References:**

Kothes *et al.* 2001, A&A, 376, 641. DRAO at 408 MHz ( $4'.4 \times 2'.8 : S < 0.9$  Jy) and 1.4 GHz ( $1'.1 \times 0'.8$ ), plus H<sub>i</sub>, X-ray and optical data.

**G89.0+4.7**

HB21

**RA:** 20<sup>h</sup>45<sup>m</sup>00<sup>s</sup>  
**Dec:** +50°35'

**1-GHz flux/Jy:** 220  
**Spectral index:** 0.40

**Size/arcmin:** 120 × 90  
**Type:** S

**Radio:** Distorted shell (4C50.52, an extragalactic double, is within the boundary of the remnant).

**Optical:** Filaments possibly associated.

**X-ray:** Centrally brightened.

**Distance:** Various associations suggest 0.8 kpc.

**References:**

Hirabayashi & Takahashi 1972, PASJ, 24, 231. 30-m dish at 4.2 GHz ( $11' : S = 160 \pm 40$  Jy).  
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ( $5' : S = 148 \pm 16$  Jy), plus optical filaments.  
Hill 1974, MNRAS, 169, 59. Half-Mile Telescope at 1.4 GHz ( $3' \times 3'.9$ ).  
Haslam *et al.* 1975, A&A, 39, 453. Effelsberg 100-m at 2.7 GHz ( $4'.4$ ).  
Fukui & Tatematsu 1988, in SNRISM, p261. CO observations of the vicinity ( $2'.7$ ).  
Tatematsu *et al.* 1990, A&A, 237, 189. DRAO at 408 MHz ( $3'.5 \times 4'.5$ ) and 1.4 GHz ( $1'.0 \times 1'.3$ ), including H<sub>i</sub>, plus CO observations of adjacent molecular cloud.  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Leahy & Aschenbach 1996, A&A, 315, 260. ROSAT observations.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
Koo *et al.* 2001, ApJ, 552, 175. NRAO 12-m and other CO observations ( $27''$  and  $45''$ ) of eastern part.

<b>G93.3+6.9</b>	<b>DA 530, 4C(T)55.38.1</b>
<b>RA:</b> 20 <sup>h</sup> 52 <sup>m</sup> 25 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 9
<b>Dec:</b> +55°21'	<b>Spectral index:</b> 0.54
Has been called G93.2+6.7.	
<b>Radio:</b> Shell, with two bright limbs, highly polarized.	
<b>Distance:</b> HI observations suggest 2.5 kpc.	
<b>References:</b>	
Roger & Costain 1976, A&A, 51, 151. DRAO at 1.42 GHz ( $2' \times 2'.4 : S = 6.9$ Jy).	
Haslam <i>et al.</i> 1980, A&A, 92, 57. Effelsberg 100-m at 1.72 GHz ( $7'.6 : S = 6.47 \pm 0.52$ Jy) and 2.7 GHz ( $4'.4 : S = 5.64 \pm 0.64$ Jy), plus review of flux densities.	
Lalitha <i>et al.</i> 1984, A&A, 131, 196. Effelsberg 100-m at 4.75 GHz (smoothed to $3' : S = 4.01 \pm 0.57$ Jy).	
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.	
Landecker <i>et al.</i> 1999, ApJ, 527, 866. DRAO at 408 MHz ( $3'.5 \times 4'.3$ ) and 1.4 GHz ( $1'.0 \times 1'.2$ ), including polarisation and HI.	
<b>G93.7–0.2</b>	<b>CTB 104A, DA 551</b>
<b>RA:</b> 21 <sup>h</sup> 29 <sup>m</sup> 20 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 65
<b>Dec:</b> +50°50'	<b>Spectral index:</b> 0.3
Has been called G93.6–0.2 and G93.7–0.3.	
<b>Radio:</b> Faint shell with possible a steep spectrum core.	
<b>References:</b>	
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 18.4 \pm 1.0$ Jy).	
Mantovani <i>et al.</i> 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ( $7'.6 : S = 53.5 \pm 5.0$ Jy), plus review of flux densities.	
Landecker <i>et al.</i> 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 58 \pm 6$ Jy).	
Mantovani <i>et al.</i> 1991, A&A, 247, 545. Effelsberg 100-m at 4.75 GHz (smoothed to $3' : S = 33.5 \pm 4.0$ Jy), including polarization, plus review of flux densities.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.	
Koralesky <i>et al.</i> 1998, AJ, 116, 1323. VLA search for OH emission.	
<b>G94.0+1.0</b>	<b>3C434.1</b>
<b>RA:</b> 21 <sup>h</sup> 24 <sup>m</sup> 50 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 15
<b>Dec:</b> +51°53'	<b>Spectral index:</b> 0.44
<b>Radio:</b> Incomplete shell, in complex region.	<b>Size/arcmin:</b> 30 × 25
<b>References:</b>	<b>Type:</b> S
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ( $5' : S = 6.1 \pm 0.8$ Jy), and VRO 37-m at 1.7 GHz ( $S = 11 \pm 3$ Jy).	
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 5.8 \pm 0.4$ Jy). Also NRAO 140-ft at 5 GHz ( $6'$ ).	
Mantovani <i>et al.</i> 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ( $7'.6 : S = 12.0 \pm 1.3$ Jy), plus review of flux densities.	
Goss <i>et al.</i> 1984, A&A, 138, 469. WSRT at 610 MHz (smoothed to $100'' : S = 16 \pm 1.7$ Jy) and Effelsberg 100-m at 4.75 GHz ( $2'.4 : S = 7.2 \pm 0.5$ Jy).	
Landecker <i>et al.</i> 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 16 \pm 3$ Jy).	
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.	

**G106.3+2.7**

**RA:** 22<sup>h</sup>27<sup>m</sup>30<sup>s</sup>  
**Dec:** +60°50'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.6

**Size/arcmin:** 60 × 24  
**Type:** ?

Incorporates the proposed smaller remnant G106.6+2.9.

**Radio:** Faint extended source, which brighter ‘head’ to NE.

**Point sources:** Pulsar in ‘head’.

**References:**

- Pineault & Joncas 2000, AJ, 120, 3218. DRAO at 408 MHz ( $3'.9 \times 3'.5 : S = 10.5 \pm 0.3$  Jy) and 1.4 GHz ( $1'.2 \times 1'.0 : S = 4.9 \pm 0.6$  Jy), plus HI.  
 Halpern *et al.* 2001, ApJ, 547, 323. X-ray and radio observations of the ‘head’.  
 Halpern *et al.* 2001, ApJ, 552, L125. Pulsar detection.  
 Kothes *et al.* 2001, ApJ, 560, 236. DRAO at 1.4 GHz, including HI, plus CO and other observations.

**G109.1–1.0**

**RA:** 23<sup>h</sup>01<sup>m</sup>35<sup>s</sup>  
**Dec:** +58°53'

**1-GHz flux/Jy:** 20  
**Spectral index:** 0.50

**CTB 109**  
**Size/arcmin:** 28  
**Type:** S

**Radio:** Semicircular shell, with the Molecular cloud S152 is to the immediate W.

**X-ray:** Incomplete shell, with pulsar at W edge.

**Point sources:** Long period X-ray pulsar.

**References:**

- Hughes *et al.* 1981, ApJ, 246, L127. WSRT at 610 MHz ( $1' : S = 40 \pm 5$  Jy) shows bad CLEAN artefacts.  
 Blair & Kirshner 1981, Natur, 291, 132. Optical spectra.  
 Downes 1983, MNRAS, 203, 695. Effelsberg 100-m at 2.7 GHz ( $4'.4 : S = 13.0 \pm 1.5$  Jy).  
 Sofue *et al.* 1983, PASJ, 35, 447. NRO 45-m at 10.2 GHz ( $2'.7$ ).  
 Hughes *et al.* 1984, ApJ, 283, 147. WSRT at 610 MHz ( $20'' : S = 26 \pm 3$  Jy) 1.4 GHz ( $10''$ ), DRAO at 1.4 GHz ( $1' \times 1'.2 : S = 16.8 \pm 2$  Jy) and Algonquin 46-m at 6.5 GHz ( $4'.5 : S = 6.7 \pm 1$  Jy).  
 Braun & Strom 1986, A&AS, 63, 345. WSRT HI observations.  
 Tatematsu *et al.* 1987, PASJ, 39, 755. NRO 45-m at 10 GHz ( $2'.7$ ), plus polarization.  
 Tatematsu *et al.* 1987, A&A, 184, 279. CO observations of the surroundings ( $2'.7$ ).  
 Hanson *et al.* 1988, A&A, 195, 114. EXOSAT of pulsar.  
 Morini *et al.* 1988, ApJ, 333, 777. EXOSAT observations.  
 Koyama *et al.* 1989, PASJ, 41, 461. X-ray observations of pulsar.  
 Coe *et al.* 1989, MNRAS, 238, 649. IRAS observations of surroundings.  
 Green 1989, MNRAS, 238, 737. OH observations.  
 Tatematsu *et al.* 1990, ApJ, 351, 157. CO of surroundings, plus X-ray observations.  
 Davies & Coe 1991, MNRAS, 249, 313. Optical and IR observations near pulsar.  
 Hurford & Fesen 1995, MNRAS, 277, 549. ROSAT imaging.  
 Rho & Petre 1997, ApJ, 484, 828. ROSAT observations.  
 Parmar *et al.* 1998, A&A, 330, 175. X-ray observations.  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.  
 Patel *et al.* 2001, ApJ, 563, L45. Chandra observations of pulsar.

**G111.7–2.1**

**RA:** 23<sup>h</sup>23<sup>m</sup>26<sup>s</sup>  
**Dec:** +58°48'

**1-GHz flux/Jy:** 2720  
**Spectral index:** 0.77

**Cassiopeia A, 3C461**

**Size/arcmin:** 5  
**Type:** S

Presumably the remnant of a late 17th century SN.

**Radio:** Bright shell with compact knots and extended plateau of emission.

**Optical:** Fast knots and quasi-stationary flocculli, with many filaments at large radii, and NE ‘jet’.

**X-ray:** Incomplete shell, with hard spectral component, and compact central source.

**Distance:** Optical expansion, plus proper motions indicate 3.4 kpc.

**References:**

- Woan & Duffett-Smith 1990, MNRAS, 243, 87. Structure at 81.5 MHz.  
Rees 1990, MNRAS, 243, 637. Flux density decrease at 38 MHz.  
Anderson *et al.* 1991, ApJ, 373, 146. VLA at 1.4 and 5 GHz (1''.3) for spectral variations.  
Hook *et al.* 1992, A&A, 255, 285. Flux density decrease at 81.5 MHz.  
Woan & Duffett-Smith 1993, MNRAS, 260, 693. Deep pulsar search.  
Guame *et al.* 1994, ApJ, 425, 127. NH<sub>3</sub> absorption and CO observations.  
Reed *et al.* 1995, ApJ, 440, 706. Optical spectra, for 3-d structure.  
Anderson *et al.* 1995, ApJ, 441, 300. VLA at 1.4 GHz and 4.8 GHz (both 1''.3), for polarization studies.  
Anderson & Rudnick 1995, ApJ, 441, 307. VLA at 1.4 GHz and 4.8 GHz, for proper motion studies.  
The *et al.* 1995, ApJ, 444, 244.  $\gamma$ -ray observations.  
Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.  
Liszt & Lucas 1995, A&A, 294, 811. NRAO 12-m observations of molecular emission and absorption.  
Lawrence *et al.* 1995, AJ, 109, 2635. Optical imaging spectroscopy.  
Kassim *et al.* 1995, ApJ, 455, L59. VLA at 74 (25'') and 332 MHz (6''), including spectral index comparisons.  
Anderson & Rudnick 1996, ApJ, 456, 234. VLA for spectral index comparisons.  
Agafanov 1996, A&A, 306, 578. Flux density decrease at 102 MHz.  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Vink *et al.* 1996, A&A, 307, L41. ASCA spectrum.  
Lagage *et al.* 1996, A&A, 315, L273. ISO observations.  
Keohane *et al.* 1996, ApJ, 466, 309. Radio and X-ray comparison.  
Hurford & Fesen 1996, ApJ, 469, 246. Optical spectroscopy and reddening studies.  
Fesen & Gunderson 1996, ApJ, 470, 967. Optical observations of NE ‘jet’.  
Schwarz *et al.* 1997, A&AS, 123, 43. WSRT at 1.4 GHz (30'') for HI absorption.  
Reynoso *et al.* 1997, A&A, 317, 203. VLA at 1.4 GHz (11''  $\times$  14'') for HI absorption.  
Dupraz *et al.* 1997, A&A, 324, 683.  $\gamma$ -ray line detection.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.  
Vink *et al.* 1998, A&A, 339, 201. Comparison of ROSAT and Einstein images for expansion studies.  
Keohane *et al.* 1998, ApJ, 503, L175. Deep ROSAT image, and comparison with radio.  
Koralesky *et al.* 1998, ApJ, 505, L27. Comparison of ROSAT and Einstein images for expansion studies.  
Wright *et al.* 1999, ApJ, 518, 284. BIMA at 83 GHz (6''.5  $\times$  6''.2) at 28 GHz, for comparison with other radio images for spectral index studies.  
Arendt *et al.* 1999, ApJ, 521, 234. ISO observations.  
O’Sullivan & Green 1999, MNRAS, 303, 575. Flux density changes at 13.5, 15.5 and 16.5 GHz.  
Agüeros & Green 1999, MNRAS, 305, 957. CLFST at 151 MHz for bulk expansion studies.  
Mason *et al.* 1999, AJ, 118, 2908. Flux density at 32 GHz ( $S=194 \pm 5$  Jy).  
Vink *et al.* 1999, A&A, 344, 289. X-ray line and continuum observations.  
Liszt & Lucas 1999, A&A, 347, 258. NRAO 12-m at 86 GHz (72'':  $S=101 \pm 3.3$  Jy) and 140 GHz (44'':  $S=69.6 \pm 6.3$  Jy), plus CO observations.  
Douvion *et al.* 1999, A&A, 352, L111. ISO observations.  
Hughes *et al.* 2000, ApJ, 528, L109. Chandra observations.  
Reichart & Stephens 2000, ApJ, 537, 904. Flux density changes at 1405 MHz.  
Hwang *et al.* 2000, ApJ, 537, L119. Chandra imaging, including lines.  
McLaughlin *et al.* 2001, ApJ, 547, L41. Radio upper limit for central source.  
Chakrabarty *et al.* 2001, ApJ, 548, 800. Chandra observations, including central compact source.  
Ryan *et al.* 2001, ApJ, 548, 811. Optical limits in compact central source.  
Gothelf *et al.* 2001, ApJ, 552, L39. Chandra observations, showing outer shock.  
Kaplan *et al.* 2001, ApJ, 558, 270. Search for near-IR counterpart to central source.  
Vink *et al.* 2001, ApJ, 560, L79. BeppoSAX X-ray line observations.  
Fesen 2001, ApJS, 133, 161. Optical imaging and spectroscopy.  
Bleeker *et al.* 2001, A&A, 365, L225. XMM-Newton observations.  
Maccarone *et al.* 2001, A&A, 368, 267. BeppoSAX observations.  
Douvion *et al.* 2001, A&A, 369, 589. ISO observations.  
Aharonian *et al.* 2001, A&A, 370, 112. High energy  $\gamma$ -ray emission detection.  
Gerardy & Fesen 2001, AJ, 121, 2781. IR spectroscopy and imaging.  
Thorstensen *et al.* 2001, AJ, 122, 297. Optical expansion studies.  
Hwang *et al.* 2001, ApJ, 560, L175. Chandra observations of Doppler shifted lines.  
Fesen *et al.* 2001, AJ, 122, 2644. HST observations.

**G114.3+0.3**

**RA:** 23<sup>h</sup>37<sup>m</sup>00<sup>s</sup>  
**Dec:** +61°55'

**1-GHz flux/Jy:** 6?  
**Spectral index:** 0.3?

**Size/arcmin:** 90 × 55  
**Type:** S

**Radio:** Shell, with HII region S165 within the boundary of the remnant.

**Distance:** Possible association with HI features suggests 3.0–3.8 kpc.

**Point sources:** Pulsar near centre of remnant.

**References:**

- Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4'.4 :  $S = 3.6$  Jy) and  $S_{1.4 \text{ GHz}} = 4.4$  Jy from 1.4 GHz survey data, plus HI from Maryland-Green Bank survey.  
Kulkarni *et al.* 1993, Natur, 362, 135. Pulsar association.  
Fürst *et al.* 1993, A&A, 276, 470. Pulsar association.  
Becker *et al.* 1996, A&A, 306, 464. ROSAT of pulsar.  
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.

**G116.5+1.1**

**RA:** 23<sup>h</sup>53<sup>m</sup>40<sup>s</sup>  
**Dec:** +63°15'

**1-GHz flux/Jy:** 11?  
**Spectral index:** 0.8?

**Size/arcmin:** 80 × 60  
**Type:** S

**Radio:** Distinct shell, with high polarization.

**Distance:** Possible association with HI features suggests 3.6–5.2 kpc.

**References:**

- Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4'.4 :  $S = 4.7 \pm 0.4$  Jy) and  $S_{1.4 \text{ GHz}} = 8.0 \pm 0.8$  Jy from 1.4 GHz survey data, plus HI from Maryland-Green Bank survey.  
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

**G116.9+0.2**

CTB 1

**RA:** 23<sup>h</sup>59<sup>m</sup>10<sup>s</sup>  
**Dec:** +62°26'

**1-GHz flux/Jy:** 9?  
**Spectral index:** 0.5?

**Size/arcmin:** 34  
**Type:** S

Has been called G117.3+0.1 and G116.9+0.1.

**Radio:** Incomplete shell.

**Optical:** Filaments on sky survey.

**X-ray:** Centrally brightened, with NE ‘breakout’.

**Point sources:** Pulsar to NE.

**Distance:** Possible association with HI features suggests 2.8–4.0 kpc, mean optical velocity suggests 2.7 kpc.

**References:**

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' :  $S = 3.9 \pm 1.0$  Jy), and VRO 37-m at 1.7 GHz ( $S = 5.5 \pm 2.0$  Jy), plus review of flux densities.  
van den Bergh *et al.* 1973, ApJS, 26, 19. Optical.  
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5' :  $S = 4.2 \pm 0.2$  Jy).  
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 :  $S = 3.0 \pm 0.3$  Jy).  
Dicke & Willis 1980, A&A, 85, 55. WSRT at 610 MHz (56'' × 64'') and 1.4 GHz (22'' × 25'').  
Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4'.4 :  $S = 4.8 \pm 0.4$  Jy) and  $S_{1.4 \text{ GHz}} = 7.8 \pm 0.8$  Jy from 1.4 GHz survey data, plus HI from Maryland-Green Bank survey.  
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.  
Landeker *et al.* 1982, AJ, 87, 1379. DRAO at 1.42 GHz and HI (2' × 2'.3 :  $S = 8.3 \pm 0.5$  Jy), plus review of flux densities.  
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.  
Hailey & Craig 1994, ApJ, 434, 635. Optical spectroscopy.  
Hailey & Craig 1995, ApJ, 455, L151. ROSAT of nearby pulsar.  
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.  
Craig *et al.* 1997, ApJ, 488, 307. ROSAT observations.  
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

<b>G119.5+10.2</b>		<b>CTA 1</b>
<b>RA:</b> 00 <sup>h</sup> 06 <sup>m</sup> 40 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 36	<b>Size/arcmin:</b> 90?
<b>Dec:</b> +72°45'	<b>Spectral index:</b> 0.6	<b>Type:</b> S
Has been called G119.5+10.3.		
<b>Radio:</b> Incomplete shell, with ‘breakout’ to NW.		
<b>Optical:</b> Faint diffuse nebulosities.		
<b>X-ray:</b> Centrally brightened.		
<b>Point sources:</b> Compact, central X-/γ-ray source.		
<b>References:</b>		
Sieber <i>et al.</i> 1979, A&A, 74, 361. Effelsberg 100-m at 2.7 GHz ( $4'.4 : S = 23.6 \pm 2.8$ Jy).		
Sieber <i>et al.</i> 1981, A&A, 103, 393. Effelsberg 100-m at 2.7 GHz and 1.7 GHz ( $7'.6 : S = 31.6 \pm 2.5$ Jy), and 6C at 151 MHz ( $4' : S = 62.6 \pm 6$ Jy).		
Fesen <i>et al.</i> 1981, ApJ, 247, 148. Optical, including spectra.		
Fesen <i>et al.</i> 1983, ApJS, 51, 337. Deep [OIII] imagery.		
Seward 1990, ApJS, 73, 781. Einstein observations.		
Simonetti 1992, ApJ, 386, 170. VLA observations of background sources for rotation measure studies.		
Pineault <i>et al.</i> 1993, AJ, 105, 1060. DRAO at 1.4 GHz ( $1' : S = 34 \pm 4$ Jy), plus HI and IRAS.		
Seward <i>et al.</i> 1995, ApJ, 453, 284. ROSAT observations.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Slane <i>et al.</i> 1997, ApJ, 485, 221. ASCA and ROSAT observations.		
Pineault <i>et al.</i> 1997, A&A, 324, 1152. DRAO at 408 MHz ( $3'.5$ ) and 1.4 GHz ( $1'.0$ ).		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
Brazier <i>et al.</i> 1998, MNRAS, 295, 819. Studies of central, compact γ-ray source.		
Mavromatakis <i>et al.</i> 2000, A&A, 353, 371. Optical imagery.		
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.		

**G120.1+1.4**

**RA:** 00<sup>h</sup>25<sup>m</sup>18<sup>s</sup>  
**Dec:** +64°09'

**1-GHz flux/Jy:** 56  
**Spectral index:** 0.61

Tycho, 3C10, SN1572

**Size/arcmin:** 8  
**Type:** S

This is the remnant of the Tycho's SN of AD1572.

**Radio:** Shell, brightest to the NE.

**Optical:** Faint filaments/knots to the NNW, NE and E.

**X-ray:** Shell, coincident with the radio shell, brighter to the NE.

**Point sources:** Faint radio source near centre of the remnant, thought to be extragalactic.

**Distance:** HI absorption gives 2–5 kpc, optical proper motion and shock velocity gives 2.4 kpc.

**References:**

- Duin & Strom 1975, A&A, 39, 33. WSRT at 610 MHz ( $57'' \times 64''$ ) and 5 GHz ( $7'' \times 8''$ ).  
 Klein *et al.* 1979, A&A, 76, 120. Effelsberg 100-m at 10.7 GHz ( $1''.2 : S = 13.1 \pm 0.8$  Jy), plus review of flux densities.  
 Strom *et al.* 1982, MNRAS, 200, 473. WSRT at 1415 MHz ( $27'' \times 31''$ ) from 1971 and 1979, for expansion.  
 Dickel *et al.* 1982, ApJ, 257, 145. Comparison of Radio, X-ray and Optical.  
 Reid *et al.* 1982, ApJ, 261, 485. Einstein observations.  
 Seward *et al.* 1983, ApJ, 266, 287. Einstein observations.  
 Tan & Gull 1985, MNRAS, 216, 949. 5km at 2.7 GHz ( $4''$ ) from 1980 and 1983, and One-Mile Telescope at 1.4 GHz ( $23''$ ) from 1965 and 1980, for expansion.  
 Albinson *et al.* 1986, MNRAS, 219, 427. HI observations.  
 Green & Gull 1987, MNRAS, 224, 1055. VLA HI absorption observations towards central radio source.  
 Smith *et al.* 1988, ApJ, 325, 288. EXOSAT spectrum and image (smoothed to  $13''$ ).  
 Dickel *et al.* 1991, AJ, 101, 2151. VLA at 1.4 and 5 GHz ( $1''.5$ ).  
 Wood *et al.* 1992, AJ, 103, 1338. VLA at 5 GHz ( $1''.5$ ) polarization studies.  
 Fink *et al.* 1994, A&A, 283, 635. X-ray spectra.  
 Vancura *et al.* 1995, ApJ, 441, 680. X-ray spectra and ROSAT image.  
 Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.  
 Schwarz *et al.* 1995, A&A, 299, 193. WSRT and Effelsberg 100-m at 1.4 GHz ( $50''$ ), and VLA at 1.4 GHz ( $13''$ ), for neutral hydrogen studies.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
 Hwang & Gotthelf 1997, ApJ, 475, 665. ASCA observations.  
 Reynoso *et al.* 1997, ApJ, 491, 816. VLA at 1.4 GHz ( $1''.4 \times 1''.5$ ) from 1984 and 1994 for expansion studies.  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
 Reynoso *et al.* 1999, AJ, 117, 1827. VLA at 1.4 GHz ( $13''$ ), for HI studies.  
 Katz-Stone *et al.* 2000, ApJ, 529, 453. VLA at 333 MHz and 1.4 GHz ( $9''.5 \times 7''.7$ ), for spectral index studies.  
 Ghavamian *et al.* 2000, ApJ, 535, 266. Optical observations of shock precursors.  
 Hughes 2000, ApJ, 545, L53. ROSAT X-ray expansion.  
 Decourchelle *et al.* 2001, A&A, 365, L218. XMM-Newton observations.  
 Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.  
 Douvion *et al.* 2001, A&A, 373, 281. ISO observations.  
 Aharonian *et al.* 2001, A&A, 373, 292. Upper limit on high energy  $\gamma$ -rays.

**G126.2+1.6**

**RA:** 01<sup>h</sup>22<sup>m</sup>00<sup>s</sup>  
**Dec:** +64°15'

**1-GHz flux/Jy:** 7  
**Spectral index:** varies

**Size/arcmin:** 70  
**Type:** S?

**Radio:** Poorly defined shell.

**Optical:** Filaments detected.

**References:**

- Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz ( $9' : S = 6.8 \pm 0.7$  Jy) and 2.7 GHz ( $4'.4 : S = 3.9 \pm 0.4$  Jy).  
 Blair *et al.* 1980, ApJ, 242, 592. Optical detection and spectra.  
 Rosado 1982, RMxAA, 5, 127. Optical spectra.  
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.  
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz ( $4'.4$ ) and 4.8 GHz ( $2'.6$ ).  
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz ( $3'.5 \times 3'.9 : S = 12 \pm 2.5$  Jy) and part at 1.4 GHz ( $1'.1 \times 1'.0$ ), plus review of flux densities.  
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

G127.1+0.5	R5
<b>RA:</b> 01 <sup>h</sup> 28 <sup>m</sup> 20 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 13
<b>Dec:</b> +63°10'	<b>Spectral index:</b> 0.6

Has been called G127.3+0.7.

**Radio:** Distinct shell, with bright central source.

**Point sources:** Flat radio spectrum (extragalactic) source at centre of remnant.

**Optical:** Detected.

**Distance:** 1.2–1.3 kpc if associated with NGC559.

**References:**

- Caswell 1977, MNRAS, 181, 789. Half-Mile Telescope at 1.42 GHz ( $3'.5 \times 3'.9$ ), plus other observations of central source.  
 Salter *et al.* 1978, A&A, 66, 77. Effelsberg 100-m at 2.7 GHz ( $4'.4$ ), plus 5 and 8.7 GHz of the central source.  
 Pauls 1977, A&A, 59, L13. Effelsberg 100-m at 1.4 GHz ( $9':S=8\pm1$  Jy).  
 Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz ( $9':S=10.8\pm1.3$  Jy).  
 Pauls *et al.* 1982, A&A, 112, 120. WSRT at 610 MHz ( $56'' \times 62''$ ) and HI absorption to the point source with the VLA.  
 Geldzahler & Shaffer 1982, ApJ, 260, L69. Observations of central source.  
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz ( $4'.4$ ) and 4.8 GHz ( $2'.6$ ).  
 Goss & van Gorkom 1984, JApA, 5, 425. WSRT HI absorption of central source.  
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz ( $3'.5 \times 3'.9 : S = 17.9 \pm 2.0$  Jy) and 1.4 GHz ( $1'.13 \times 1'.0 : S = 10.1 \pm 0.8$  Jy), plus review of flux densities.  
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

G130.7+3.1	3C58, SN1181
<b>RA:</b> 02 <sup>h</sup> 05 <sup>m</sup> 41 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 33
<b>Dec:</b> +64°49'	<b>Spectral index:</b> 0.10

This is the remnant of the SN of AD1181.

**Radio:** Filled-centre, highly polarized, with high frequency turnover.

**Optical:** Faint filaments.

**X-ray:** Centrally brightened with power-law spectrum.

**Point sources:** Compact source in X-rays.

**Distance:** HI absorption indicates 3.2 kpc.

**References:**

- Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz ( $58'' : S = 26.7 \pm 0.5$  Jy).  
 Wilson & Weiler 1976, A&A, 49, 357. WSRT at 610 MHz ( $58'' \times 64''$ ), 1.4 GHz ( $24'' \times 27''$ ) and 5 GHz ( $7'' \times 8''$ ).  
 van den Bergh 1978, ApJ, 220, L9. Optical observations.  
 Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.  
 Green & Gull 1982, Natur, 299, 606. HI absorption distance.  
 Fesen 1983, ApJ, 270, L53. Optical spectra.  
 Reynolds & Aller 1985, AJ, 90, 2312. VLA at 1.4 GHz ( $2'$ ), for limits of shell.  
 Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT spectrum.  
 Green 1986, MNRAS, 218, 533. CLFST at 151 MHz ( $1'.2 \times 1'.3 : S = 36 \pm 4$  Jy), 5km at 2.7 GHz ( $4''$ ), plus Einstein observations for limit on shell.  
 Green 1987, MNRAS, 225, 11P. Flux density increase at 408 MHz.  
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz ( $26''.5 : S = 24.2 \pm 1.4$  Jy).  
 Reynolds & Aller 1988, ApJ, 327, 845. VLA at 1.4 ( $2''.4$ ) and 4.9 GHz ( $2''.5$ ).  
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ( $90'' : S = 15.0 \pm 2.0$  Jy), plus review of flux densities.  
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.  
 Green & Scheuer 1992, MNRAS, 258, 833. IRAS upper limits.  
 Roberts *et al.* 1993, A&A, 274, 427. HI absorption.  
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.  
 Helfand *et al.* 1995, ApJ, 453, 741. ROSAT observations.  
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.  
 Torii *et al.* 2000, PASJ, 52, 875. ASCA observations.  
 Bietenholz *et al.* 2001, ApJ, 560, 772. VLA at 74 MHz ( $26'' : S = 33.6$  Jy) and 327 MHz ( $8''.2 : S = 33.9$  Jy), for spectral index studies, and comparison with earlier observations for expansion studies.  
 Bocchino *et al.* 2001, A&A, 369, 1078. XMM-Newton observations.

---

<b>G132.7+1.3</b>		HB3
<b>RA:</b> 02 <sup>h</sup> 17 <sup>m</sup> 40 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 45	<b>Size/arcmin:</b> 80
<b>Dec:</b> +62°45'	<b>Spectral index:</b> 0.6	<b>Type:</b> S

Has been called G132.4+2.2.

**Radio:** Faint shell, adjacent to W3/4/5 complex.

**Optical:** Complete, filamentary shell, shock excited spectra.

**X-ray:** Partial shell.

**Point sources:** Pulsar nearby.

**Distance:** Interaction with surroundings suggests 2.2 kpc.

**References:**

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical.

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ( $5' : S = 33.8 \pm 7.0$  Jy).

d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.

Read 1981, MNRAS, 194, 863. 6C at 151 MHz ( $4'.4$ ) and Half-Mile Telescope at 1.4 GHz ( $2'$ ) showing H $\alpha$  shell.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Fesen & Gull 1983, PASP, 95, 196. Optical image.

Leahy *et al.* 1985, ApJ, 294, 183. Einstein observations.

Landecker *et al.* 1987, AJ, 94, 111. DRAO at 408 MHz ( $3'.5 \times 4' : S = 75 \pm 15$  Jy), plus review of flux densities.

Routledge *et al.* 1991, A&A, 247, 529. DRAO at 1.4 GHz ( $1'.0 \times 1'.1$ ) for H $\alpha$ , plus CO observations.

Fesen *et al.* 1995, AJ, 110, 2876. Optical imaging and spectroscopy, DRAO at 408 MHz ( $3'.5 \times 4'$ ) and 1.4 GHz ( $1'.0 \times 1'.1$ ).

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.

---

<b>G156.2+5.7</b>		
<b>RA:</b> 04 <sup>h</sup> 58 <sup>m</sup> 40 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 5	<b>Size/arcmin:</b> 110
<b>Dec:</b> +51°50'	<b>Spectral index:</b> 0.5	<b>Type:</b> S

**Radio:** Faint shell.

**X-ray:** Faint shell.

**References:**

Pfeffermann *et al.* 1991, A&A, 246, L28. ROSAT detection.

Reich *et al.* 1992, A&A, 256, 214. Effelsberg 100-m at 1.4 ( $9' : S = 4.2 \pm 1.0$  Jy) and 2.7 GHz ( $4'.3 : S = 3.0 \pm 1.0$  Jy), plus H $\alpha$  and IRAS.

Yamauchi *et al.* 1993, PASJ, 45, 795. Hard X-ray observations.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Yamauchi *et al.* 1999, PASJ, 51, 13. ASCA observations of some regions.

G160.9+2.6		HB9
<b>RA:</b> 05 <sup>h</sup> 01 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 110	<b>Size/arcmin:</b> 140 × 120
<b>Dec:</b> +46°40'	<b>Spectral index:</b> 0.6	<b>Type:</b> S
Has been called G160.5+2.8 and G160.4+2.8.		
<b>Radio:</b> Large, filamentary shell.		
<b>Optical:</b> Incomplete shell.		
<b>X-ray:</b> Centrally brightened.		
<b>Point sources:</b> Pulsar within boundary of the remnant, plus several nearby compact radio sources.		
<b>Distance:</b> Various observations suggests less than 4 kpc.		
<b>References:</b>		
d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.		
Damashek <i>et al.</i> 1978, ApJ, 225, L31. Pulsar.		
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.		
Dwarkanath <i>et al.</i> 1982, JApA, 3, 207. Radio observations at 34.5 MHz ( $26' \times 40' : S = 750 \pm 150$ Jy), plus review of flux densities.		
van Gorkom <i>et al.</i> 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source.		
Sequoia & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.		
Leahy 1988, ApJ, 322, 917. Einstein observations.		
Leahy & Roger, 1991, AJ, 101, 1033. DRAO at 408 MHz ( $3'.5 \times 4'.8$ ) and 1.4 GHz ( $1'.0 \times 1'.4$ ), including HI and discussion of distance.		
Yamauchi & Koyama 1993, PASJ, 45, 545. Hard X-ray observations.		
Leahy & Aschenbach 1995, A&A, 293, 853. ROSAT observations.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Leahy <i>et al.</i> 1998, A&A, 339, 601. Miyun at 232 MHz ( $3'.8 \times 5'.2$ ), CLFST at 151 MHz ( $4'.2 \times 5'.8$ ) and Effelsberg 100-m at 4.7 GHz ( $2'.5$ ) for spectral index studies.		
Roger <i>et al.</i> 1999, A&AS, 137, 7. 22 MHz flux density ( $S = 1130 \pm 340$ Jy).		

G166.0+4.3		VRO 42.05.01
<b>RA:</b> 05 <sup>h</sup> 26 <sup>m</sup> 30 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 7?	<b>Size/arcmin:</b> 55 × 35
<b>Dec:</b> +42°56'	<b>Spectral index:</b> 0.4?	<b>Type:</b> S
<b>Radio:</b> Two arcs of strikingly different radii.		
<b>Optical:</b> Nearly complete ring.		
<b>X-ray:</b> Predominantly in SW.		
<b>Distance:</b> HI indicates 4.5 kpc.		
<b>References:</b>		
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical.		
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ( $5' : S = 5.2 \pm 1.0$ Jy).		
Lozinskaya 1979, AuPh, 32, 113. H $\alpha$ interferometry.		
Landecker <i>et al.</i> 1982, ApJ, 261, L41. DRAO at 1.4 GHz ( $1'.0 \times 1'.4$ ), plus review of flux densities.		
Fesen <i>et al.</i> 1983, ApJS, 51, 337. Deep [OIII] imagery.		
Pineault <i>et al.</i> 1985, A&A, 151, 52. VLA at 1.4 GHz ( $16'' \times 20''$ ) of part of remnant, and optical.		
Pineault <i>et al.</i> 1987, ApJ, 315, 580. DRAO and VLA combined at 1.4 GHz ( $20''$ ).		
Fesen <i>et al.</i> 1985, ApJ, 292, 29. Optical spectra.		
Braun & Strom 1986, A&AS, 63, 345. WSRT HI Observations.		
Landecker <i>et al.</i> 1989, MNRAS, 237, 277. DRAO at 1.4 GHz ( $1'.0 \times 1'.4$ ), including HI.		
Burrows & Guo 1994, ApJ, 421, L19. ROSAT images and spectra.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Guo & Burrows 1997, ApJ, 480, L51. ASCA observations.		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		

<b>G166.2+2.5</b>		OA 184
<b>RA:</b> 05 <sup>h</sup> 19 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 11	<b>Size/arcmin:</b> 90 × 70
<b>Dec:</b> +41°55'	<b>Spectral index:</b> 0.5	<b>Type:</b> S
<b>Radio:</b> Large, faint shell.		
<b>Optical:</b> Nearly complete ring.		
<b>Distance:</b> HI indicates 8 ± 2 kpc.		
<b>References:</b>		
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ( $5' : S = 6.0 \pm 2.2$ Jy).		
Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo 111 MHz.		
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.		
Fesen <i>et al.</i> 1985, ApJ, 292, 29. Optical spectra.		
Routledge <i>et al.</i> 1986, MNRAS, 221, 809. DRAO at 1.4 GHz ( $1' \times 1'.4 : S = 9.0 \pm 0.5$ Jy) including HI, plus review of flux densities.		
Kim <i>et al.</i> 1988, AJ, 96, 704. VLA of background sources for magnetic field structure of the remnant.		
Leahy & Marshall 1988, MNRAS, 235, 805. IRAS observations.		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
<b>G179.0+2.6</b>		
<b>RA:</b> 05 <sup>h</sup> 53 <sup>m</sup> 40 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 7	<b>Size/arcmin:</b> 70
<b>Dec:</b> +31°05'	<b>Spectral index:</b> 0.4	<b>Type:</b> S?
<b>Radio:</b> Thick shell, with background extragalactic sources near centre.		
<b>References:</b>		
Fürst & Reich 1986, A&A, 154, 303. Effelsberg 100-m at 1.4 (9'.4), 2.7 (4'.3) and 4.75 GHz (2'.4).		
Fürst <i>et al.</i> 1989, A&A, 223, 66. Observations of central, extragalactic source.		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
<b>G180.0–1.7</b>		S147
<b>RA:</b> 05 <sup>h</sup> 39 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 65	<b>Size/arcmin:</b> 180
<b>Dec:</b> +27°50'	<b>Spectral index:</b> varies	<b>Type:</b> S
<b>Radio:</b> Large faint shell, with spectral break.		
<b>Optical:</b> Wispy ring.		
<b>X-ray:</b> Possible detection.		
<b>Point sources:</b> Pulsar within boundary.		
<b>References:</b>		
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical.		
Sofue <i>et al.</i> 1980, PASJ, 32, 1. Effelsberg 100-m at 5 GHz (2'.6) of parts.		
Kundu <i>et al.</i> 1980, A&A, 92, 225. Effelsberg 100-m at 2.7 GHz ( $5'.5 : S = 34.9 \pm 4$ Jy) and 1.6 GHz ( $10' : S = 60.2 \pm 6$ Jy).		
Angerhofer & Kundu 1981, AJ, 86, 1003. Arecibo at 430 MHz (9' : $S = 97 \pm 20$ Jy).		
Fürst <i>et al.</i> 1982, A&A, 115, 428. Observations of compact radio sources near the remnant.		
Fesen <i>et al.</i> 1985, ApJ, 292, 29. Optical spectra.		
Fürst & Reich 1986, A&A, 163, 185. Effelsberg 100-m at 1.4, 2.7 and 4.7/5.0 GHz (9'.4, 4'.3 and 2'.4/2'.6).		
Sauvageot <i>et al.</i> 1990, A&A, 227, 183. EXOSAT possible detection.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Anderson <i>et al.</i> 1996, ApJ, 468, L55. Pulsar detection.		
<b>G182.4+4.3</b>		
<b>RA:</b> 06 <sup>h</sup> 08 <sup>m</sup> 10 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 1.2	<b>Size/arcmin:</b> 50
<b>Dec:</b> +29°00'	<b>Spectral index:</b> 0.4	<b>Type:</b> S
<b>Radio:</b> Incomplete shell.		
<b>References:</b>		
Kothes <i>et al.</i> 1998, A&A, 331, 661. Effelsberg 100-m at 1.4, 2.7, 4.9 and 10.5 GHz ( $9'.4 : S = 0.36 \pm 0.08$ Jy, $4'.4 : S = 0.25 \pm 0.04$ Jy, $2'.5 : S = 0.20 \pm 0.02$ Jy and $1'.2 : S = 0.15 \pm 0.03$ Jy), plus X-ray upper limit.		

**G184.6–5.8**

**RA:** 05<sup>h</sup>34<sup>m</sup>31<sup>s</sup>  
**Dec:** +22°01'

**1-GHz flux/Jy:** 1040  
**Spectral index:** 0.30

**Crab Nebula, 3C144, SN1054**

**Size/arcmin:** 7 × 5  
**Type:** F

This is the remnant of the SN of AD1054.

**Radio:** Filled-centre, central pulsar, with faint ‘jet’ (or tube) extending from the N edge.

**Optical:** Strongly polarized filaments, diffuse synchrotron emission, with ‘jet’ faintly visible.

**X-ray:** Central ‘torus’ around the pulsar.

**Point sources:** Pulsar powering the remnant.

**Distance:** Proper motions and radial velocities give 2 kpc.

**References:**

- Velusamy 1984, Natur, 308, 251. VLA at 1.4 GHz, radio detection of ‘jet’.  
 Harden & Seward 1984, ApJ, 283, 279. X-ray observations.  
 Velusamy 1985, MNRAS, 212, 359. VLA at 1.4 GHz (15'').  
 Aller & Reynolds 1985, ApJ, 293, L73. Flux density decrease.  
 Fesen & Gull 1986, ApJ, 306, 259. Deep [OIII] imagery of ‘jet’.  
 Marcelin *et al.* 1990, A&A, 228, 471. Optical of ‘jet’.  
 Fesen & Blair 1990, ApJ, 351, L45. Optical identification of dust in filaments.  
 Hester *et al.* 1990, ApJ, 357, 539. Optical and IR images.  
 Graham *et al.* 1990, ApJ, 352, 172. IR spectroscopy and imaging.  
 Hickson & van den Bergh 1990, ApJ, 365, 224. Optical polarization.  
 Bietenholz & Kronberg 1990, ApJ, 357, L13. VLA at 1.4 GHz (1''.8).  
 MacAlpine & Uomoto 1991, AJ, 102, 218. Photometry.  
 Michel *et al.* 1991, ApJ, 368, 463. Optical photometry.  
 Bietenholz *et al.* 1991, ApJ, 373, L59. VLA at multiple epochs for expansion studies.  
 Véron-Cetty & Woltjer 1991, A&A, 251, L31. Limits on optical variations.  
 Hennessy *et al.* 1992, ApJ, 395, L13. UV imagery.  
 Fesen *et al.* 1992, ApJ, 399, 599. Optical studies of ‘dark bays’.  
 Blair *et al.* 1992, ApJ, 399, 611. Far UV observations.  
 Bietenholz & Kronberg 1992, ApJ, 393, 206. VLA at 1.5 and 5 GHz (1''.8) and 1.5 and 14 GHz (6''.5) for spectral studies.  
 Strom & Greidanus 1992, Natur, 358, 654. IRAS observations.  
 Fesen & Staker 1993, MNRAS, 263, 69. [OIII] imaging of ‘jet’, and proper motion studies.  
 Kassim *et al.* 1993, AJ, 106, 2218. VLA at 74 MHz (20'').  
 Véron-Cetty & Woltjer 1993, A&A, 270, 370. Continuum and [OIII] photometry.  
 Nolan *et al.* 1993, ApJ, 409, 697.  $\gamma$ -ray observations.  
 Strong *et al.* 1993, A&AS, 97, 133.  $\gamma$ -ray observations.  
 Murdin 1994, MNRAS, 269, 89. Deep H $\alpha$  plates and spectroscopy of halo.  
 Wallace *et al.* 1994, A&A, 286, 565. H $\alpha$  of surroundings.  
 MacAlpine *et al.* 1994, ApJ, 432, L131. Prominent optical knots.  
 Rudy *et al.* 1994, ApJ, 426, 646. IR spectroscopy.  
 Tanimori *et al.* 1994, ApJ, 429, L61.  $\gamma$ -rays from pulsar.  
 Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.  
 Lawrence *et al.* 1995, AJ, 109, 2635. Optical imaging spectroscopy.  
 Frail *et al.* 1995, ApJ, 454, L129. VLA at 333 MHz (20'') for limits on shell.  
 MacAlpine *et al.* 1996, ApJ, 463, 650. Optical spectroscopy of N and S rich filaments.  
 Nasuti *et al.* 1996, A&A, 314, 849. Spectrophotometry of pulsar.  
 Bietenholz *et al.* 1997, ApJ, 490, 291. Comparison of VLA observations at 74 MHz, 327 MHz, 1.5 GHz and 5 GHz for spectral index studies.  
 Blair *et al.* 1997, ApJS, 109, 473. HST imaging.  
 van der Meulen *et al.* 1998, A&A, 330, 321.  $\gamma$ -ray observations.  
 Tanimori *et al.* 1998, ApJ, 492, L33.  $\gamma$ -ray observations.  
 Nugent 1998, PASP, 110, 831. Optical expansion.  
 Hillas *et al.* 1998, ApJ, 503, 744.  $\gamma$ -ray spectrum.  
 Sankrit *et al.* 1998, ApJ, 504, 344. HST images.  
 Wallace *et al.* 1999, ApJS, 124, 181. DRAO at 1.4 GHz (1''.0 × 2''.8), plus Effelsberg 100-m, for H $\alpha$  studies.  
 Greiveldinger & Aschenbach 1999, ApJ, 510, 305. X-ray variability of torus.  
 Weisskopf *et al.* 2000, ApJ, 536, L81. Chandra observations.  
 Sollerman *et al.* 2000, ApJ, 537, 861. HST observations.  
 Aharonian *et al.* 2000, ApJ, 539, 317. High energy  $\gamma$ -ray observations.  
 Carramiñana *et al.* 2000, ApJ, 542, 974. Optical spectra of pulsar.  
 Oser *et al.* 2001, ApJ, 547, 949.  $\gamma$ -ray observations.  
 Willingale *et al.* 2001, A&A, 365, L212. XMM-Newton observations.  
 Douvion *et al.* 2001, A&A, 373, 281. ISO observations.  
 Bietenholz *et al.* 2001, ApJ, 560, 254. Multi-epoch VLA observations, showing variations near the pulsar.

**G189.1+3.0**

**RA:** 06<sup>h</sup>17<sup>m</sup>00<sup>s</sup>  
**Dec:** +22°34'

**1-GHz flux/Jy:** 160  
**Spectral index:** 0.36

**IC443, 3C157**

**Size/arcmin:** 45  
**Type:** C

**Radio:** Limb-brightened to NE, with faint extension to the E.

**Optical:** Brightest to the NE, with faint filaments outside the NE boundary.

**X-ray:** Shell, brightest to the NE, with nebula and compact source.

**Point sources:** Compact X-ray source in S.

**Distance:** Mean optical velocity suggests 0.7–1.5 kpc, association with S249 gives 1.5–2 kpc.

**References:**

- Duin & van der Laan 1975, A&A, 40, 111. WSRT at 610 MHz (1' × 2'.5), 1415 MHz (24'' × 63''), and part at 5 GHz (12'' × 31'').  
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.  
Fesen 1984, ApJ, 281, 658. Optical of filament to far NE.  
Erickson & Mahoney 1985, ApJ, 290, 596. TPT at 4 frequencies between 31 and 74 MHz (13' × 11' to 5'.4 × 4'.7), plus review of flux densities.  
Braun & Strom 1986, A&A, 164, 193. WSRT at 327 MHz (72'' × 185'') and 1.4 GHz (17'' × 43''), plus H<sub>I</sub> and IRAS.  
Green 1986, MNRAS, 221, 473. CLFST at 151 MHz (1'.2 × 3'.1) and Half-Mile Telescope at 1.4 GHz (2'.1 × 5'.4).  
Mufson *et al.* 1986, AJ, 92, 1349. Radio, IR, optical, UV and X-ray comparison, including VLA at 1.6 GHz (3''.8 × 3''.3 and 40'').  
Brown *et al.* 1988, ApJ, 334, 852. [FeX] optical imaging.  
Dickele *et al.* 1989, AJ, 98, 1363. VLA at 1.4 GHz (1''.1 × 1''.2) of NE.  
Petre *et al.* 1988, ApJ, 335, 215. Einstein and other X-ray observations.  
Burton *et al.* 1990, ApJ, 355, 197. IR observations of shocked O<sub>2</sub>.  
Sauvageot *et al.* 1990, A&A, 232, 203. [FeX] and [FeXIV] imaging.  
Moorhouse *et al.* 1991, MNRAS, 253, 662. Observations of shocked molecular H<sub>2</sub>.  
Wood *et al.* 1991, AJ, 102, 224. VLA at 5 GHz (3''.6 × 3''.8) of northeast, including polarization.  
Teske 1991, ApJ, 383, 233. [FeX] and [FeXIV] imaging.  
Wang & Scoville 1992, ApJ, 386, 158. Observations of shocked molecular species.  
Wang *et al.* 1992, PASJ, 44, 303. Ginga X-ray observations.  
Turner *et al.* 1992, ApJ, 399, 114. Observations of shocked molecular species.  
Dickman *et al.* 1992, ApJ, 400, 203. Observations of shocked molecular species.  
Inoue *et al.* 1993, PASJ, 45, 539. Observations of shocked molecular H<sub>2</sub>.  
van Dishoeck *et al.* 1993, A&A, 279, 541. Observations of shocked molecular species.  
Asaoka & Aschenbach 1994, A&A, 284, 573. X-ray, including possible overlapping remnant.  
Tauber *et al.* 1994, ApJ, 421, 570. Observations of shocked molecular species.  
Richter *et al.* 1995, ApJ, 454, 277. Observations of shocked molecular H<sub>2</sub>.  
Esposito *et al.* 1996, ApJ, 461, 820. Associated  $\gamma$ -ray emission.  
Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.  
Keohane *et al.* 1997, ApJ, 484, 350. ASCA observations.  
Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.  
Claussen *et al.* 1997, ApJ, 522, 349. High resolution observations of OH masers.  
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.  
Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.  
Oliva *et al.* 1999, A&A, 341, L75. ISO observations.  
Cesarsky *et al.* 1999, A&A, 348, 945. ISO observations of shocked molecular H<sub>2</sub>.  
Bocchino & Bykov 2000, A&A, 362, L29. BeppoSAX observations.  
Rho *et al.* 2001, ApJ, 547, 885. ISAO and 2Mass IR spectroscopy and imaging.  
Olbert *et al.* 2001, ApJ, 554, L205. Chandra of compact X-ray source and surrounding nebula.  
Bocchino & Bykov 2001, A&A, 376, 248. XMM-Newton of compact X-ray source and surrounding nebula.  
Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.

**G192.8–1.1**

**RA:** 06<sup>h</sup>09<sup>m</sup>20<sup>s</sup>  
**Dec:** +17°20'

**1-GHz flux/Jy:** 20?  
**Spectral index:** 0.6?

**PKS 0607+17**

**Size/arcmin:** 78  
**Type:** S

Has been called G193.3–1.5. Has been regarded as part of the Origem Loop, a supposed larger remnant.

**Radio:** In complex region.

**Optical:** Encompasses S261 and S254–258.

**References:**

- Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (9': S = 13 ± 15% Jy).  
Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 318 MHz (15': S = 74 ± 32 Jy), and  $S_{610 \text{ MHz}} = 40 \text{ Jy}$ .  
Caswell 1985, AJ, 90, 1076. DRAO at 1.4 GHz (1' × 3'.3: S = 18 ± 3 Jy).  
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.  
Welsh *et al.* 2001, A&A, 372, 516. Far-UV spectroscopy.

**G205.5+0.5**

**RA:** 06<sup>h</sup>39<sup>m</sup>00<sup>s</sup>  
**Dec:** +06°30'

**1-GHz flux/Jy:** 160  
**Spectral index:** 0.5

Monoceros Nebula  
**Size/arcmin:** 220  
**Type:** S

**Radio:** In complex region, parts may be HII regions.

**Optical:** Large ring, near Rosette nebula.

**X-ray:** Possibly detected.

**Distance:** Mean optical velocity suggests 0.8 kpc, low frequency radio absorption suggests 1.6 kpc.

**References:**

- Milne & Dickel 1974, *AuJPh*, 27, 549. Parkes 64-m at 2.7 GHz (9').  
 Velusamy & Kundu 1974, *A&A*, 32, 375. NRAO 300-ft at 2.7 GHz (5'), part only.  
 Dickel & DeNoyer 1975, *AJ*, 80, 437. Arecibo at 111 MHz (1':  $S = 462 \pm 180$  Jy) and  $S_{610 \text{ MHz}} = 245$  Jy.  
 Davies *et al.* 1978, *A&AS*, 31, 271. Deep optical plates.  
 Lozinskaya 1981, *SvAL*, 7, 17. Mean optical velocity.  
 Graham *et al.* 1982, *A&A*, 109, 145. Effelsberg 100-m at 2.7 GHz (4'.4:  $S = 97.6 \pm 12.5$  Jy), plus review of flux densities.  
 Fesen *et al.* 1985, *ApJ*, 292, 29. Optical spectra.  
 Leahy *et al.* 1986, *MNRAS*, 220, 501. Einstein observations.  
 Odegard 1986, *ApJ*, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24', 19' and 16').  
 Esposito *et al.* 1996, *ApJ*, 461, 820. Possible associated  $\gamma$ -ray emission.  
 Biggs & Lyne 1996, *MNRAS*, 282, 691. Pulsar search.  
 Oliver *et al.* 1996, *A&A*, 315, 578. CO observations of some of surroundings.  
 Jaffe *et al.* 1997, *ApJ*, 484, L129.  $\gamma$ -ray detection.

**G206.9+2.3**

**RA:** 06<sup>h</sup>48<sup>m</sup>40<sup>s</sup>  
**Dec:** +06°26'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.5

PKS 0646+06

**Size/arcmin:** 60 × 40  
**Type:** S?

**Radio:** Diffuse source near the Monoceros Nebula.

**Optical:** Filaments detected.

**X-ray:** Possibly detected.

**References:**

- Davies & Meaburn 1978, *A&A*, 69, 443. Optical.  
 Nosek *et al.* 1981, *ApJ*, 248, 152. HEAO-1 X-ray limit.  
 Graham *et al.* 1982, *A&A*, 109, 145. Effelsberg 100-m at 2.7 GHz (4'.4:  $S = 4.1 \pm 0.6$  Jy), plus review of flux densities.  
 Rosado 1982, *RMxAA*, 5, 127. Optical.  
 Fesen *et al.* 1985, *ApJ*, 292, 29. Optical spectra.  
 Leahy *et al.* 1986, *MNRAS*, 220, 501. Einstein observations.  
 Odegard 1986, *ApJ*, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24', 19' and 16').  
 Biggs & Lyne 1996, *MNRAS*, 282, 691. Pulsar search.

**G260.4–3.4**

**RA:** 08<sup>h</sup>22<sup>m</sup>10<sup>s</sup>  
**Dec:** −43°00'

**1-GHz flux/Jy:** 130  
**Spectral index:** 0.5

Puppis A, MSH 08–44

**Size/arcmin:** 60 × 50  
**Type:** S

This remnant overlaps the Vela SNR (G263.9–3.3).

**Radio:** Angular shell, brightest to the E, poorly defined to the W.

**Optical:** Nebulosity and wisps.

**X-ray:** Brightest to the E.

**Point sources:** Central source, a possible pulsar in X-rays.

**Distance:** Association with HI gives  $2.2 \pm 0.3$  kpc; OH absorption/emission in vicinity implies 0.5 to 1.9 kpc.

**References:**

- Green 1971, *AuJPh*, 24, 773. Molonglo at 408 MHz ( $3':S=198 \pm 20$  Jy).  
 Goudis & Meaburn 1978, *A&A*, 62, 283. H $\alpha$ +N $\text{II}$ ] optical image.  
 Winkler *et al.* 1981, *ApJ*, 245, 574. X-ray spectroscopy of O lines.  
 Winkler *et al.* 1981, *ApJ*, 245, L27. X-ray spectrum.  
 Petre *et al.* 1982, *ApJ*, 258, 22. Einstein observations.  
 Milne *et al.* 1983, *MNRAS*, 204, 237. FIRST at 1415 MHz ( $50''$ ), large scale emission missing.  
 Teske & Petre 1987, *ApJ*, 318, 370. Coronal optical line emission.  
 Dubner & Arnal 1988, *A&AS*, 75, 363. HI and CO observations of surroundings.  
 Arendt *et al.* 1990, *ApJ*, 350, 266. MOST at 843 MHz ( $44'' \times 65''$ ), with large scale emission added, plus IR, optical and X-ray observations.  
 Dubner *et al.* 1991, *AJ*, 101, 1466. VLA at 327 MHz ( $43'' \times 73''$ ) and 1.5 GHz ( $43'' \times 77''$ ).  
 Arendt *et al.* 1991, *ApJ*, 368, 474. IR observations.  
 Milne *et al.* 1993, *MNRAS*, 261, 366. Parkes 64-m at 4.75 ( $4'.5:S=59 \pm 5$  Jy) and 8.4 GHz ( $3':S=38 \pm 4$  Jy), plus polarization.  
 Berthiaume *et al.* 1994, *ApJ*, 425, 132. X-ray spectroscopy.  
 Sutherland & Dopita 1995, *ApJ*, 439, 365. Spectrophotometry.  
 Reynolds *et al.* 1995, *AJ*, 110, 318. VLA at 1.4 GHz ( $90''$ ) including neutral hydrogen.  
 Blair *et al.* 1995, *ApJ*, 454, L35. Far UV spectroscopy.  
 Petre *et al.* 1996, *ApJ*, 465, L43. ROSAT of central source.  
 Bock *et al.* 1998, *AJ*, 116, 1886. MOST at 843 MHz ( $43'' \times 60''$ ).  
 Pavlov *et al.* 1999, *ApJ*, 511, L45. Possible pulsation detection from central X-ray source.  
 Zavlin *et al.* 1999, *ApJ*, 525, 959. X-ray observations of central source.  
 Bocchino *et al.* 2000, *A&A*, 359, 316. Optical studies of selected filaments in N.  
 Woermann *et al.* 2000, *MNRAS*, 317, 421. OH observations.  
 Gaensler *et al.* 2000, *ApJ*, 537, L35. Radio limit for nebula around possible pulsar.

**G261.9+5.5**

**RA:** 09<sup>h</sup>04<sup>m</sup>20<sup>s</sup>  
**Dec:** −38°42'

**1-GHz flux/Jy:** 10?  
**Spectral index:** 0.4?

**Size/arcmin:** 40 × 30  
**Type:** S

**Radio:** Faint shell with little limb brightening.

**References:**

- Hill 1967, *AuJPh*, 20, 297. Parkes 64-m at 2650 MHz ( $7'.5:S=7$  Jy) also  $S_{1410\text{ MHz}}=8$  Jy,  $S_{81.5\text{ MHz}}=25$  Jy.  
 Colomb & Dubner 1980, *A&A*, 82, 244. Argentine 30-m dish at 1.4 GHz, for HI possibly associated with remnant.  
 Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz ( $44'' \times 71''$ ).  
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

**G263.9–3.3**

**RA:** 08<sup>h</sup>34<sup>m</sup>00<sup>s</sup>  
**Dec:** −45°50'

**1-GHz flux/Jy:** 1750  
**Spectral index:** varies

**Vela (XYZ)**  
**Size/arcmin:** 255  
**Type:** C

This refers to the whole Vela XYZ complex, of which X has at times been classified as a separate (filled-centre) remnant. This remnant is overlapped by G260.4–3.4 and G266.2–1.2.

**Radio:** Large shell, with flatter spectrum component (Vela X).

**Optical:** Filaments.

**X-ray:** Patchy shell, with extensions, central nebula and pulsar.

**Point sources:** Pulsar within Vela X, with one-sided ‘jet’.

**Distance:** Vela pulsar parallax gives 0.3 kpc, optical spectra and HI studies suggest 0.25 kpc.

**References:**

- Milne 1968, AuJPh, 21, 201. Parkes 64-m at various frequencies, including 408 MHz ( $48' : S = 2300 \pm 300$  Jy), 635 MHz ( $31' : S = 2360 \pm 300$  Jy), 1410 MHz ( $14' : S = 1640 \pm 300$  Jy) and 2650 MHz ( $7'.5 : S = 1400 \pm 250$  Jy), plus discussion of the distance.  
van den Bergh *et al.* 1973, ApJS, 26, 19. Optical.  
Dopita *et al.* 1977, ApJ, 214, 179. Optical spectroscopy.  
Hearn *et al.* 1980, ApJ, 235, L67. X-ray map.  
Milne 1980, A&A, 81, 293. Maps of Vela X with Parkes 64-m at 1.66, 2.7 and 5 GHz ( $12', 8'.4$  and  $4'.4$ ).  
Weiler & Panagia 1980, A&A, 90, 269. Clarify notation of this region and review previous observations.  
Kahn *et al.* 1985, ApJ, 299, 821. Einstein observations.  
Harnden *et al.* 1985, ApJ, 299, 828. Einstein observations of Vela X and pulsar.  
Bignami & Caraveo 1988, ApJ, 325, L5. Pulsar proper motion from optical observations.  
Dwarakanath 1991, JApA, 12, 199. Gauribidanur ‘T’ array at 34.5 MHz ( $26' \times 84' : S = 1800$  Jy for Vela X and  $S = 3900$  Jy for Vela YZ), plus review of flux densities.  
Bietenholz *et al.* 1991, ApJ, 376, L41. VLA at 5 GHz in vicinity of pulsar.  
Raymond *et al.* 1991, ApJ, 383, 226. UV spectroscopy.  
Willmore *et al.* 1992, MNRAS, 254, 139. Hard X-ray observations.  
Dubner *et al.* 1992, A&AS, 96, 505. Argentine 30-m at 1.4 GHz ( $30'$ ) of surrounding HI.  
Oberlack *et al.* 1994, ApJS, 92, 433.  $\gamma$ -ray observations.  
Bocchino *et al.* 1994, ApJ, 437, 209. ROSAT observations of NE.  
Aschenbach *et al.* 1995, Natur, 373, 587. ROSAT of X-ray extensions.  
Strom *et al.* 1995, Natur, 373, 590. Radio of X-ray extensions.  
Markwardt & Ögelman 1995, Natur, 375, 40. X-ray jet from pulsar.  
Jenkins & Wallerstein 1995, ApJ, 440, 227. Optical absorption of associated neutral carbon cloud.  
Milne 1995, MNRAS, 277, 1435. Parkes 64-m at 8.4 GHz ( $3'$ ), including polarization, of Vela X.  
Duncan *et al.* 1996, MNRAS, 280, 252. Parkes 64-m at 2.4 GHz ( $8'.9$ ).  
Danks & Sembach 1995, AJ, 109, 2627. Optical spectroscopy of background stars.  
de Jager *et al.* 1996, ApJ, 460, 729.  $\gamma$ -ray detection.  
Blair *et al.* 1995, AJ, 110, 312. UV spectroscopy.  
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 62''$ ) of part.  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.  
Frail *et al.* 1997, ApJ, 475, 224. VLA at 330 MHz ( $1' \times 1'.8$ ) and comparison with ROSAT observations of Vela X.  
Markwardt & Ögelman 1997, ApJ, 480, L13. ASCA observations of pulsar ‘jet’.  
Raymond *et al.* 1997, ApJ, 482, 881. UV spectroscopy.  
Yoshikoshi *et al.* 1997, ApJ, 487, L65.  $\gamma$ -ray observations of pulsar.  
Jenkins *et al.* 1998, ApJ, 492, L147. UV absorption to background star.  
Dubner *et al.* 1998, AJ, 116, 813. Parkes 64-m at 1.4 GHz ( $15'$ ) for HI studies.  
Bock *et al.* 1998, AJ, 116, 1886. MOST at 843 MHz ( $43'' \times 60''$ ).  
Cha *et al.* 1999, ApJ, 515, L25. Optical spectra, for distance.  
Lu & Aschenbach 2000, A&A, 362, 1083. ROSAT spatially resolved spectroscopy.  
Chadwick *et al.* 2000, ApJ, 537, 414. Limit on high energy  $\gamma$ -rays from pulsar.  
Sankrit *et al.* 2001, ApJ, 549, 416. Far-UV observations of selected region.  
Pavlov *et al.* 2001, ApJ, 554, L189. Chandra two-epoch observations of pulsar nebula.  
Helfand *et al.* 2001, ApJ, 556, 380. Chandra observations of pulsar and its nebula.  
Miyata *et al.* 2001, ApJ, 559, L45. Chandra of X-ray extension.  
Alvarez *et al.* 2001, A&A, 372, 636. Radio spectral index studies.  
Caraveo *et al.* 2001, ApJ, 561, 930. HST parallax observations of pulsar.

**G266.2–1.2**

**RA:** 08<sup>h</sup>52<sup>m</sup>00<sup>s</sup>  
**Dec:** −46°20'

**1-GHz flux/Jy:** 50?  
**Spectral index:** 0.3?

**Size/arcmin:** 120  
**Type:** S

This remnant overlaps the Vela SNR (G263.9–3.3).

**Radio:** Incomplete shell, confused by the Vela SNR.

**Optical:** Possible nebulosity associated.

**X-ray:** Non-thermal shell, confused by the Vela SNR, with central source.

**Point sources:** Central X-ray source.

**Distance:** X-ray data suggest an upper limit of 1 kpc.

**References:**

- Aschenbach 1998, Natur, 396, 141. ROSAT identification.  
Iyudin *et al.* 1998, Natur, 396, 142.  $\gamma$ -ray observations.  
Combi *et al.* 1999, ApJ, 519, L177. Radio observations.  
Duncan & Green 2000, A&A, 364, 732. Parkes 64-m at 1.4 GHz (14'.0), and comparison with other observations.  
Redman *et al.* 2000, ApJ, 543, L153. Optical of nearly nebulosity.  
Slane *et al.* 2001, ApJ, 548, 814. ASCA observations.  
Mereghetti *et al.* 2001, ApJ, 548, L213. BeppoSAX observations of central sources.  
Tsunemi *et al.* 2000, PASJ, 52, 887. ASCA spectral observations.  
Pavlov *et al.* 2001, ApJ, 559, L131. Chandra of central X-ray source.

**G272.2–3.2**

**RA:** 09<sup>h</sup>06<sup>m</sup>50<sup>s</sup>  
**Dec:** −52°02'

**1-GHz flux/Jy:** 0.4  
**Spectral index:** 0.6

**Size/arcmin:** 15?  
**Type:** S?

**Radio:** Diffuse shell.

**X-ray:** Centrally brightened.

**Optical:** Detected.

**References:**

- Greiner *et al.* 1994, A&A, 286, L35. ROSAT observations, plus optical.  
Duncan *et al.* 1997, MNRAS, 289, 97. Parkes 64-m at 1.4 GHz (18' :  $S = 0.38 \pm 0.09$  Jy) 2.4 GHz (10'.6 :  $S = 0.25 \pm 0.04$  Jy) and 4.8 GHz (5'.7 :  $S = 0.17 \pm 0.02$  Jy), MOST at 843 MHz (45''  $\times$  70'' :  $S = 0.45 \pm 0.10$  Jy), and ATCA at 2.4 GHz (37''  $\times$  52''), plus ROSAT observations.  
Harrus *et al.* 2001, ApJ, 552, 614. ASCA and ROSAT observations, plus review of earlier observations.

**G279.0+1.1**

**RA:** 09<sup>h</sup>57<sup>m</sup>40<sup>s</sup>  
**Dec:** −53°15'

**1-GHz flux/Jy:** 30?  
**Spectral index:** 0.6?

**Size/arcmin:** 95  
**Type:** S

**Radio:** Faint, incomplete shell.

**Point sources:** Pulsar nearby.

**References:**

- Woermann & Jonas 1988, MNRAS, 234, 971. Hartesbeesthoek 26-m at 1.6 (30' :  $S = 25.2 \pm 4$  Jy) and 2.3 GHz (20' :  $S = 20.7 \pm 3$  Jy).  
Duncan *et al.* 1995, MNRAS, 277, 319. Parkes 64-m at 1.4 (18' :  $S = 28 \pm 3$  Jy) and 2.4 GHz (11' :  $S = 20 \pm 2$  Jy), including polarization.  
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  53'') of part.  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G284.3–1.8**

**RA:**  $10^{\text{h}}18^{\text{m}}15^{\text{s}}$   
**Dec:**  $-59^{\circ}00'$

**1-GHz flux/Jy:** 11?  
**Spectral index:** 0.3?

**MSH 10–53**

**Size/arcmin:** 24?  
**Type:** S

Has been called G284.2–1.8.

**Radio:** Incomplete, poorly defined shell.

**Point sources:** Pulsar nearby.

**References:**

- Ruiz & May 1986, ApJ, 309, 667. CO and optical observations.  
 Milne *et al.* 1989, PASAu, 8, 187. MOST at 843 MHz ( $43'' \times 50''$ ) and Parkes 64-m at 8.4 GHz ( $3' : S = 5.4 \pm 0.8$  Jy) including polarization, plus earlier flux densities.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.  
 Camilo *et al.* 2001, ApJ, 557, L51. Observations of nearby pulsar.

**G286.5–1.2**

**RA:**  $10^{\text{h}}35^{\text{m}}40^{\text{s}}$   
**Dec:**  $-59^{\circ}42'$

**1-GHz flux/Jy:** 1.4?  
**Spectral index:** ?

**Size/arcmin:**  $26 \times 6$   
**Type:** S?

**Radio:** Double, elongated arc.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 50'' : S = 1.6$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G289.7–0.3**

**RA:**  $11^{\text{h}}01^{\text{m}}15^{\text{s}}$   
**Dec:**  $-60^{\circ}18'$

**1-GHz flux/Jy:** 6.2  
**Spectral index:** 0.2?

**Size/arcmin:**  $18 \times 14$   
**Type:** S

**Radio:** Incomplete shell.

**Point sources:** Compact radio source near centre.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 50'' : S = 6.4 \pm 0.5$  Jy), plus Parkes 64-m at 4.5 GHz ( $S = 7.5 \pm 2.5$  Jy) and 8.55 GHz ( $S = 3.6 \pm 0.9$  Jy).

**G290.1–0.8**

**RA:**  $11^{\text{h}}03^{\text{m}}05^{\text{s}}$   
**Dec:**  $-60^{\circ}56'$

**1-GHz flux/Jy:** 42  
**Spectral index:** 0.4

**MSH 11–61A**

**Size/arcmin:**  $19 \times 14$   
**Type:** S

**Radio:** Elongated, clumpy shell.

**Optical:** Filaments detected.

**X-ray:** Detected.

**Point sources:** Pulsar nearby.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ( $3'$ ) and Parkes 64-m at 5 GHz ( $4'$ ).  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ( $4'.4 : S = 20.2$  Jy).  
 Elliot & Malin 1979, MNRAS, 186, 45P. Optical image and spectra.  
 Kirshner & Winkler 1979, ApJ, 227, 853. Optical.  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 50''$ ).  
 Milne *et al.* 1989, PASAu, 8, 187. MOST at 843 MHz ( $43'' \times 49'' : S = 45 \pm 11$  Jy), and Parkes 64-m at 8.4 GHz ( $3' : S = 19.5 \pm 1.0$  Jy), including polarization.  
 Seward 1990, ApJS, 73, 781. Einstein observations.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 49'' : S = 43$  Jy).  
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.  
 Kaspi *et al.* 1997, ApJ, 485, 820. Pulsar detection.  
 Gotthelf & Kaspi 1998, ApJ, 497, L29. ASCA observations of pulsar.

**G291.0–0.1** (MSH 11–62)

**RA:** 11<sup>h</sup>11<sup>m</sup>54<sup>s</sup>  
**Dec:** −60°38'

**1-GHz flux/Jy:** 16  
**Spectral index:** 0.29

**Size/arcmin:** 15 × 13  
**Type:** C

**Radio:** Centrally brightened core, with surrounding arcs.

**X-ray:** Centrally brightened.

**Point sources:** Central compact X-ray source.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Roger *et al.* 1986, MNRAS, 219, 815. MOST at 843 MHz (43'' × 50'':  $S = 17.2 \pm 1.0$  Jy), and Parkes 64-m at 5 and 8.4 GHz (4'.6 and 3':  $S = 10.4 \pm 0.4$  Jy and 9'.1 ± 0.2), with polarization.  
 Wilson 1986, ApJ, 302, 718. Einstein observations.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'':  $S = 12.7$  Jy).  
 Harrus *et al.* 1998, ApJ, 499, 273. ASCA observations.

**G292.0+1.8** MSH 11–54

**RA:** 11<sup>h</sup>24<sup>m</sup>36<sup>s</sup>  
**Dec:** −59°16'

**1-GHz flux/Jy:** 15  
**Spectral index:** 0.4

**Size/arcmin:** 12 × 8  
**Type:** C

**Radio:** Centrally brightened source surrounded by a plateau of faint emission.

**Optical:** Oxygen rich.

**X-ray:** Ring of emission, with diffuse central nebula and compact source.

**Point sources:** Compact X-ray source.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Goss *et al.* 1979, MNRAS, 188, 357. Optical spectra.  
 Lockhart *et al.* 1977, MNRAS, 179, 147. Fleurs at 1415 MHz (50'':  $S = 13.0$  Jy).  
 Clark *et al.* 1980, MNRAS, 193, 129. X-ray spectrum.  
 Tuohy *et al.* 1982, ApJ, 260, L65. Einstein observations.  
 Dopita & Tuohy 1984, ApJ, 282, 135. Optical spectra.  
 Braun *et al.* 1986, A&A, 162, 259. MOST at 843 MHz (0'.8) and IRAS.  
 Hughes & Singh 1994, ApJ, 422, 126. EXOSAT spectrum.  
 Sutherland & Dopita 1995, ApJ, 439, 365. Spectrophotometry.  
 Hughes *et al.* 2001, ApJ, 559, L153. Chandra observations, including central nebula.

**G292.2–0.5**

**RA:** 11<sup>h</sup>19<sup>m</sup>20<sup>s</sup>  
**Dec:** −61°28'

**1-GHz flux/Jy:** 7?  
**Spectral index:** 0.6?

**Size/arcmin:** 20 × 15  
**Type:** S

**Radio:** Shell.

**X-ray:** Detected.

**Point sources:** Central, young pulsar.

**References:**

- Crawford *et al.* 2001, ApJ, 554, 152. ATCA at 1.4 GHz (29'' × 25'':  $S = 5.6 \times 0.3$  Jy) and 2.5 GHz (21'' × 20'').  
 Pivovaroff *et al.* 2001, ApJ, 554, 161. ROSAT and ASCA observations.

**G293.8+0.6**

**RA:** 11<sup>h</sup>35<sup>m</sup>00<sup>s</sup>  
**Dec:** −60°54'

**1-GHz flux/Jy:** 5?  
**Spectral index:** 0.6?

**Size/arcmin:** 20  
**Type:** C

**Radio:** Central source, with faint extended plateau.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3':  $S = 9.0$  Jy) and Parkes 64-m at 5 GHz (4':  $S = 2.1$  Jy).  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 51'').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'':  $S = 2.6$  Jy).

**G294.1–0.0**

**RA:** 11<sup>h</sup>36<sup>m</sup>10<sup>s</sup>  
**Dec:** −61°38'

**1-GHz flux/Jy:** >2?  
**Spectral index:** ?

**Size/arcmin:** 40  
**Type:** S

**Radio:** Faint shell.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 49'' : S > 2$ ).

**G296.1–0.5**

**RA:** 11<sup>h</sup>51<sup>m</sup>10<sup>s</sup>  
**Dec:** −62°34'

**1-GHz flux/Jy:** 8?  
**Spectral index:** 0.6?

**Size/arcmin:** 37 × 25  
**Type:** S

Incorporates the previously catalogued remnant G296.1–0.7. Has been called G296.05–0.50.

**Radio:** Irregular shell, with nearby HII regions.

**Optical:** Detected.

**X-ray:** Detected.

**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 6.9$  Jy) and Parkes 64-m at 5 GHz ( $4' : S > 0.74$ ) poor 5-GHz map (of G296.1–0.7).  
 Longmore *et al.* 1977, MNRAS, 181, 541. Optical spectra.  
 van den Berg 1978, ApJS, 38, 119. Optical.  
 Markert *et al.* 1981, ApJ, 248, L17. Einstein observations.  
 Caswell & Barnes 1983, ApJ, 271, L55. Molonglo at 408 MHz ( $3' : S = 12.4$  Jy).  
 Bignami *et al.* 1986, ApJ, 302, 606. EXOSAT and Einstein observations.  
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 48'' : S > 2.4$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G296.5+10.0**

**RA:** 12<sup>h</sup>09<sup>m</sup>40<sup>s</sup>  
**Dec:** −52°25'

**1-GHz flux/Jy:** 48  
**Spectral index:** 0.5

**PKS 1209–51/52**

**Size/arcmin:** 90 × 65  
**Type:** S

Has been called G296.5+9.7.

**Radio:** Shell with two bright limbs.

**Optical:** Detected.

**X-ray:** Incomplete shell, with central pulsar.

**Point sources:** Central pulsar.

**References:**

Irvine & Irvine 1974, ApJ, 192, L111. Optical.  
 Danziger & Dennefeld 1976, PASP, 88, 44. Optical spectra.  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ( $8'.4$ ) and 5 GHz ( $4'.4$ ).  
 Tuohy *et al.* 1979, ApJ, 230, L27. X-ray detection with HEAO-1 A2 experiment.  
 Ruiz 1983, AJ, 88, 1210. Optical spectra.  
 Dubner *et al.* 1986, AJ, 91, 343. Argentine 30-m dish at 1.4 GHz ( $34'$ ), plus HI.  
 Kellett *et al.* 1987, MNRAS, 225, 199. EXOSAT of the west of the remnant, including the compact source.  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 56''$ ).  
 Matsui *et al.* 1988, ApJ, 329, 838. Einstein observations, including compact source.  
 Roger *et al.* 1988, ApJ, 332, 940. MOST at 843 MHz ( $44'' \times 56''$ ).  
 Bignami *et al.* 1992, ApJ, 389, L67. Optical in vicinity of X-ray source.  
 Milne & Haynes 1994, MNRAS, 270, 106. Parkes 64-m at 2.4 GHz ( $8'.3 : S = 33 \pm 3$  Jy), 4.8 GHz ( $4'.5 : S = 23.3 \pm 3$  Jy) and 8.4 GHz ( $3'.0 : 18.8 \pm 3$ ), including polarization and review of flux densities.  
 Mereghetti *et al.* 1996, ApJ, 464, 842. Radio, optical and X-ray observations of central source.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.  
 Vasisht *et al.* 1997, ApJ, 476, L43. ASCA observations of neutron star.  
 Zavlin *et al.* 1998, A&A, 331, 821. ROSAT and ASCA observations of neutron star.  
 Giacani *et al.* 2000, AJ, 119, 281. ATCA at 1.4 GHz ( $2'.7 \times 4'.0$ ) for HI studies.  
 Zavlin *et al.* 2000, ApJ, 540, L25. Chandra observations of central pulsar.

<b>G296.8–0.3</b>		1156–62
<b>RA:</b> 11 <sup>h</sup> 58 <sup>m</sup> 30 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 9	<b>Size/arcmin:</b> 20 × 14
<b>Dec:</b> −62°35'	<b>Spectral index:</b> 0.6	<b>Type:</b> S

**Radio:** Shell, brighter to the NW.

**Distance:** HI absorption gives  $9.6 \pm 0.6$  kpc.

**X-ray:** Detected.

**References:**

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 15.0$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 3.2$  Jy).  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ( $8'.4$ ) and 5 GHz ( $4'.4$ ).  
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 48'' : S = 9.2$  Jy).  
 Gaensler *et al.* 1998, MNRAS, 296, 813. ATCA at 1.3 GHz ( $22'' \times 24'' : S = 7.0 \pm 0.3$  Jy), including polarisation and HI observations, plus review of flux densities.

---

<b>G298.5–0.3</b>		
<b>RA:</b> 12 <sup>h</sup> 12 <sup>m</sup> 40 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 5?	<b>Size/arcmin:</b> 5?
<b>Dec:</b> −62°52'	<b>Spectral index:</b> 0.4?	<b>Type:</b> ?

**Radio:** Not well resolved, may be part of a larger ring?

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ( $3'$ ) and Parkes 64-m at 5 GHz ( $4'$ ).  
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 48'' : S = 1.8$  Jy).

---

<b>G298.6–0.0</b>		
<b>RA:</b> 12 <sup>h</sup> 13 <sup>m</sup> 41 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 5?	<b>Size/arcmin:</b> 12 × 9
<b>Dec:</b> −62°37'	<b>Spectral index:</b> 0.3	<b>Type:</b> S

Has been called G298.6–0.1.

**Radio:** Incomplete shell, in complex region.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ( $3'$ ) and Parkes 64-m at 5 GHz ( $4'$ ).  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 50''$ ).  
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 48'' : S = 7.4$  Jy).

---

<b>G299.2–2.9</b>		
<b>RA:</b> 12 <sup>h</sup> 15 <sup>m</sup> 13 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 0.5?	<b>Size/arcmin:</b> 18 × 11
<b>Dec:</b> −65°30'	<b>Spectral index:</b> ?	<b>Type:</b> S

**Radio:** Faint source.

**X-ray:** Centrally brightened.

**Optical:** Filaments in W.

**References:**

- Busser *et al.* 1996, A&A, 310, L1. ROSAT detection, plus optical studies.  
 Slane *et al.* 1996, ApJ, 465, 840. Einstein, IRAS and radio observations.  
 Bai & Wang 2000, ApJ, 539, 760. ASCA observations.

**G299.6–0.5**

**RA:** 12<sup>h</sup>21<sup>m</sup>45<sup>s</sup>  
**Dec:** −63°09'

**1-GHz flux/Jy:** 1.0?  
**Spectral index:** ?

**Size/arcmin:** 13  
**Type:** S

**Radio:** Faint shell, brightest to E.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'' :  $S=1.1$  Jy).

**G301.4–1.0**

**RA:** 12<sup>h</sup>37<sup>m</sup>55<sup>s</sup>  
**Dec:** −63°49'

**1-GHz flux/Jy:** 2.1?  
**Spectral index:** ?

**Size/arcmin:** 37 × 23  
**Type:** S

**Radio:** Faint, incomplete shell, with possible extensionm to southwest.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'' :  $S=2.3$  Jy).

**G302.3+0.7**

**RA:** 12<sup>h</sup>45<sup>m</sup>55<sup>s</sup>  
**Dec:** −62°08'

**1-GHz flux/Jy:** 5?  
**Spectral index:** 0.4?

**Size/arcmin:** 17  
**Type:** S

**Radio:** Distorted shell, in complex region, with possibly associated filament.

**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' :  $S=7.5$  Jy) and Parkes 64-m at 5 GHz (4' :  $S=3.0$  Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'' :  $S=3.2$  Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

**G304.6+0.1**

Kes 17

**RA:** 13<sup>h</sup>05<sup>m</sup>59<sup>s</sup>  
**Dec:** −62°42'

**1-GHz flux/Jy:** 14  
**Spectral index:** 0.5

**Size/arcmin:** 8  
**Type:** S

**Radio:** Incomplete shell.

**Distance:** Possible limit of > 9.7 kpc from HI absorption.

**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4 :  $S=6.9$  Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'' :  $S=18$  Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

**G308.1–0.7**

**RA:** 13<sup>h</sup>37<sup>m</sup>37<sup>s</sup>  
**Dec:** −63°04'

**1-GHz flux/Jy:** 1.2?  
**Spectral index:** ?

**Size/arcmin:** 13  
**Type:** S

**Radio:** Faint shell.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'' :  $S=1.3$  Jy).

**G308.8–0.1**

**RA:** 13<sup>h</sup>42<sup>m</sup>30<sup>s</sup>  
**Dec:** −62°23'

**1-GHz flux/Jy:** 15?  
**Spectral index:** 0.4?

**Size/arcmin:** 30 × 20?  
**Type:** C?

Incorporates previous catalogued remnant G308.7+0.0.

**Radio:** Bright ridge in north, and arc to south.

**Point sources:** Pulsar near centre of remnant.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 16.7$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 7.0$  Jy).  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ( $4'.4 : S = 6.5$  Jy).  
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'').  
 Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.  
 Caswell *et al.* 1992, ApJ, 399, L151. MOST at 843 MHz (43'' × 49'').  
 Kaspi *et al.* 1992, ApJ, 399, L155. Pulsar observations.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G309.2–0.6**

**RA:** 13<sup>h</sup>46<sup>m</sup>31<sup>s</sup>  
**Dec:** −62°54'

**1-GHz flux/Jy:** 7?  
**Spectral index:** 0.4?

**Size/arcmin:** 15 × 12  
**Type:** S

Has been called G309.2–0.7.

**Radio:** Distorted shell.

**X-ray:** Extended emission, with central source.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 10.0$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 3.9$  Jy).  
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1'.5).  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 50'').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'':  $S = 6$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Gaensler *et al.* 1998, MNRAS, 299, 812. ATCA at 1.3 GHz (24'' × 23'':  $S = 5.2 \pm 0.2$  Jy).  
 Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT observations.

**G309.8+0.0**

**RA:** 13<sup>h</sup>50<sup>m</sup>30<sup>s</sup>  
**Dec:** −62°05'

**1-GHz flux/Jy:** 17  
**Spectral index:** 0.5

**Size/arcmin:** 25 × 19  
**Type:** S

**Radio:** Distorted shell.

**Point sources:** Steep radio spectrum source near the centre of the remnant.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 26.4$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 7.4$  Jy).  
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (1').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'':  $S > 8.8$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G310.6–0.3**

**RA:** 13<sup>h</sup>58<sup>m</sup>00<sup>s</sup>  
**Dec:** −62°09'

**1-GHz flux/Jy:** 5?  
**Spectral index:** ?

Kes 20B  
**Size/arcmin:** 8  
**Type:** S

**Radio:** Asymmetric shell.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'':  $S = 5.4$  Jy).

---

<b>G310.8–0.4</b>		Kes 20A
<b>RA:</b> 14 <sup>h</sup> 00 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 6?	<b>Size/arcmin:</b> 12
<b>Dec:</b> −62°17'	<b>Spectral index:</b> ?	<b>Type:</b> S
<b>Radio:</b> Arc in E, in complex region.		
<b>References:</b> Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'': $S=6.9$ Jy).		
<hr/>		
<b>G311.5–0.3</b>		
<b>RA:</b> 14 <sup>h</sup> 05 <sup>m</sup> 38 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 3?	<b>Size/arcmin:</b> 5
<b>Dec:</b> −61°58'	<b>Spectral index:</b> 0.5	<b>Type:</b> S
<b>Radio:</b> Shell, not well resolved.		
<b>References:</b> Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4'). Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3'). Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'': $S=2.9$ Jy).		
<hr/>		
<b>G312.4–0.4</b>		
<b>RA:</b> 14 <sup>h</sup> 13 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 45	<b>Size/arcmin:</b> 38
<b>Dec:</b> −61°44'	<b>Spectral index:</b> 0.36	<b>Type:</b> S
<b>Radio:</b> Irregular, incomplete shell.		
<b>Point sources:</b> Nearby $\gamma$ -ray sources.		
<b>References:</b> Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3': $S=56$ Jy). Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'': $S>19$ ), plus Parkes 64-m at 4.5 GHz ( $S=30±2$ Jy) and 8.55 GHz ( $S=17±4$ Jy). Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant. Case & Bhattacharya 1998, ApJ, 521, 246. Nearby $\gamma$ -ray sources. Roberts <i>et al.</i> 1999, ApJ, 515, 712. MOST at 843 MHz (43'' × 49'').		

**G315.4–2.3**

**RA:** 14<sup>h</sup>43<sup>m</sup>00<sup>s</sup>  
**Dec:** −62°30'

**1-GHz flux/Jy:** 49  
**Spectral index:** 0.6

RCW 86, MSH 14–63

**Size/arcmin:** 42  
**Type:** S

Possibly the remnant of the SN of AD185?

**Radio:** Shell, brightest to the SW.

**Optical:** Bright, radiative filaments, with some faint Balmer dominated filaments.

**X-ray:** Partial shell and bright patch.

**Distance:** Possible association with OB stars suggests 2.5 kpc.

**References:**

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical.  
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 86$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 18.2$  Jy).  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ( $8'.4$ ) and 5 GHz ( $4'.4$ ).  
 Winkler 1978, ApJ, 221, 220. X-ray detection.  
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.  
 Pisarski *et al.* 1984, ApJ, 277, 710. Einstein observations.  
 Nugent *et al.* 1984, ApJ, 284, 612. X-ray spectrum.  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 50''$ ).  
 Long & Blair 1990, ApJ, 358, L13. Balmer dominated optical filaments.  
 Greidanus & Strom 1990, A&A, 240, 385. IRAS observations.  
 Kaastra *et al.* 1992, A&A, 264, 654. Ginga X-ray spectra.  
 Strom 1994, MNRAS, 268, L5. Historical association  
 Chin & Huang 1994, Natur, 371, 398. Questioning of historical association.  
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 48'' : S > 22$ ).  
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Smith *et al.* 1997, AJ, 114, 2664. Observations of Balmer dominated filaments.  
 Vink *et al.* 1997, A&A, 328, 628. ASCA spectroscopy.  
 Bocchino *et al.* 2000, A&A, 360, 671. ASCA observations of N and SW.  
 Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.  
 Dickel *et al.* 2001, ApJ, 546, 447. ATCA at 1.34 GHz ( $8''$ ), including polarisation.  
 Bamba *et al.* 2000, PASJ, 52, 1157. ASCA observations.  
 Bokowski *et al.* 2001, ApJ, 550, 334. ASCA observations.

**G315.4–0.3**

**RA:** 14<sup>h</sup>35<sup>m</sup>55<sup>s</sup>  
**Dec:** −60°36'

**1-GHz flux/Jy:** 8  
**Spectral index:** 0.4

**Size/arcmin:** 24 × 13  
**Type:** ?

**Radio:** Irregular non-thermal emission, with HII region superposed in E.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 15.9$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 4.9$  Jy).  
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz ( $50'' : S = 6.25$  Jy), re-assessment of earlier flux densities.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 49'' : S = 3.1$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

**G315.9–0.0**

**RA:** 14<sup>h</sup>38<sup>m</sup>25<sup>s</sup>  
**Dec:** −60°11'

**1-GHz flux/Jy:** 0.8?  
**Spectral index:** ?

**Size/arcmin:** 25 × 14  
**Type:** S

Has been called G315.8–0.0.

**Radio:** Faint, distorted shell, with steep-spectrum ‘jet’?

**References:**

- Kesteven *et al.* 1987, AuJPh, 40, 855. MOST at 843 MHz ( $44'' \times 50''$ ).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 50'' : S = 0.9$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.

**G316.3–0.0**

(MSH 14–57)

**RA:** 14<sup>h</sup>41<sup>m</sup>30<sup>s</sup>  
**Dec:** −60°00'

**1-GHz flux/Jy:** 20?  
**Spectral index:** 0.4

**Size/arcmin:** 29 × 14  
**Type:** S

**Radio:** Distorted shell, with possible ‘blowout’.

**X-ray:** Detected.

**Distance:** HI absorption data suggests > 7.2 kpc.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4 :  $S = 16.7$  Jy).  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 51'').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 50'':  $S = 20$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G317.3–0.2**

**RA:** 14<sup>h</sup>49<sup>m</sup>40<sup>s</sup>  
**Dec:** −59°46'

**1-GHz flux/Jy:** 4.7?  
**Spectral index:** ?

**Size/arcmin:** 11  
**Type:** S

**Radio:** Incomplete shell.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 50'':  $S = 5.2$  Jy).

**G318.2+0.1**

**RA:** 14<sup>h</sup>54<sup>m</sup>50<sup>s</sup>  
**Dec:** −59°04'

**1-GHz flux/Jy:** >3.9?  
**Spectral index:** ?

**Size/arcmin:** 40 × 35  
**Type:** S

**Radio:** Faint shell, with central HII region.

**X-ray:** Sources within remnant.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 50'':  $S > 4.3$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Bocchino *et al.* 2001, A&A, 367, 629. BeppoSAX and ROSAT observations.

**G318.9+0.4**

**RA:** 14<sup>h</sup>58<sup>m</sup>30<sup>s</sup>  
**Dec:** −58°29'

**1-GHz flux/Jy:** 4?  
**Spectral index:** 0.2?

**Size/arcmin:** 30 × 14  
**Type:** C

May not be a SNR?

**Radio:** Complex arcs, with off-centre core.

**References:**

- Whiteoak 1990, Natur, 347, 157. MOST at 843 MHz (43'' × 51'').  
 Whiteoak 1993, ApJ, 415, 701. MOST at 843 MHz (43'' × 51'':  $S = 4.8 \pm 0.6$  Jy), Parkes 64-m at 4.5 GHz (4'.7 :  $S = 3.7 \pm 0.2$  Jy) and 8.4 GHz (2'.8 :  $S = 3.0 \pm 0.4$  Jy) including polarization, and ATCA at 1.4 GHz and 4.8 GHz (11'' × 13'') of core.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 50'':  $S = 4.8$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G320.4–1.2**

**RA:** 15<sup>h</sup>14<sup>m</sup>30<sup>s</sup>  
**Dec:** −59°08'

**1-GHz flux/Jy:** 60?  
**Spectral index:** 0.4

MSH 15–52, RCW 89

**Size/arcmin:** 35  
**Type:** C

Has been suggested as the remnant of the SN of AD185?

**Radio:** Ragged shell.

**Optical:** RCW 89 is the H $\alpha$  emitting region to the NW.

**X-ray:** Partial shell, central nebula and pulsar and possible ‘jet’.

**Point sources:** Radio and X-ray pulsar, plus IR compact source.

**Distance:** HI absorption indicates 5.2 kpc.

**References:**

- Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).  
 Dopita *et al.* 1977, ApJ, 214, 179. Optical spectra.  
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'').  
 Seward & Harnden 1982, ApJ, 256, L45. X-ray of pulsar.  
 Manchester *et al.* 1982, ApJ, 262, L31. Radio of pulsar.  
 Seward *et al.* 1983, ApJ, 267, 698. X-ray, Optical and IR.  
 van den Bergh & Kamper, 1984, ApJ, 280, L51. Optical expansion.  
 Seward *et al.* 1984, ApJ, 281, 650. X-ray observations of pulsar and nebulosity.  
 Lortet *et al.* 1987, A&A, 180, 65. Optical observations.  
 Trussoni *et al.* 1990, A&A, 234, 403. EXOSAT observations.  
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.  
 Arendt 1991, AJ, 101, 2160. IRAS observations, including compact source.  
 Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz (4'.5 :  $S = 37 \pm 7$  Jy) and 8.4 GHz (3'.0 :  $S = 24 \pm 4$  Jy), including polarization and review of flux densities.  
 Strom 1994, MNRAS, 268, L5. Historical association.  
 Chin & Huang 1994, Natur, 371, 398. Questioning of historical association.  
 Matz *et al.* 1994, ApJ, 434, 288. X-ray observations of pulsar.  
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.  
 Du Plessis *et al.* 1995, ApJ, 453, 746. Hartesbeesthoek 26-m at 2.3, 5 and 8.5 GHz ( $S = 42, 35.6, 14.5$  Jy).  
 Greiveldinger *et al.* 1995, ApJ, 454, 855. ROSAT observations.  
 Trassoni *et al.* 1996, A&A, 306, 581. ROSAT observations.  
 Tamura *et al.* 1996, PASJ, 48, L33. ASCA observations.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  50'':  $S = 62$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Marsden *et al.* 1997, ApJ, 491, L39. X-ray spectroscopy.  
 Brazier & Becker 1997, MNRAS, 284, 335. ROSAT observations.  
 Gaensler *et al.* 1999, MNRAS, 305, 724. ATCA at 1.4 GHz (24''  $\times$  21''), plus HI observations, and 5.3 GHz (15''  $\times$  10'').  
 Sako *et al.* 2000, ApJ, 537, 422. Possible high energy  $\gamma$ -ray detection of pulsar.

**G320.6–1.6**

**RA:** 15<sup>h</sup>17<sup>m</sup>50<sup>s</sup>  
**Dec:** −59°16'

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:** 60  $\times$  30  
**Type:** S

**Radio:** Faint shell, overlapping G320.4–1.2 in W.

**References:**

- Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz (4'.5) and 8.4 GHz (3'.0), including polarization.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  50'':  $S > 9.3$ ).

**G321.9–1.1**

**RA:** 15<sup>h</sup>23<sup>m</sup>45<sup>s</sup>  
**Dec:** −58°13'

**1-GHz flux/Jy:** >3.4?  
**Spectral index:** ?

**Size/arcmin:** 28  
**Type:** S

**Radio:** Faint shell.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  51'':  $S > 3.8$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G321.9–0.3**

**RA:** 15<sup>h</sup>20<sup>m</sup>40<sup>s</sup>  
**Dec:** −57°34'

**1-GHz flux/Jy:** 13  
**Spectral index:** 0.3

**Size/arcmin:** 31 × 23  
**Type:** S

**Radio:** Shell brighter to the W, with Cir X-1 to N.

**Point sources:** Compact, probably thermal source at S edge.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 18.3$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 7.8$  Jy).  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 52''$ ).  
 Stewart *et al.* 1993, MNRAS, 261, 593. ATCA at 1.5 GHz (21'').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 51'' : S > 8.3$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G322.5–0.1**

**RA:** 15<sup>h</sup>23<sup>m</sup>23<sup>s</sup>  
**Dec:** −57°06'

**1-GHz flux/Jy:** 1.5  
**Spectral index:** 0.4

**Size/arcmin:** 15  
**Type:** C

**Radio:** Shell with central extended source.

**Point sources:** PN Pe 2-8 within boundary.

**References:**

- Whiteoak 1992, MNRAS, 256, 121. MOST at 843 MHz ( $43'' \times 51'' : S = 2.0 \pm 0.3$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 51'' : S = 1.7$  Jy), plus Parkes 64-m at 4.5 GHz ( $= 0.89 \pm 0.13$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G323.5+0.1**

**RA:** 15<sup>h</sup>28<sup>m</sup>42<sup>s</sup>  
**Dec:** −56°21'

**1-GHz flux/Jy:** 3?  
**Spectral index:** 0.4?

**Size/arcmin:** 13  
**Type:** S

**Radio:** Distorted shell, confused with thermal emission.

**Point sources:** Compact, probably thermal source near centre.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 4.2$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 1.5$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 52'' : S = 4.2$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G326.3–1.8**

**RA:**  $15^{\text{h}}53^{\text{m}}00^{\text{s}}$   
**Dec:**  $-56^{\circ}10'$

**1-GHz flux/Jy:** 145  
**Spectral index:** varies

**MSH 15–56**

**Size/arcmin:** 38  
**Type:** C

Has been called G326.2–1.7.

**Radio:** Shell, with elongated, flat-spectrum core.

**Optical:** Emission around the shell.

**X-ray:** Shell, with central extended emission.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ( $3' : S = 180$  Jy).  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz ( $0'.8 : S > 95$ ) and Parkes 64-m at 14.7 GHz ( $2'.2 : S = 69 \pm 8$  Jy).  
 van den Berg 1979, ApJ, 227, 497. Optical observations.  
 Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.  
 Dennefeld 1980, PASP, 92, 603. Optical spectra.  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 53''$ ).  
 Milne *et al.* 1989, PASAu, 8, 187. MOST at 843 MHz ( $43'' \times 52'' : S = 153 \pm 40$  Jy), and Parkes 64-m at 8.4 GHz ( $3' : S = 68 \pm 5$  Jy), including polarization.  
 Seward 1990, ApJS, 73, 781. Einstein observations.  
 Kassim *et al.* 1993, ApJ, 419, 733. ROSAT image.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 52'' : S > 130$ ).  
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Dickel *et al.* 2000, ApJ, 543, 840. ATCA at 1.34 GHz ( $8''.5 \times 6''.4 : S > 60$  Jy), 4.80 GHz ( $3''.8 : S = 25$  Jy for core only) and 8.64 GHz ( $3''.2 : S = 15$  Jy for core only).

**G327.1–1.1**

**RA:**  $15^{\text{h}}54^{\text{m}}25^{\text{s}}$   
**Dec:**  $-55^{\circ}09'$

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:** 18  
**Type:** C

**Radio:** Shell, with off-centre core.

**X-ray:** Diffuse, with core.

**References:**

- Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz ( $8'.4 : S = 10 \pm 15\%$  Jy).  
 Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 10.6$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 4.3$  Jy).  
 Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 52'' : S = 7.6$  Jy).  
 Seward *et al.* 1996, ApJ, 471, 887. ROSAT observations.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Sun *et al.* 1999, ApJ, 511, 274. ASCA and ROSAT observations.

<b>G327.4+0.4</b>		<b>Kes 27</b>
<b>RA:</b> 15 <sup>h</sup> 48 <sup>m</sup> 20 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 30?	<b>Size/arcmin:</b> 21
<b>Dec:</b> -53°49'	<b>Spectral index:</b> 0.6	<b>Type:</b> S
Has been called G327.3+0.4 and G327.3+0.5.		
<b>Radio:</b> Incomplete, multi-arc shell, brightest to the SE. <b>X-ray:</b> Diffuse, best defined to E. <b>Distance:</b> HI absorption indicates 4.3 to 5.4 kpc. <b>References:</b> Caswell <i>et al.</i> 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 58$ Jy) and Parkes 64-m at 5 GHz ( $4' : S = 12.4$ Jy). Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ( $8'.4$ ) and 5 GHz ( $4'.4$ ). Lamb & Markert 1981, ApJ, 244, 94. Einstein observations. Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 55''$ ). Milne <i>et al.</i> 1989, PASAu, 8, 187. MOST at 843 MHz ( $43'' \times 53'' : S = 32.2 \pm 6$ Jy), and Parkes 64-m at 8.4 GHz ( $3' : S = 9.4 \pm 0.8$ Jy), including polarization. Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 53'' : S = 25$ Jy). Seward <i>et al.</i> 1996, ApJ, 471, 887. ROSAT observations. Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations. McClure-Griffiths <i>et al.</i> 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ( $2'.0 \times 1'.8$ ), plus HI.		

---

<b>G327.4+1.0</b>		<b>Size/arcmin:</b> 14
<b>RA:</b> 15 <sup>h</sup> 46 <sup>m</sup> 48 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 1.9?	<b>Type:</b> S
<b>Dec:</b> -53°20'	<b>Spectral index:</b> ?	
<b>Radio:</b> Asymmetric shell. <b>References:</b> Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 54'' : S = 2.1$ Jy). Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations. McClure-Griffiths <i>et al.</i> 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ( $2'.0 \times 1'.8$ ), plus HI.		

<b>G327.6+14.6</b>		SN1006, PKS 1459–41
<b>RA:</b> 15 <sup>h</sup> 02 <sup>m</sup> 50 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 19	<b>Size/arcmin:</b> 30
<b>Dec:</b> −41°56'	<b>Spectral index:</b> 0.6	<b>Type:</b> S
This is the remnant of the SN of AD1006.		
<b>Radio:</b> Shell, with two bright arcs.		
<b>Optical:</b> Filaments to the NW, with broad H $\alpha$ component.		
<b>X-ray:</b> Thermal shell, with non-thermal limb-brightened arcs.		
<b>Point sources:</b> The background Schweizer–Middleditch star is near the middle of the remnant.		
<b>Distance:</b> Optical spectra and proper motion indicate 1.7–3.1 kpc, modelling spectra gives $1.8 \pm 0.3$ kpc.		
<b>References:</b>		
van den Bergh 1976, ApJ, 208, L17. Optical.		
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).		
Becker <i>et al.</i> 1980, ApJ, 240, L33. X-ray spectroscopy.		
Schweizer & Middleditch 1980, ApJ, 241, 1039. Possible stellar remnant.		
Pye <i>et al.</i> 1981, MNRAS, 194, 569. Einstein observations.		
Caswell <i>et al.</i> 1983, MNRAS, 204, 921. FIRST at 1415 MHz (77").		
Reynolds & Gilmore 1986, AJ, 92, 1138. VLA at 1.37 and 1.67 GHz (16" $\times$ 20").		
Kirshner <i>et al.</i> 1987, ApJ, 315, L135. Broad H $\alpha$ optical component.		
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44" $\times$ 66").		
Fesen <i>et al.</i> 1988, ApJ, 327, 164. UV absorption spectra of the Schweizer–Middleditch star.		
Roger <i>et al.</i> 1988, ApJ, 332, 940. MOST at 843 MHz (44" $\times$ 66") : $S = 17.5 \pm 1.5$ Jy).		
Long <i>et al.</i> 1988, ApJ, 333, 749. Optical proper motion for distance.		
Jones & Pye 1989, MNRAS, 238, 567. EXOSAT observations.		
Leahy <i>et al.</i> 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.		
Wu <i>et al.</i> 1993, ApJ, 416, 247. UV spectra.		
Ozaki <i>et al.</i> 1994, PASJ, 46, 367. X-ray observations.		
Reynolds & Gilmore 1993, AJ, 106, 272. VLA at 1.37 and 1.67 GHz (24"), including polarization.		
Moffett <i>et al.</i> 1993, AJ, 106, 1566. VLA at 1.37 and 1.67 GHz (10" $\times$ 14"), for proper motion studies.		
Raymond <i>et al.</i> 1995, ApJ, 454, L31. Far UV spectroscopy.		
Koyama <i>et al.</i> 1995, Natur, 378, 255. ASCA observations.		
Willingale <i>et al.</i> 1996, MNRAS, 278, 749. ROSAT observations.		
Blair <i>et al.</i> 1996, ApJ, 468, 871. Optical absorption studies.		
Laming <i>et al.</i> 1996, ApJ, 472, 267. Modelling of optical spectra, including distance.		
Wu <i>et al.</i> 1997, ApJ, 477, L53. Far UV absorption spectra of the Schweizer–Middleditch star.		
Winkler & Long 1997, ApJ, 486, L137. UV absorption spectra of background quasar.		
Winkler & Long 1997, ApJ, 491, 829. ROSAT and optical images.		
Tanimori <i>et al.</i> 1998, ApJ, 497, L25. $\gamma$ -ray detection.		
Vink <i>et al.</i> 2000, A&A, 354, 931. X-ray spectroscopy.		
Burleigh <i>et al.</i> 2000, A&A, 356, 585. Optical spectroscopy of the Schweizer–Middleditch star.		
Allen <i>et al.</i> 2001, ApJ, 558, 739. ASCA, ROSAT and other X-ray observations.		

<b>G328.4+0.2</b>		(MSH 15–57)
<b>RA:</b> 15 <sup>h</sup> 55 <sup>m</sup> 30 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 15	<b>Size/arcmin:</b> 5
<b>Dec:</b> −53°17'	<b>Spectral index:</b> 0.12	<b>Type:</b> F
<b>Radio:</b> Amorphous emission, with central bar.		
<b>X-ray:</b> Detected at high energies.		
<b>Distance:</b> HI absorption indicates at least $17.4 \pm 0.9$ kpc.		
<b>References:</b>		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Caswell <i>et al.</i> 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50").		
Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43" $\times$ 54") : $S = 15$ Jy).		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.		
Gaensler <i>et al.</i> 2000, ApJ, 542, 380. ATCA at 1.4 GHz (19'.5 $\times$ 16''.5 : $S = 14.3 \pm 0.1$ Jy) and 4.5 GHz (2''.0 $\times$ 1''.5 : $S = 12.5 \pm 0.2$ Jy).		
Hughes <i>et al.</i> 2000, ApJ, 542, 386. ASCA observations.		
McClure-Griffiths <i>et al.</i> 2001, ApJ, 551, 394. ATCA and Parks 64-m at 1.4 GHz (2'.0 $\times$ 1'.8), plus HI.		

**G329.7+0.4**

**RA:** 16<sup>h</sup>01<sup>m</sup>20<sup>s</sup>  
**Dec:** −52°18'

**1-GHz flux/Jy:** >34?  
**Spectral index:** ?

**Size/arcmin:** 40 × 33  
**Type:** S

**Radio:** Diffuse shell, in complex region.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 53'' : S > 38$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parks 64-m at 1.4 GHz ( $2'.0 \times 1'.8$ ), plus HI.

**G330.0+15.0**

**RA:** 15<sup>h</sup>10<sup>m</sup>00<sup>s</sup>  
**Dec:** −40°00'

**1-GHz flux/Jy:** 350?  
**Spectral index:** 0.5?

Lupus Loop  
**Size/arcmin:** 180?  
**Type:** S

**Radio:** Low surface brightness loop with HI shell.

**X-ray:** Detected.

**References:**

Milne 1971, AuJPh, 24, 757. Parkes 64-m at 408 MHz (48'), 635 MHz (31') and 1410 MHz (15').  
 Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (8'.4 :  $S = 120 \pm 30$ % Jy).  
 Toor 1980, A&A, 85, 184. X-ray image and spectrum.  
 Colomb & Dubner 1982, A&A, 112, 141. Argentine 30-m dish at 1.42 GHz (30'), HI observations.  
 Leahy *et al.* 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.  
 Ozaki *et al.* 1994, PASJ, 46, 367. X-ray observations.

**G330.2+1.0**

**RA:** 16<sup>h</sup>01<sup>m</sup>06<sup>s</sup>  
**Dec:** −51°34'

**1-GHz flux/Jy:** 5?  
**Spectral index:** 0.3

**Size/arcmin:** 11  
**Type:** S?

**Radio:** Clumpy non-thermal emission, possibly a distorted shell.

**Distance:** HI absorption indicates at least 4.9 kpc.

**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 8.6$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 4.0$  Jy).  
 Caswell *et al.* 1983, MNRAS, 204, 915. FIRST at 1415 MHz ( $52'' \times 47''$ ), and MOST at 843 MHz ( $43'' \times 55''$ ).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 55'' : S = 4.7$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parks 64-m at 1.4 GHz ( $2'.0 \times 1'.8$ ), plus HI.

**G332.0+0.2**

**RA:** 16<sup>h</sup>13<sup>m</sup>17<sup>s</sup>  
**Dec:** −50°53'

**1-GHz flux/Jy:** 8?  
**Spectral index:** 0.5

**Size/arcmin:** 12  
**Type:** S

**Radio:** Incomplete shell.

**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ( $44'' \times 57''$ ).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 55'' : S = 8.9$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parks 64-m at 1.4 GHz ( $2'.0 \times 1'.8$ ), plus HI.

<b>G332.4–0.4</b>		RCW 103
<b>RA:</b> 16 <sup>h</sup> 17 <sup>m</sup> 33 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 28	<b>Size/arcmin:</b> 10
<b>Dec:</b> −51°02'	<b>Spectral index:</b> 0.5	<b>Type:</b> S
<b>Radio:</b> Shell, brightest to the S.		
<b>Optical:</b> Filaments correspond well to the radio shell.		
<b>X-ray:</b> Detected, with point source near centre.		
<b>Point sources:</b> Central, variable X-ray source, and nearby pulsar.		
<b>Distance:</b> HI absorption indicates 3.3 kpc.		
<b>References:</b>		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Caswell <i>et al.</i> 1975, A&A, 45, 239. Parkes HI absorption.		
Tuohy & Garmire 1980, ApJ, 239, L107. Einstein detection of X-ray point source.		
Caswell <i>et al.</i> 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50'').		
Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.		
Ruiz 1983, AJ, 88, 1210. Optical spectra.		
Tuohy <i>et al.</i> 1983, ApJ, 268, 778. X-ray observations.		
Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.		
Nugent <i>et al.</i> 1984, ApJ, 284, 612. X-ray spectrum (and Einstein image from Tuohy, private communication).		
Meaburn & Allen 1986, MNRAS, 222, 593. Optical spectra.		
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 57'').		
Oliva <i>et al.</i> 1990, A&A, 240, 453. IR spectroscopy.		
Dickel <i>et al.</i> 1996, AJ, 111, 340. ATCA at 1.36 (8'') and 2.37 GHz (4''.5''), including polarization.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 55'': S=34 Jy).		
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant.		
Gotthelf <i>et al.</i> 1997, ApJ, 487, L175. ASCA observations of compact X-ray source.		
Torii <i>et al.</i> 1998, ApJ, 494, L207. ASCA detection of nearby pulsar.		
Kaspi <i>et al.</i> 1998, ApJ, 503, L161. Pulsar observations.		
Gotthelf <i>et al.</i> 1999, ApJ, 514, L107. X-ray variability of central source.		
Oliva <i>et al.</i> 1999, A&A, 343, 943. ISO spectroscopy.		
Torii <i>et al.</i> 2000, ApJ, 534, L71. X-ray timing on pulsar, including glitch.		

<b>G332.4+0.1</b>		MSH 16–51, Kes 32
<b>RA:</b> 16 <sup>h</sup> 15 <sup>m</sup> 17 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 26	<b>Size/arcmin:</b> 15
<b>Dec:</b> −50°42'	<b>Spectral index:</b> 0.5	<b>Type:</b> S
Has been called G332.4+0.2.		
<b>Radio:</b> Distorted shell, with thermal jet and plume adjacent.		
<b>Point sources:</b> Pulsar nearby.		
<b>References:</b>		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Roger <i>et al.</i> 1985, Natur, 316, 44. MOST at 843 MHz (44'' × 57'').		
Caraveo 1993, ApJ, 415, L111. Nearby pulsar.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 56'': S=29 Jy).		
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant.		
Brinkmann <i>et al.</i> 1999, A&A, 346, 599. ROSAT image of remnant and nearby pulsar.		

<b>G335.2+0.1</b>		
<b>RA:</b> 16 <sup>h</sup> 27 <sup>m</sup> 45 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 16	<b>Size/arcmin:</b> 21
<b>Dec:</b> −48°47'	<b>Spectral index:</b> 0.5	<b>Type:</b> S
<b>Radio:</b> Well defined shell.		
<b>Point sources:</b> Old pulsar within remnant boundary.		
<b>References:</b>		
Clark <i>et al.</i> 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S=27.1 Jy) and Parkes 64-m at 5 GHz (4': S=8.6 Jy).		
Kaspi <i>et al.</i> 1996, AJ, 111, 2028. Pulsar observations.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 57'': S=16 Jy).		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.		

**G336.7+0.5**

**RA:** 16<sup>h</sup>32<sup>m</sup>11<sup>s</sup>  
**Dec:** -47°19'

**1-GHz flux/Jy:** 6  
**Spectral index:** 0.5

**Size/arcmin:** 14 × 10  
**Type:** S

**Radio:** Irregular shell.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 60'').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 58''):  $S = 6.1$  Jy.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G337.0-0.1**

(CTB 33)

**RA:** 16<sup>h</sup>35<sup>m</sup>57<sup>s</sup>  
**Dec:** -47°36'

**1-GHz flux/Jy:** 1.5  
**Spectral index:** 0.6?

**Size/arcmin:** 1.5  
**Type:** S

This entry refers to a small (1'.5) SNR, not the larger previously catalogued G337.0-0.1.

**Radio:** Shell, in a complex region.

**Distance:** Association with CTB 33 gives 11 kpc.

**Point sources:** Associated with a soft gamma repeater.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 58''):  $S = 21$  Jy.  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.  
 Sarma *et al.* 1997, ApJ, 483, 335. ATCA at 1.4 GHz (12'') including H<sub>i</sub>, and 8.9 GHz (13'' × 15'') for recombination lines, clarifying extent of the remnant.  
 Woods *et al.* 1999, ApJ, 519, L139. Soft gamma repeater observations.  
 Hurley *et al.* 2000, ApJ, 528, L21. ASCA observations of soft gamma repeater.  
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

**G337.2-0.7**

**RA:** 16<sup>h</sup>39<sup>m</sup>28<sup>s</sup>  
**Dec:** -47°51'

**1-GHz flux/Jy:** 2?  
**Spectral index:** 0.7

**Size/arcmin:** 6  
**Type:** S

**Radio:** Shell, not well resolved.

**X-ray:** Extended emission.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3':  $S = 3.8$  Jy) and Parkes 64-m at 5 GHz (4':  $S = 0.70$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 58''):  $S = 2.0$  Jy.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT observations.

**G337.3+1.0**

Kes 40

**RA:** 16<sup>h</sup>32<sup>m</sup>39<sup>s</sup>  
**Dec:** -46°36'

**1-GHz flux/Jy:** 16  
**Spectral index:** 0.55

**Size/arcmin:** 15 × 12  
**Type:** S

**Radio:** Nearly complete shell.

**References:**

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3':  $S = 24.6$  Jy) and Parkes 64-m at 5 GHz (4':  $S = 7.2$  Jy).  
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).  
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 61'').  
 Milne *et al.* 1989, PASAu, 8, 187. MOST at 843 MHz (43'' × 59''):  $S = 14.8 \pm 3.0$  Jy), and Parkes 64-m at 8.4 GHz (3':  $S = 5.1 \pm 0.6$  Jy), including polarization.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59''):  $S = 20$  Jy.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

---

<b>G337.8–0.1</b>		Kes 41
<b>RA:</b> 16 <sup>h</sup> 39 <sup>m</sup> 01 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 18	<b>Size/arcmin:</b> 9 × 6
<b>Dec:</b> −46°59'	<b>Spectral index:</b> 0.5	<b>Type:</b> S

**Radio:** Distorted shell.

**Distance:** HI absorption suggests > 9.3 kpc.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'':  $S = 18$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.

---

<b>G338.1+0.4</b>		
<b>RA:</b> 16 <sup>h</sup> 37 <sup>m</sup> 59 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 4?	<b>Size/arcmin:</b> 15?
<b>Dec:</b> −46°24'	<b>Spectral index:</b> 0.4	<b>Type:</b> S

**Radio:** Arc in NE, merging with thermal emission in S.

**Optical:** Detected.

**X-ray:** Detected.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'':  $S = 3.8$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

---

<b>G338.3–0.0</b>		
<b>RA:</b> 16 <sup>h</sup> 41 <sup>m</sup> 00 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 7?	<b>Size/arcmin:</b> 8
<b>Dec:</b> −46°34'	<b>Spectral index:</b> ?	<b>Type:</b> S

**Radio:** Irregular shell, in complex region.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'':  $S = 7.4$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

---

<b>G338.5+0.1</b>		
<b>RA:</b> 16 <sup>h</sup> 41 <sup>m</sup> 09 <sup>s</sup>	<b>1-GHz flux/Jy:</b> 12?	<b>Size/arcmin:</b> 9
<b>Dec:</b> −46°19'	<b>Spectral index:</b> ?	<b>Type:</b> ?

**Radio:** Circle of non-thermal emission in complex region, not well defined.

**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'':  $S = 13$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G340.4+0.4****RA:** 16<sup>h</sup>46<sup>m</sup>31<sup>s</sup>  
**Dec:** -44°39'**1-GHz flux/Jy:** 5  
**Spectral index:** 0.4**Size/arcmin:** 10 × 7  
**Type:** S**Radio:** Distorted shell, elongated east–west.**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 8.2$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 2.9$  Jy).  
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50'').  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz ( $75'' \times 48'' : S = 9.8 \pm 0.9$  Jy) and 1.4 GHz ( $27'' \times 9'' : S = 3.6 \pm 0.1$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 61'' : S = 5.9$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G340.6+0.3****RA:** 16<sup>h</sup>47<sup>m</sup>41<sup>s</sup>  
**Dec:** -44°34'**1-GHz flux/Jy:** 5?  
**Spectral index:** 0.4?**Size/arcmin:** 6  
**Type:** S**Radio:** Incomplete shell.**Optical:** Possible associated filaments.**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 7.0$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 2.8$  Jy).  
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50'').  
 Zealey *et al.* 1979, A&AS, 38, 39. Optical.  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz ( $75'' \times 48'' : S = 9.2 \pm 0.9$  Jy) and 1.4 GHz ( $27'' \times 9'' : S = 5.8 \pm 0.1$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 61'' : S = 4.5$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G341.2+0.9****RA:** 16<sup>h</sup>47<sup>m</sup>35<sup>s</sup>  
**Dec:** -43°47'**1-GHz flux/Jy:** 1.5?  
**Spectral index:** 0.6?**Size/arcmin:** 16 × 22  
**Type:** C**Radio:** Incomplete shell, with extension to SW.**Point sources:** Pulsar in W, with wind nebula.**References:**

- Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz ( $54'' \times 116'' : S = 3.0 \pm 0.1$  Jy) and 1.4 GHz ( $21'' \times 25'' : S = 12.5 \pm 0.05$  Jy).  
 Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4 and 4.9 GHz (25'') of pulsar wind nebula.

**G341.9-0.3****RA:** 16<sup>h</sup>55<sup>m</sup>01<sup>s</sup>  
**Dec:** -44°01'**1-GHz flux/Jy:** 2.5  
**Spectral index:** 0.5**Size/arcmin:** 7  
**Type:** S**Radio:** Incomplete shell, brightest to NE.**References:**

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ( $3' : S = 7.4$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 1.7$  Jy), see Caswell & Clark 1975, AuJPA, 37, 57 for the maps.  
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50''), revision of previous flux densities.  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ( $35'' \times 11'' : S = 2.2 \pm 0.1$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 62'' : S = 2.7$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G342.0–0.2**

**RA:** 16<sup>h</sup>54<sup>m</sup>50<sup>s</sup>  
**Dec:** −43°53'

**1-GHz flux/Jy:** 3.5?  
**Spectral index:** 0.4?

**Size/arcmin:** 12 × 9  
**Type:** S

**Radio:** Distorted shell.

**References:**

- Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50''), estimate  $S_{408 \text{ MHz}} = 5 \text{ Jy}$ ,  $S_{5 \text{ GHz}} = 2 \text{ Jy}$  from previous maps.  
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (35'' × 11'').  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 62'':  $S = 3.5 \text{ Jy}$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G342.1+0.9**

**RA:** 16<sup>h</sup>50<sup>m</sup>43<sup>s</sup>  
**Dec:** −43°04'

**1-GHz flux/Jy:** 0.5?  
**Spectral index:** ?

**Size/arcmin:** 10 × 9  
**Type:** S

**Radio:** Incomplete shell.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 63'':  $S = 0.6 \text{ Jy}$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G343.0–6.0**

**RA:** 17<sup>h</sup>25<sup>m</sup>00<sup>s</sup>  
**Dec:** −46°30'

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:** 250  
**Type:** S

**Radio:** Faint, poorly defined.

**Optical:** Filamentary shell.

**References:**

- Walker & Zealey 2001, MNRAS, 325, 287. Optical observations, and review of earlier observations.

**G343.1–2.3**

**RA:** 17<sup>h</sup>08<sup>m</sup>00<sup>s</sup>  
**Dec:** −44°16'

**1-GHz flux/Jy:** 8?  
**Spectral index:** 0.5?

**Size/arcmin:** 32?  
**Type:** C?

**Radio:** Incomplete shell?

**Point sources:** Pulsar near edge, with wind nebula.

**References:**

- McAdam *et al.* 1993, Natur, 361, 516. MOST at 843 MHz (smoothed to 2').  
 Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (56'' × 11'':  $S = 10.6 \text{ Jy}$ ) and 1.4 GHz (22'' × 27'') near pulsar.  
 Becker *et al.* 1995, A&A, 298, 528. ROSAT of pulsar, and limit for remnant.  
 Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4, 4.9 and 8.5 GHz (25'') of pulsar wind nebula.

**G343.1–0.7**

**RA:** 17<sup>h</sup>00<sup>m</sup>25<sup>s</sup>  
**Dec:** −43°14'

**1-GHz flux/Jy:** 7.8  
**Spectral index:** 0.55

**Size/arcmin:** 27 × 21  
**Type:** S

**Radio:** Shell, with smaller thermal shell adjacent.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 63'':  $S = 8.5 \pm 0.6 \text{ Jy}$ ), plus Parkes 64-m at 4.5 GHz ( $S = 3.9 \pm 0.6 \text{ Jy}$ ) and 8.55 GHz ( $S = 2.4 \pm 0.5 \text{ Jy}$ ).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G344.7–0.1**

**RA:** 17<sup>h</sup>03<sup>m</sup>51<sup>s</sup>  
**Dec:** −41°42'

**1-GHz flux/Jy:** 2.5?  
**Spectral index:** 0.5

**Size/arcmin:** 10  
**Type:** C?

**Radio:** Asymmetric shell, with possible core.

**X-ray:** Detected.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 4.7$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 1.3$  Jy).  
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ( $30'' \times 43'' : S = 1.7 \pm 0.1$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 65'' : S = 2.5$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G345.7–0.2**

**RA:** 17<sup>h</sup>07<sup>m</sup>20<sup>s</sup>  
**Dec:** −40°53'

**1-GHz flux/Jy:** 0.6?  
**Spectral index:** ?

**Size/arcmin:** 6  
**Type:** S

**Radio:** Poorly defined diffuse shell.

**Point sources:** Old pulsar nearby.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 66'' : S = 0.7$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G346.6–0.2**

**RA:** 17<sup>h</sup>10<sup>m</sup>19<sup>s</sup>  
**Dec:** −40°11'

**1-GHz flux/Jy:** 8?  
**Spectral index:** 0.5?

**Size/arcmin:** 8  
**Type:** S

**Radio:** Irregular shell.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S = 14.9$  Jy) and Parkes 64-m at 5 GHz ( $4' : S = 4.3$  Jy).  
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ( $31'' \times 43'' : S = 8.1 \pm 0.9$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 67'' : S = 8.7$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.

**G347.3–0.5**

**RA:** 17<sup>h</sup>13<sup>m</sup>50<sup>s</sup>  
**Dec:** −39°45'

**1-GHz flux/Jy:** ?  
**Spectral index:** ?

**Size/arcmin:**  $65 \times 55$   
**Type:** S?

**Radio:** Faint emission.

**X-ray:** Non-thermal, limb-brightened to W, with central source.

**Distance:** Association with molecular clouds, and HII region, suggests 6 kpc.

**References:**

- Koyama *et al.* 1997, PASJ, 49, L7. ASCA of NW.  
 Slane *et al.* 1999, ApJ, 525, 357. ASCA and other observations.  
 Muraishi *et al.* 2000, A&A, 354, L57. High energy  $\gamma$ -ray detection.  
 Butt *et al.* 2001, ApJ, 562, L167. Associated  $\gamma$ -ray emission.

**G348.5–0.0**

**RA:**  $17^{\text{h}}15^{\text{m}}26^{\text{s}}$   
**Dec:**  $-38^{\circ}28'$

**1-GHz flux/Jy:** 10?  
**Spectral index:** 0.4?

**Size/arcmin:** 10?  
**Type:** S?

**Radio:** Arc, overlapping G348.5+0.1.

**References:**

- Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz ( $46'' \times 53''$ ), 1.4 GHz ( $18'' \times 33''$ ) and part at 5 GHz ( $2''.5 \times 3''.9$ ).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 69''$ :  $S=10.2$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.  
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G348.5+0.1**

**RA:**  $17^{\text{h}}14^{\text{m}}06^{\text{s}}$   
**Dec:**  $-38^{\circ}32'$

**1-GHz flux/Jy:** 72  
**Spectral index:** 0.3

CTB 37A

**Size/arcmin:** 15  
**Type:** S

**Radio:** Shell, poorly define to S and W, overlapping G348.5–0.0 in E.

**Distance:** HI absorption indicates  $10.2 \pm 3.5$  kpc.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ( $3' : S=97$  Jy).  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ( $4'.4 : S=43$  Jy).  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz ( $0'.8 : S>50$ ) and Parkes 64-m at 14.7 GHz ( $2'.2 : S=18 \pm 5$  Jy).  
 Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz ( $20'' \times 45''$ ).  
 Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz ( $46'' \times 53''$ ), 1.4 GHz ( $18'' \times 33''$ ) and part at 5 GHz ( $2''.5 \times 3''.9$ ).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 69''$ :  $S=71$  Jy).  
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

**G348.7+0.3**

**RA:**  $17^{\text{h}}13^{\text{m}}55^{\text{s}}$   
**Dec:**  $-38^{\circ}11'$

**1-GHz flux/Jy:** 26  
**Spectral index:** 0.3

CTB 37B

**Size/arcmin:** 17?  
**Type:** S

**Radio:** Incomplete shell with faint eastern extensions.

**Distance:** HI absorption indicates  $10.2 \pm 3.5$  kpc.

**References:**

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ( $3' : S=34$  Jy).  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ( $4'.4 : S=32$  Jy).  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz ( $0'.8 : S>20$ ) and Parkes 64-m at 14.7 GHz ( $2'.2 : S=8 \pm 3$  Jy).  
 Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz ( $20'' \times 45''$ ).  
 Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz ( $46'' \times 53''$ ).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 69''$ :  $S=33$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

**G349.2–0.1**

**RA:**  $17^{\text{h}}17^{\text{m}}15^{\text{s}}$   
**Dec:**  $-38^{\circ}04'$

**1-GHz flux/Jy:** 1.4?  
**Spectral index:** ?

**Size/arcmin:**  $9 \times 6$   
**Type:** S

**Radio:** Elongated shell, adjacent to bright HII region.

**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ( $43'' \times 70''$ :  $S=1.6$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

**G349.7+0.2****RA:** 17<sup>h</sup>17<sup>m</sup>59<sup>s</sup>  
**Dec:** −37°26'**1-GHz flux/Jy:** 20  
**Spectral index:** 0.5**Size/arcmin:** 2.5 × 2  
**Type:** S**Radio:** Incomplete clumpy shell, with enhancement to the S.**Distance:** HI absorption indicates  $18.3 \pm 4.6$  kpc, association with CO features gives 23 kpc.**References:**

- Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').  
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' :  $S=31.0$  Jy) and Parkes 64-m at 5 GHz (4' :  $S=9.1$  Jy), no maps.  
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.  
 Shaver *et al.* 1985, Natur, 313, 113. VLA at 1.4 GHz (3''  $\times$  14''.5).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  71'' :  $S=22$  Jy).  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.  
 Reynoso & Mungum 2001, AJ, 121, 347. CO observations of the vicinity.  
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.  
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

**G350.0–2.0****RA:** 17<sup>h</sup>27<sup>m</sup>50<sup>s</sup>  
**Dec:** −38°32'**1-GHz flux/Jy:** 26  
**Spectral index:** 0.4**Size/arcmin:** 45  
**Type:** S

Incorporates the previously catalogued G350.0–1.8 in the NW.

**Radio:** Shell, brightest in NW.**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' :  $S=49.5$  Jy) and Parkes 64-m at 5 GHz (4' :  $S=13.6$  Jy).  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4).  
 Gaensler 1998, ApJ, 493, 781. VLA and Parkes 64-m at 1.4 GHz (18''  $\times$  21'' :  $S=22.3 \pm 0.3$  Jy), clarifying extent of remnant.

**G351.2+0.1****RA:** 17<sup>h</sup>22<sup>m</sup>27<sup>s</sup>  
**Dec:** −36°11'**1-GHz flux/Jy:** 5?  
**Spectral index:** 0.4**Size/arcmin:** 7  
**Type:** C?

Has been called G351.3+0.2.

**Radio:** Distorted shell, with possible flat-spectrum core.**References:**

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' :  $S=8.1$  Jy) and Parkes 64-m at 5 GHz (4' :  $S=3.1$  Jy).  
 Caswell *et al.* 1984, PASAu, 5, 227. MOST at 843 MHz (43''  $\times$  77'').  
 Becker & Helfand 1988, AJ, 95, 883. VLA at 5 GHz (15''), and at 15 GHz of core.  
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (32''  $\times$  36'' :  $S=4.8 \pm 0.2$  Jy).  
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  73'' :  $S=5.5$  Jy).  
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G351.7+0.8****RA:** 17<sup>h</sup>21<sup>m</sup>00<sup>s</sup>  
**Dec:** −35°27'**1-GHz flux/Jy:** 10?  
**Spectral index:** ?**Size/arcmin:** 18 × 14  
**Type:** S**Radio:** Elongated shell, adjacent to bright HII region.**Point sources:** Pulsar nearby.**References:**

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''  $\times$  74'' :  $S=11$  Jy).

**G351.9–0.9**

**RA:** 17<sup>h</sup>28<sup>m</sup>52<sup>s</sup>  
**Dec:** −36°16'

**1-GHz flux/Jy:** 1.8?  
**Spectral index:** ?

**Size/arcmin:** 12 × 9  
**Type:** S

**Radio:** Asymmetric shell.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 73'':  $S=2.0$  Jy).

**G352.7–0.1**

**RA:** 17<sup>h</sup>27<sup>m</sup>40<sup>s</sup>  
**Dec:** −35°07'

**1-GHz flux/Jy:** 4  
**Spectral index:** 0.6

**Size/arcmin:** 8 × 6  
**Type:** S

**Radio:** Distorted shell.

**X-ray:** Detected.

**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3':  $S=9.6$  Jy) and Parkes 64-m at 5 GHz (4':  $S=2.3$  Jy).

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (1'.1).

Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (34'':  $S=3.4 \pm 0.4$  Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 75'':  $S=4.4$  Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Kinugasa *et al.* 1998, PASJ, 50, 249. ASCA observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G353.9–2.0**

**RA:** 17<sup>h</sup>38<sup>m</sup>55<sup>s</sup>  
**Dec:** −35°11'

**1-GHz flux/Jy:** 1?  
**Spectral index:** 0.5?

**Size/arcmin:** 13  
**Type:** S

**Radio:** Shell, with central double source.

**References:**

Green 2001, MNRAS, 326, 283. VLA at 327 MHz (3'.0 × 2'.7) and 1.4 GHz (42'' × 36''), plus 8.4 GHz (8''.4 × 6''.1) of central source only.

**G354.1+0.1**

**RA:** 17<sup>h</sup>30<sup>m</sup>28<sup>s</sup>  
**Dec:** −33°46'

**1-GHz flux/Jy:** ?  
**Spectral index:** varies?

**Size/arcmin:** 15 × 3?  
**Type:** C?

Is this a SNR?

**Radio:** Elongated N–S.

**Point sources:** Pulsar at S tip.

**References:**

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (47'' × 99'') and 1.4 GHz (8''.8 × 21'').

**G354.8–0.8**

**RA:** 17<sup>h</sup>36<sup>m</sup>00<sup>s</sup>  
**Dec:** −33°42'

**1-GHz flux/Jy:** 2.8?  
**Spectral index:** ?

**Size/arcmin:** 19  
**Type:** S

**Radio:** Distorted shell.

**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 78'':  $S=3.1$  Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

**G355.6–0.0**

**RA:** 17<sup>h</sup>35<sup>m</sup>16<sup>s</sup>  
**Dec:** −32°38'

**1-GHz flux/Jy:** 3?  
**Spectral index:** ?

**Size/arcmin:** 8 × 6  
**Type:** S

**Radio:** Well defined shell.

**X-ray:** Detected.

**References:**

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ( $43'' \times 80''$ :  $S=2.6$  Jy).  
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

**G355.9–2.5**

**RA:** 17<sup>h</sup>45<sup>m</sup>53<sup>s</sup>  
**Dec:** −33°43'

**1-GHz flux/Jy:** 8  
**Spectral index:** 0.5

**Size/arcmin:** 13  
**Type:** S

**Radio:** Distorted shell, brightest to SE.

**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ( $3' : S=12.3$  Jy) and Parkes 64-m at 5 GHz ( $4' : S=3.4$  Jy).  
Caswell *et al.* 1984, PASAu, 5, 227. MOST at 843 MHz ( $43'' \times 77''$ ).  
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ( $32'' \times 34''$ :  $S=5.0 \pm 0.3$  Jy).  
Gray 1994, MNRAS, 270, 836. MOST at 843 MHz ( $43'' \times 77''$ ).  
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

**G356.2+4.5**

**RA:** 17<sup>h</sup>19<sup>m</sup>00<sup>s</sup>  
**Dec:** −29°40'

**1-GHz flux/Jy:** 4  
**Spectral index:** 0.7

**Size/arcmin:** 25  
**Type:** S

**Radio:** Faint shell.

**References:**

Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz ( $3' \times 1'.5 : S=8.1 \pm 1.7$  Jy), and NVSS at 1.4 GHz.

**G356.3–0.3**

**RA:** 17<sup>h</sup>37<sup>m</sup>56<sup>s</sup>  
**Dec:** −32°16'

**1-GHz flux/Jy:** 3?  
**Spectral index:** ?

**Size/arcmin:** 11 × 7  
**Type:** S

**Radio:** Elongated shell, brighter in N.

**References:**

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ( $43'' \times 81'' : S=2.6$  Jy).

**G356.3–1.5**

**RA:** 17<sup>h</sup>42<sup>m</sup>35<sup>s</sup>  
**Dec:** −32°52'

**1-GHz flux/Jy:** 3?  
**Spectral index:** ?

**Size/arcmin:** 20 × 15  
**Type:** S

**Radio:** Double arc.

**References:**

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ( $43'' \times 79'' : S=2.8$  Jy).

**G357.7–0.1**

**RA:** 17<sup>h</sup>40<sup>m</sup>29<sup>s</sup>  
**Dec:** −30°58'

**1-GHz flux/Jy:** 37  
**Spectral index:** 0.4

**MSH 17–39**  
**Size/arcmin:** 8 × 3?  
**Type:** ?

Has been suggested that this is not a SNR.

**Radio:** Multiple arcs and filaments.

**Point sources:** Peripheral compact source (possibly a compact HII region).

**References:**

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' :  $S = 54.2$  Jy) and Parkes 64-m at 5 GHz (4' :  $S = 18.5$  Jy).  
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4 :  $S = 14.6$  Jy).  
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).  
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50'').  
 Weiler & Panagia 1980, A&A, 90, 269. Effelsberg 100-m at 9 GHz (1'.5) (private communication from Baker).  
 Shaver *et al.* 1985, Natur, 313, 113. VLA at 1.4 GHz (3''.8 × 10''.9) and 5 GHz (12'' × 26'').  
 Becker & Helfand 1985, Natur, 313, 115. VLA at 1.4 GHz and 5 GHz, suggest it is not a SNR, see Helfand & Becker 1985 Natur, 313, 118.  
 Shaver *et al.* 1985, A&A, 147, L23. Observations of peripheral compact source.  
 Caswell *et al.* 1989, PASAu, 8, 184. MOST at 843 MHz (43'' × 83'').  
 Gray 1994, MNRAS, 270, 836. MOST at 843 MHz (43'' × 84'').  
 Stewart *et al.* 1994, ApJ, 432, L39. ATCA at 4.79 and 5.84 GHz (12'' × 22'') and Effelsberg 100-m at 10.6 GHz (1'), including polarization.  
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.  
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.  
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').  
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

**G357.7+0.3**

**RA:** 17<sup>h</sup>38<sup>m</sup>35<sup>s</sup>  
**Dec:** −30°44'

**1-GHz flux/Jy:** 10  
**Spectral index:** 0.4?

**Size/arcmin:** 24  
**Type:** S

**Radio:** Non-thermal shell in complex region.

**References:**

- Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz (4'.3 :  $S = 7 \pm 1.5$  Jy),  $S_{5\text{ GHz}} = 5.5 \pm 1.5$  Jy from surveys.  
 Gray 1994, MNRAS, 270, 836. MOST at 843 MHz (43'' × 84'').  
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.

**G358.0+3.8**

**RA:** 17<sup>h</sup>26<sup>m</sup>00<sup>s</sup>  
**Dec:** −28°36'

**1-GHz flux/Jy:** 1.5?  
**Spectral index:** ?

**Size/arcmin:** 38  
**Type:** S

**Radio:** Faint shell.

**References:**

- Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (2'.2 × 1'.3 :  $S = 2.5 \pm 1.3$  Jy), and NVSS at 1.4 GHz.

**G359.0–0.9**

**RA:** 17<sup>h</sup>46<sup>m</sup>50<sup>s</sup>  
**Dec:** −30°16'

**1-GHz flux/Jy:** 23  
**Spectral index:** 0.5

**Size/arcmin:** 23  
**Type:** S

**Radio:** Incomplete shell.

**X-ray:** Partial shell.

**References:**

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.  
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).  
 Gray 1994, MNRAS, 270, 836. MOST at 843 MHz (43'' × 86'').  
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').  
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.

**G359.1–0.5****RA:** 17<sup>h</sup>45<sup>m</sup>30<sup>s</sup>  
**Dec:** −29°57'**1-GHz flux/Jy:** 14  
**Spectral index:** 0.4?**Size/arcmin:** 24  
**Type:** S**Radio:** Non-thermal shell in complex region, crossed by the ‘snake’.**X-ray:** Centrally brightened.**Point sources:** Several compact radio sources near centre, OH masers around edge.**References:**

Downes *et al.* 1979, A&AS, 35, 1. From observations by Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6:S=13 Jy). Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz (4'.3:S=10±1.5 Jy) and 4.8 GHz (2'.4:S=8.1±0.5 Jy).

Uchida *et al.* 1992, ApJ, 398, 128. VLA at 1.5 GHz (11''×10''), and observations of nearby molecular material.

Uchida *et al.* 1992, AJ, 104, 1533. VLA at 1.4 GHz.

Gray 1994, MNRAS, 270, 836. MOST at 843 MHz (43''×85'').

Yusef-Zadeh *et al.* 1995, Sci, 270, 1801. VLA at 1.4 GHz (33''×31''), and 1.7 GHz for OH survey.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43''×24'').

Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.

**G359.1+0.9****RA:** 17<sup>h</sup>39<sup>m</sup>36<sup>s</sup>  
**Dec:** −29°11'**1-GHz flux/Jy:** 5?  
**Spectral index:** ?**Size/arcmin:** 12×11  
**Type:** S**Radio:** Shell, brightest in E.**References:**

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×88'':S=4.3 Jy).