

**A Catalogue of
Galactic Supernova Remnants
(2001 December version)**

D.A. Green

*Mullard Radio Astronomy Observatory
Cavendish Laboratory
Madingley Road
Cambridge CB3 0HE
UNITED KINGDOM*

E-mail: D.A.Green@mrao.cam.ac.uk

Please reference this catalogue as follows.

- Green D.A., 2001, 'A Catalogue of Galactic Supernova Remnants (2001 December version)', Mullard Radio Astronomy Observatory, Cavendish Laboratory, Cambridge, UK (available on the World-Wide-Web at "<http://www.mrao.cam.ac.uk/surveys/snrs/>").

1. The Catalogue Format

This catalogue of Galactic supernova remnants (SNRs) is an updated version of those presented in detail in Green (1984, 1988), in summary form in Green (1991, 1996) — hereafter Versions I, II, III and IV respectively — and on the World-Wide-Web (versions of 1995 July, 1996 August, 1998 September and 2000 August). (Note that Version IV, although published in 1996, was produced in 1993.)

This, the 2001 December version of the catalogue, contains 231 SNRs (which is six more than in the previous, 2000 August, version: ten new remnants have been added, and four entries removed), with about thousand references in the detailed listings, plus notes on several dozen possible or probable remnants.

For each remnant in the catalogue the following parameters are given.

- **Galactic Coordinates** of the source centroid, quoted to the nearest tenth of a degree as is conventional. (Note: in this catalogue additional leading zeros are not used.)
- **Other Names** that are commonly used for the remnant. These are given in parentheses if the remnant is only a part of the source. For some remnants, notably the Crab Nebula, not all common names are given.
- **Right Ascension** and **Declination** of the source centroid. The accuracy of the quoted values depends on the size of the remnant; for small remnants they are to the nearest few seconds of time and the nearest minute of arc respectively, whereas for larger remnants they are rounded to coarser values, but are in every case sufficient to specify a point within the boundary of the remnant. These coordinates are almost always deduced from radio maps rather than from X-ray or optical observations, and are for J2000.0.
- **Angular Size** of the remnant, in arcminutes, usually taken from the highest resolution radio map available. The boundary of most remnants approximates reasonably well to a circle or an ellipse; a single value is quoted for the angular size of the more nearly circular remnants, which is the diameter of a circle with an area equal to that of the remnant, but for elongated remnants the product of two values is quoted, and these are the major and minor axes of the remnant boundary modelled as an ellipse. In a few cases an ellipse is not a satisfactory description of the boundary of the object (refer to the description of the individual object given in its catalogue entry), although an angular size is still quoted for information. For 'filled-centre' remnants the size quoted is for the largest extent of the observed radio emission, not, as at times has been used by others, the half-width of the centrally brightened peak.

- **Flux Density** of the remnant at 1 GHz in jansky. This is *not* a measured value, but is deduced from the observed radio frequency spectrum of the source. The frequency of 1 GHz is chosen because flux density measurements at frequencies both above and below this value are usually available.
- **Spectral Index** of the integrated radio emission from the remnant, α (here defined in the sense, $S \propto \nu^{-\alpha}$, where S is the flux density at a frequency ν), either a value that is quoted in the literature, or one deduced from the available integrated flux densities of the remnant. For several SNRs a simple power law is not adequate to describe their radio spectra, either because there is evidence that the integrated spectrum is curved or the spectral index varies across the face of the remnant. In these cases the spectral index is given as ‘varies’ (refer to the description of the remnant and recent references in the catalogue entry for more information). In some cases, for example where the remnant is highly confused with thermal emission, the spectral index is given as ‘?’ since no value can be deduced with any confidence.
- **Type** of the SNR, either ‘S’, ‘F’ or ‘C’ if the remnant shows a ‘shell’, ‘filled-centre’ or ‘composite’ (or ‘combination’) radio structure (or ‘S?’, ‘F?’ or ‘C?’, respectively, if there is some uncertainty), or ‘?’ in several cases where an object is conventionally regarded as an SNR even though its nature is poorly known or not well understood. (Note: the term ‘composite’ has been used in a different sense by some authors, to describe SNRs with shell radio and centrally-brightened X-ray morphologies. An alternative term used to describe such remnants is ‘mixed morphology’, see Rho & Petre 1998.)

In the detailed listings, for each remnant, notes on a variety of topics are given. First, it is noted if other Galactic coordinates have at times been used to label it (usually before good observations have revealed the full extent of the object), if the SNR is thought to be the remnant of a historical SN, or if the nature of the source as an SNR has been questioned (in which case an appropriate reference is usually given later in the entry). Brief descriptions of the remnant from the available radio, optical and X-ray observations as applicable are then given, together with notes on available distance determinations, and any point sources or pulsars in the field of the object (although they may not necessarily be related to the remnant). Finally, appropriate references to observations are given for each remnant, complete with journal, volume, page, and a short description of what information each paper contains (for radio observations these include the telescopes used, the observing frequencies and resolutions, together with any flux density determinations). These references are *not* complete, but cover representative and recent observations of the remnant, and they should themselves include references to earlier work. The references do not generally include large observational surveys — of particular interest in this respect are: the Effelsberg 100-m survey at 2.7 GHz of the Galactic plane $358^\circ < l < 240^\circ$, $|b| < 5^\circ$ by Reich *et al.* (1990) and Fürst *et al.* (1990); reviews of the radio spectra of some SNRs by Kassim (1989), Kovalenko, Pynzar’ & Udal’tsov (1994) and Trushkin (1998); the Parkes 64-m survey at 2.4 GHz of the Galactic plane $238^\circ < l < 365^\circ$, $|b| < 5^\circ$ by Duncan *et al.* (1995) and Duncan *et al.* (1997); the Molonglo Galactic plane survey at 843 MHz of $245^\circ < l < 355^\circ$, $|b| < 1.5^\circ$ by Green *et al.* (1999); the survey of $345^\circ < l < 255^\circ$, $|b| < 5^\circ$ at 8.35 and 14.35 GHz by Langston *et al.* (2000); reviews of *Einstein* X-ray imaging and spectroscopic observations of Galactic SNRs by Seward (1990) and Lum *et al.* (1992) respectively; surveys of *IRAS* observations of SNRs and their immediate surroundings by Arendt (1989) and by Saken, Fesen & Shull (1992); the survey of HI emission towards SNRs by Koo & Heiles (1991); and the catalogue by Fesen & Hurford (1996) of UV/optical/infra-red lines identified in SNRs.

A summary of the data available for all 231 remnants in the catalogue is given in Table I. The other names for SNRs are listed in Table II, and the abbreviations for journals, proceedings and telescopes are listed in Table III. The detailed listings for each SNR are given in Table IV.

2. Revisions and Notes

2.1 Objects no longer thought to be SNRs

The following objects, which were listed in Version I of the catalogue were removed because they were no longer thought to be remnants, or were poorly observed (see Version II for references and further details): G2.4+1.4 (see also Gray 1994a; Goss & Lozinskaya 1995; Polcaro *et al.* 1995), G41.9–4.1 (=CTB 73, PKS 1920+06), G47.6+6.1 (=CTB 63), G53.9+0.3 (part of HC40), G93.4+1.8 (=NRAO 655), G123.2+2.9, G194.7+0.4 (the Origem Loop), G287.8–0.5 (see below), G322.3–1.2 (=Kes 24) and G343.0–6.0 (but see below). G350.1–0.3, which was listed in Version II of the catalogue, was removed as it is no longer thought to be a SNR (see Version III for details). G358.4–1.9, which was listed in Version IV of the catalogue, was removed, as following the discussion of Gray (1994a), as it is not clear that this is a SNR. G240.9–0.9, G299.0+0.2 and G328.0+0.3, which were listed in 1995 July version of the catalogue, were removed from the 1996 August version, following the improved observations of Duncan *et al.* (1996) and Whiteoak & Green (1996). For the 1998 September revision of the catalogue G350.0–1.8 was incorporated into G350.0–2.0, and G337.0–0.1 refers to a smaller remnant than that previously catalogued with the same name.

G112.0+1.2, G117.4+5.0, G152.2–1.2 and G211.7–1.1 – which were reported as SNRs by Bonsignori-Facondi & Tomasi (1979) – have been removed from this version of the catalogue, as the first three of these are not confirmed as SNRs from the ongoing *Canadian Galactic Plane Survey* (Roland Kothes, private communication).

The following objects, which have been reported as SNRs, but have not been included in any of the versions of the SNR catalogue, have subsequently been shown not to be SNRs.

- G70.7+1.2, which was reported as a SNR by Reich *et al.* (1985), but this has not been confirmed by later observations (see Green 1986; de Muizon *et al.* 1988; Becker & Fesen 1988; Caswell 1988; Bally *et al.* 1989; Phillips, Onello & Kulkarni 1993; Onello *et al.* 1995).
- G81.6+1.0 a possible SNR in W75 reported by Ward-Thompson & Robson (1991). From the published data (see the observations in Wendker, Higgs & Landecker 1991) it was noted in Version IV of the catalogue that this is thermal source not a SNR, because of its thermal radio spectrum, and high infrared-to-radio emission (see also the subsequent discussion by Wendker *et al.* 1993).
- Green & Gull (1984) suggested G227.1+1.0 as a very young SNR, but subsequent observations (Channan *et al.* 1986; Green & Gull 1986) have shown that this is most likely an extragalactic source, not an SNR.
- A candidate SNR, G274.7–2.8, identified by Helfand & Channan (1989), has been shown not to be a SNR by Caswell & Stewart (1991).
- G159.6–18.5, was suggested as a SN by Pauls & Schwartz (1989), from IRAS and other observations, but is probably an HII region (see Andersson *et al.* 2000).
- G25.5+0.2, which was reported as a very young SNR by Cowan *et al.* (1989), although this identification was not certain (see White & Becker 1990; Green 1990; Zijlstra 1991). Sramek *et al.* (1992) report the detection of recombination lines from this source (also see Subrahmanyan *et al.* 1993). Becklin *et al.* (1994) identify G25.5+0.2 as a ring nebula around a luminous blue star. See also Clark, Steele & Lanfer (2000), who identified a bipolar shell around G25.5+0.2 with similarities to η Carinae.
- Most of the possible SNRs listed by Gorham (1990) — following up SNR candidates suggested by Kassim (1988) — have been shown not to be SNRs by Gorham, Kulkarni & Prince (1993).
- G203.2–12.3, a optical ring about 3 arcmin in diameter, was reported as a possible SNR by Winkler & Reipurth (1992), but was shown to be a Herbig–Haro object (HH 311) by Reipurth, Bally & Devine (1997).
- G359.87+0.18 was reported as a possible young SNR near the Galactic Centre by Yusef-Zadeh, Cotton & Reynolds (1998), but was shown to be a radio galaxy by Lazio *et al.* (1999).
- G106.6+2.9, a small remnant proposed by Halpern *et al.* (2001a), is incorporated into the larger catalogued remnant G106.3+2.7.

Some entries in the catalogue have been renamed, due to improved observations revealing a larger true extent for the object (previously G5.3–1.0 is now G5.4–1.2; G193.3–1.5 is now G192.8–1.1; G308.7+0.0 is now incorporated into G308.8–0.1). G337.0–0.1 now refers to a small (1.5 arcmin) remnant, rather than larger supposed remnant at this position (see Sarma *et al.* 1997), and G350.0–2.0 now incorporates the previously catalogued G350.0–1.8, based on the improved observations of Gaensler (1998).

2.2 New SNRs

The following remnants were added to Version II of the catalogue: G0.9+0.1, G1.9+0.3, G5.9+3.1, G6.4+4.0, G8.7–0.1, G16.8–1.1, G18.9–1.1, G20.0–0.2, G27.8+0.6, G30.7+1.0, G31.5–0.6, G36.6–0.7, G42.8+0.6, G45.7–0.4, G54.1+0.3, G73.9+0.9, G179.0+2.6, G312.4–0.4, G357.7+0.3 and G359.1–0.5.

The following remnants were added to Version III of the catalogue: G4.2–3.5, G5.2–2.6, G6.1+1.2, G8.7–5.0, G13.5+0.2, G15.1–1.6, G16.7+0.1, G17.4–2.3, G17.8–2.6, G30.7–2.0, G36.6+2.6, G43.9+1.6, G59.8+1.2, G65.1+0.6, G68.6–1.2, G69.7+1.0, G279.0+1.1, G284.3–1.8 (=MSH 10–53), G358.4–1.9 and G359.0–0.9.

The following remnants were added to Version IV of the catalogue: G59.5+0.1, G67.7+1.8, G84.9+0.5, G156.2+5.7, G318.9+0.4, G322.5–0.1, G343.1–2.3 and G348.5–0.0.

The following remnants were added to 1995 July version of the catalogue: G1.0–0.1, G1.4–0.1, G3.7–0.2, G3.8+0.3, G28.8+1.5, G76.9+1.0, G272.2–3.2, G341.2+0.9, G354.1+0.1, G355.6–0.0, G356.3–0.3, G356.3–1.5 and G359.1+0.9.

The following remnants were added to the 1996 August version of the catalogue: G13.3–1.3, G286.5–1.2, G289.7–0.3, G294.1–0.0, G299.2–2.9, G299.6–0.5, G301.4–1.0, G308.1–0.7, G310.6–0.3, G310.8–0.4, G315.9–0.0, G317.3–0.2, G318.2+0.1, G320.6–1.6, G321.9–1.1, G327.4+1.0, G329.7+0.4, G342.1+0.9, G343.1–0.7, G345.7–0.2, G349.2–0.1, G351.7+0.8, G351.9–0.9 and G354.8–0.8.

The following remnants were added to the 1998 September version of the catalogue: G0.3+0.0, G32.1–0.9, G55.0+0.3, G63.7+1.1 and G182.4+4.3.

The following remnants were added to the 2000 August version of the catalogue: G7.0–0.1, G16.2–2.7, G29.6+0.1, G266.2–1.2 and G347.3–0.5.

The following remnants have been added to this version of the catalogue.

- Three high latitude SNRs (G4.8+6.2, G356.2+4.5 and G358.0+3.8) – which were proposed as possible remnants by Duncan *et al.* (1997) – which were confirmed as remnants by Bhatnagar (2000), from radio observations made with the GMRT. (Note that G4.8+6.2 was referred to as G4.5+6.2 by Duncan *et al.* 1997)
- G28.6–0.1 – which had previously been listed as a possible SNR, based on the radio observations of Helfand *et al.* (1989) – has now been included in the catalogue, as X-ray observations by Bamba *et al.* (2001) have confirmed its nature as a SNR.
- Two remnants, G85.4+0.7 and G85.9–0.6, identified from radio observations from the ongoing *Canadian Galactic Plane Survey* by Kothes *et al.* (2001).
- G106.3+2.7, a remnant identified from radio observations by Pineault & Joncas (2000) – this incorporates a smaller proposed remnant (G106.6+2.9) that has been discussed by Halpern *et al.* (2001a,b) – see also Kothes, Uyaniker & Pineault (2001).
- G292.2–0.5, around the young pulsar J1119–6127, which was detected at radio wavelength by Crawford *et al.* (2001), and in X-rays by Pivovarov *et al.* (2001).
- G343.0–0.6 (=RCW 114) – which had been listed in Version I of the catalogue, but was removed as it was poorly defined by the available observations – has now been included in the catalogue, because of the improved observations of Walker & Zealey (2001).
- A remnant, G353.9–2.0, identified in the *NRAO VLA Sky Survey* radio data (see Condon *et al.* 1998) by Green (2001), who presented improved radio observations of this nearly circular shell SNR.

2.3 Possible and probable SNRs not listed in the catalogue

The following are possible or probable SNRs for which further observations are required to confirm their nature or parameters, or for which observations are not yet in the published literature.

2.3.1 Radio

- A possible SNR near the Galactic centre reported by Ho *et al.* (1985) from radio observations.
- Gosachinskiĭ (1985) reported evidence for non-thermal radio emission, presumably from SNRs, associated with several bright, thermal Galactic sources (also see Odegard 1986, who questions the reliability of some of Gosachinskiĭ's results).
- G300.1+9.4, a possible SNR nearly 2° in diameter reported by Dubner, Colomb & Giacani (1986) from radio observations.
- Routledge & Vaneldik (1988) report a possible faint shell SNR nearly 2° in diameter at radio wavelengths, near the young pulsar PSR 1930+22 – see also Gómez-González & del Romero (1983), who report a smaller (about 40 arcmin) possible SNR (G57.1+1.7) associated with this pulsar, and see Caswell, Landecker & Feldman (1985) and Kovalenko (1989).
- Five possible remnants (G45.9–0.1, G71.6–0.5, G72.2–0.3, G83.0–0.2 and G85.2–1.2) of the eleven reported by Taylor, Wallace & Goss (1992) from a radio survey of part of the Galactic plane. (Three of the other possible SNRs reported by Taylor *et al.*, are included in the catalogue as G55.0+0.3, G63.7+1.1 and G76.9+1.0.)
- A faint, poorly defined possible remnant G41.1+1.2 reported by Gorham, Kulkarni & Prince (1993) from radio observations.
- G9.7–0.1, a possible SNR report by Frail, Kassim & Weiler (1994) from radio observations.
- G355.4+0.7, G356.6+0.1, G357.1–0.2, G358.1+1.0, G358.5–0.9, G358.7+0.7, G359.2–1.1, G3.1–0.6 and G4.2+0.0, which are among the possible SNRs listed by Gray (1994b) from radio observations near the Galactic centre.
- G104.7+2.8, a possible SNR reported by Green & Joncas (1994) from radio observations. However, recent observations at 10.7 GHz (W. Reich, private communication) cast doubt on this identification, as they do not support a non-thermal radio spectrum for the source.
- G11.2–1.1, a possible SNR listed by Kovalenko, Pynzar' & Udal'tsov (1994), based on unpublished radio studies (Trushkin 1988, preprint).
- Duncan *et al.* (1995) and Duncan *et al.* (1997) list several large-scale (1.5 to 10 degree), and smaller, low radio surface-brightness candidate SNRs from the Parkes 2.4-GHz survey of $270^\circ < l < 360^\circ$. Several of these candidates have been confirmed as SNRs by subsequent, improved observations, and are included in the catalogue.
- Whiteoak & Green (1996), from their radio survey of much of the southern Galactic plane, list 16 possible SNRs (G308.4–1.4, G317.5+0.9, G319.9–0.7, G320.6–0.9, G322.7+0.1, G322.9–0.0, G323.2–1.0, G324.1+0.1, G325.0–0.3, G331.8–0.0, G337.2+0.1, G339.6–0.6, G345.1+0.2, G345.1–0.2, G348.8+1.1 and G350.1–0.3).
- Several candidate SNRs reported by Combi & Romero (1998), Combi, Romero & Arnal (1998), Combi, Romero & Benaglia (1998), Punsly *et al.* (2000) and Combi *et al.* (2001).
- G359.09–0.02, a possible SNR noted by LaRosa *et al.* (2000).
- A possible SNR, G313.3+0.1, near an unidentified Galactic plane γ -ray source (see Roberts *et al.* 1999).
- A likely SNR, called G359.92–0.09, adjacent to G0.0+0.0 (=Sgr A East) at the Galactic centre, see Coil & Ho (2000), and references therein.
- A possible SNR, G328.6–0.0, noted by McClure-Griffiths *et al.* (2001) in the test region of the *Southern Galactic Plane Survey*.
- G346.5–0.1, an arc of radio emission observed by Gaensler *et al.* (2001), which is potentially part of a SNR, but requires further observations to confirm its nature.
- Giacani *et al.* (2001) presented observations of a pulsar wind nebula around PSR J1709–4428, which may be part of the catalogued remnant G343.1–2.3, or may represent another object.

2.3.2 UV/Optical/Infra-red

- A possible SNR overlapping G296.1–0.5, identified from optical (and X-ray) observations by Hutchings, Crampton & Cowley (1981).
- A SNR (G260.4–3.3) about 4 arcmin in diameter within the Puppis A remnant identified optically by Winkler *et al.* (1989). This has not been detected at radio wavelengths (see Dubner *et al.* 1991).
- A possible SNR (G32.1+0.1) reported from optical spectroscopy by Thompson, Djorgovski & de Carvalho (1991), following up radio and infrared observations of Jones, Garwood & Dickey (1988).
- G75.5+2.4, a possible large (about 2°) old SNR in Cygnus suggested by Nichols-Bohlin & Fesen (1993) from infra-red and optical observations (see also Dewdney & Lozinskaya 1994; Marston 1996; Esipov *et al.* 1996).
- A possible optical SNR (G247.8+4.9) noted by Weinberger (1995), which may be Balmer dominated (see also Weinberger *et al.* 1998 and Zanin & Kerber 2000).
- An optical shell around the Coalsack Nebula (near $l = 300^\circ$, $b = 0^\circ$) identified by Walker & Zealey (1998). This coincides with one of the large possible SNRs suggested by Duncan *et al.* (1995), from radio observations.

2.3.3 X-ray

- H1538–32 a large X-ray source in Lupus, near $l = 307^\circ$, $b = +20^\circ$ (Riegler, Agrawal & Gull 1980; see also Colomb, Dubner & Giacani 1984; Gahm *et al.* 1990) which is a possible old SNR;
- The Monogem ring, near $l = 203^\circ$, $b = +12^\circ$, is a possible old SNR (see Nousek *et al.* 1981; Plucinsky *et al.* 1996; and references therein).
- X-ray emission in the Gum Nebula near $l = 250^\circ$, $b = 0^\circ$ (Leahy, Nousek & Garmire 1992, see also Reynolds 1976, Dubner *et al.* 1992, Duncan *et al.* 1996, Reynoso & Dubner 1997, Heiles 1998) which, together with optical spectroscopy indicate the existence of a possible old remnant in this region.
- An X-ray enhancement near $l = 200^\circ$, $b = -40^\circ$, which is possibly due to an old SNR in Eridanus (Naranan *et al.* 1976; see also Burrows *et al.* 1993; Snowden *et al.* 1995; Heiles 1998, Boumis *et al.* 2001).
- G189.6+3.3, a faint, possible SNR overlapping G189.1+3.0 (=IC443) identified by Asaoka & Aschenbach (1994) from ROSAT X-ray observations.
- G117.7+0.6, a faint shell of soft X-ray emission near CTB1 (=G116.9+0.2), which contains a pulsar (Hailey & Craig 1995; see also Craig, Hailey & Pisarski 1997).
- A possible SNR identified in X-rays around the pulsar B1828–13 (see Finley, Srinivasan & Park 1996).
- A possible, large SNR, G69.4+1.2, identified as an X-ray shell by Yoshita, Miyata & Tsunemi (1999, 2000).

2.3.4 Other

- G287.8–0.5, which is associated with η Carinae, was listed in Version I as a SNR, but was removed from the catalogue in Version II as its parameters are uncertain (see Jones 1973; Retallack 1984; Tateyama, Strauss & Kaufmann 1991; and the discussion in Version II).
- G359.2–0.8 (the ‘mouse’), near the Galactic centre, which has been suggested as being analogous to the central region of CTB 80 (=G69.0+2.7) by Predehl & Kulkarni (1995).

It should also be noted that some radio continuum and HI loops in the Galactic plane (see, for example, Berkhuijsen 1973) may be parts of very large, old SNRs, but they have not been included in the catalogue (see also Combi *et al.* 1995; Maciejewski *et al.* 1996; Kim & Koo 2000; Normandeau *et al.* 2000; Woermann, Gaylard & Otrupcek 2001), nor have pulsar wind nebulae (see, for example, Gaensler *et al.* 1998).

2.4 Questionable SNRs listed in the catalogue

As noted in Versions II and IV of the catalogue, the following sources are listed as SNRs, although, as discussed in each case, the identifications are not certain: G5.4–1.2, G39.7–2.0 (=W50), G65.7+1.2 (=DA 495), G69.0+2.7 (=CTB 80), G318.9+0.4 and G357.7–0.1. The nature of G76.9+1.0 (an unusual radio source similar to G65.7+1.2), of G354.1+0.1 (which appears may be similar to G357.7–0.1 (=MHS 17–39)) and G10.0–0.3 (which has been associated with a soft gamma repeater) are also uncertain (see Landecker, Higgs & Wendker 1993, Frail, Goss & Whiteoak 1994 and Gaensler *et al.* 2001 respectively).

There are also some objects that have been identified as SNRs and are listed in the catalogue, although they have been barely resolved in the available observations, or are faint, and have not been well separated from confusing background or nearby thermal emission, and their identification as SNRs, or at least their parameters remain uncertain.

Acknowledgements

I am grateful to the many colleagues who have commented on previous version of the catalogue, and have brought errors and omissions to my attention. No doubt errors remain in this version, and I am always happy to receive feedback from users of the catalogue. This research has made use of NASA's Astrophysics Data System Bibliographic Services. I am also grateful for the hospitality of IUCAA in Pune, India, during November and December 2001, when much of the revision of the catalogue was done, and to the support provided by the *The Cambridge Society, Bombay Scholarship Fund* for the visit.

References

- Arendt R.G., 1989, *ApJS*, 70, 181.
- Andersson B.-G., Wannier, P.G., Moriarty-Schieven G.H. & Bakker, E.J., 2000, *AJ*, 119, 1325.
- Asaoka I. & Aschenbach B., 1994, *A&A*, 284, 573.
- Bally J., Pound M.W., Stark A.A., Israel F., Hirano N., Kameya O., Sunada K., Hayashi M., Thronson H. & Hereld, M., 1989, *ApJ*, 338, L65.
- Bamba A., Ueno M., Koyama K. & Yamauchi S., 2001, *PASJ*, 53, L21.
- Becker R.H. & Fesen R.A., 1988, *ApJ*, 334, L35.
- Becklin E.E., Zuckerman B., McLean I.S. & Geballe T., 1994, *ApJ*, 430, 774.
- Berkhuijsen E.M., 1973, *A&A*, 24, 143.
- Bhatnagar S., 2000, *MNRAS*, 317, 453.
- Bonsignori-Facondi S.R. & Tomasi P., 1979, *A&A*, 77, 93.
- Boumis P., Dickinson C., Meaburn J., Goudis C.D., Christopoulou P.E., López J.A., Bryce M. & Redman M.P., 2001, *MNRAS*, 320, 61.
- Burrows D.N., Singh K.P., Nousek J.A., Garmire G.P. & Good, J., 1993, *ApJ*, 406, 97.
- Caswell J.L., 1988, in *SNRISM*, p269.
- Caswell J.L. & Stewart R.T., 1991, *PASAu*, 9, 103.
- Caswell J.L., Landecker T.L. & Feldman P.A., 1985, *AJ*, 90, 488.
- Channan G.A., Helfand D.J., Spinrad H. & Ebnetter K., 1986, *Natur*, 320, 41.
- Clark S.J., Steele I.A. & Langer N., 2000, *ApJ*, 541, 67.
- Coil A.L. & Ho P.T.P., 2000, *ApJ*, 533, 245.
- Colomb F.R., Dubner G.M. & Giacani E.B., 1984, *A&A*, 130, 294.
- Combi J.A. & Romero G.E., 1998, *A&AS*, 128, 423.
- Combi J.A., Testari J.C., Romero G.E. & Colomb F.R., 1995, *A&A*, 296, 514.
- Combi J.A., Romero G.E. & Arnal E.M., 1998, *A&A*, 333, 298.
- Combi J.A., Romero G.E. & Benaglia P., 1998, *A&A*, 333, L91.
- Combi J.A., Romero G.E., Benaglia P. & Jonas J.L., 2001, *A&A*, 366, 1047.
- Condon J.J., Cotton W.D., Greisen E.W., Yin Q.F., Perley R.A., Taylor G.B. & Broderick J.J., 1998, *AJ*, 115, 1693.
- Cowan J.J., Ekers R.D., Goss W.M., Sramek R.A., Roberts, D.A. & Branch D., 1989, *MNRAS*, 241, 613.
- Craig W.W., Hailey C.J. & Pisarski R.L., 1997, *ApJ*, 488, 307.
- Crawford F., Gaensler B.M., Kaspi V.M., Manchester R.N., Camilo F., Lyne A.G. & Pivovarov M.J., 2001, *ApJ*, 554, 152.
- de Muizon M., Strom R.G., Oort M.J.A., Claas J.J. & Braun R., 1988, *A&A*, 193, 248.
- Dewdney P.E. & Lozinskaya T.A., 1994, *AJ*, 108, 2212.
- Dubner G.M., Colomb F.R. & Giacani E.B., 1986, *AJ*, 91, 343.
- Dubner G.M., Braun R., Winkler P.F. & Goss 1991, *AJ*, 101, 1466.
- Dubner G., Giacani E., Cappa de Nicolau C. & Reynoso E., 1992, *A&AS*, 96, 505.
- Duncan A.R., Stewart R.T., Haynes R.F. & Jones K.L., 1995, *MNRAS*, 277, 36.
- Duncan A.R., Stewart R.T., Haynes R.F. & Jones K.L., 1996, *MNRAS*, 280, 252.
- Duncan A.R., Stewart R.T., Haynes R.F. & Jones K.L., 1997, *MNRAS*, 287, 722.
- Esipov V.F., Lozinskaya T.A., Mel'nikov V.V., Pravdikova V.V., Sitnik T.G. & Nichol-Bohlin J., 1996, *ALet*, 22, 509.
- Fesen R.A. & Hurford A.P., 1996, *ApJS*, 106, 563.
- Finley J.P., Srinivasan R. & Park S., 1996, *ApJ*, 466, 938.
- Frail D.A., Goss W.M. & Whiteoak J.B.Z., 1994, *ApJ*, 437, 781.
- Frail D.A., Kassim N.E. & Weiler K.W., 1994, *AJ*, 107, 1120.
- Fürst E., Reich W., Reich P. & Reif K., 1990, *A&AS*, 85, 691.
- Gaensler B.M., 1998, *ApJ*, 493, 781.
- Gaensler B.M., Stappers B.W., Frail D.A. & Johnston S., 1998, *ApJ*, 499, L69.
- Gaensler B.M., Slane P.O., Gotthelf E.V. & Vasisht G., 2001, *ApJ*, 559, 963.

-
- Gahm G.F., Gebeyehu M., Lindgren M., Magnusson P., Modigh P. & Nordh H.L., 1990, A&A, 228, 477.
Giacani E.B., Frail D.A., Goss W.M. & Vieytes M., 2001, AJ, 121, 3133.
Gómez-González J. & del Romero A., 1983, A&A, 123, L5.
Gorham P.W., 1990, ApJ, 364, 187.
Gorham P.W., Kulkarni S.K. & Prince T.A., 1993, AJ, 105, 314.
Gosachinskiĭ I.V., 1985, SvA, 29, 128.
Goss W.M. & Lozinskaya T.A., 1995, ApJ, 439, 637.
Gray A.D., 1994a, MNRAS, 270, 835.
Gray A.D., 1994b, MNRAS, 270, 847.
Green A.J., Cram L.E., Large M.I. & Ye T.S., 1999, ApJS, 122, 207.
Green D.A., 1984, MNRAS, 209, 449 (Version I).
Green D.A., 1986, MNRAS, 219, 39P.
Green D.A., 1988, Ap&SS, 148, 3 (Version II).
Green D.A., 1990, AJ, 100, 1241.
Green D.A., 1991, PASP, 103, 209 (Version III).
Green D.A., 1996, in *Supernovae and Supernova Remnants*, (proceedings of IAU Colloquium 145, Xi'an China, 1993 May 24–29), eds McCray R. & Wang Z., (Cambridge University Press), p.419 (Version IV).
Green D.A., 2001, MNRAS, 326, 283.
Green D.A. & Gull S.F., 1984, Natur, 312, 527.
Green D.A. & Gull S.F., 1986, Natur, 320, 42.
Green D.A. & Joncas G., 1994, A&AS, 104, 481.
Hailey C.J. & Craig W.W., 1995, ApJ, 455, L151.
Halpern J.P., Camilo F., Gotthelf E.V., Helfand D.J., Kramer M., Lyne A.G., Leighly K.M. & Eracleous M., 2001a, ApJ, 552, L125.
Halpern J.P., Gotthelf E.V., Leighly K.M. & Helfand D.J., 2001b, ApJ, 547, 323.
Heiles C., 1998, ApJ, 498, 689.
Helfand D.J. & Channan G.A., 1989, AJ, 98, 1652.
Helfand D.J., Velusamy T., Becker R.H. & Lockman F.J., 1989, ApJ, 341, 151.
Hutchings J.B., Crampton D. & Cowley P.A., 1981, AJ, 86, 871.
Ho P.T., Jackson J.M., Barrett A.H. & Armstrong J.T., 1985, ApJ, 288, 575.
Jones B.B., 1973, AuJPh, 26, 545.
Jones T.J., Garwood R. & Dickey J.M. 1988, ApJ, 328, 559.
Kassim N.E., 1988, ApJ, 328, L55.
Kassim N.E., 1989, ApJS, 71, 799.
Kim K.-T. & Koo B.-C., 2000, ApJ, 529, 229.
Koo B.-C. & Heiles C., 1991, ApJ, 382, 204.
Kothes R., Landecker T.L., Foster T. & Leahy D.A., 2001, A&A, 376, 641.
Kothes R., Uyaniker B. & Pineault S., 2001, ApJ, 560, 236.
Kovalenko A.V., 1989, SvAL, 15, 144.
Kovalenko A.V., Pynzar' A.V. & Udal'tsov V.A., 1994, ARep, 38, 95.
Landecker T.L., Higgs L.A. & Wendker H.I., 1993, A&A, 276, 522.
Langston G., Minter A., D'Addario L., Eberhart K., Koski K. & Zuber J., 2000, AJ, 119, 2801.
LaRosa T.N., Kassim N.E., Lazio T.J.W. & Hyman S.D., 2000, AJ, 119, 207.
Lazio T.J.W., Anantharamaiah K.R., Goss W.M., Kassim N.E. & Cordes J.M., 1999, ApJ, 515, 196.
Leahy D.A., Nousek J. & Garmire G., 1992, ApJ, 385, 561.
Lum K.S.K., Canizares C.R., Clark S.W., Coyne J.M., Markert T.H., Saez P.J., Schattenburg M.L. & Winkler P.F., 1992, ApJS, 78, 423.
McClure-Griffiths N.M., Green A.J., Dickey J.M., Gaensler B.M., Haynes R.F. & Wieringa M.H., 2001, ApJ, 551, 394.
Maciejewski W., Murphy E.M., Lockman F.J. & Savage B.D., 1996, ApJ, 469, 238.
Marston A.P., 1996, AJ, 112, 2828.

-
- Naranan S., Shulman S., Friedman H. & Fritz G., 1976, ApJ, 208, 718.
Nichols-Bohlin J. & Fesen R.A., 1993, AJ, 105, 672.
Normandeau M., Taylor A.R., Dewdney P.E. & Basu S., 2000, AJ, 119, 2982.
Nousek J.A., Cowie L.L., Hu E., Lindblad C.J. & Garmire G.P., 1981, ApJ, 248, 152.
Odegard N., 1986, AJ, 92, 1372.
Onello J.S., DePree C.G., Phillips J.A. & Goss W.M., 1995, ApJ, 449, L127.
Pauls T. & Schwarz P.R., 1989, in *The Physics and Chemistry of Interstellar Molecular Clouds*, eds Winnewisser G. & Armstrong T.J. (Springer), p.225.
Phillips J.A., Onello J.S. & Kulkarni S.R., 1993, ApJ, 415, 143.
Pineault S. & Joncas G., 2000, AJ, 120, 3218.
Pivovarov M.J., Kaspi V.M., Camilo F., Gaensler B.M. & Crawford F., 2001, ApJ, 554, 161.
Plucinsky P.P., Snowden S.L., Aschenbach B., Eggar R., Edgar R.J. & McCammon D., 1996, ApJ, 463, 224.
Polcaro V.F., Rossi C., Norci L. & Viotti R., 1995, A&A, 303, 211.
Predehl P. & Kulkarni S.R., 1995, A&A, 294, L29.
Punsly B., Romero G.E., Torres D.F. & Combi J.A., 2000, A&A, 364, 552.
Reich W., Fürst E., Altenhoff W.J., Reich P. & Junkes N., 1985, A&A, 151, L10.
Reich W., Fürst E., Reich P. & Reif K., 1990, A&AS, 85, 633.
Reipurth B., Bally J. & Divine D., 1997, AJ, 114, 2708.
Retallack D.S., 1983, MNRAS, 204, 669.
Reynolds R., 1976, ApJ, 206, 679.
Reynoso E.M. & Dubner G.M. 1997, A&AS, 123, 31.
Rho J. & Petre R., 1998, ApJ, 503, L167.
Riegler G.R., Agrawal P.C. & Gull S.F., 1980, ApJ, 235, L71.
Roberts M.S.E., Romani R.W., Johnston S. & Green A.J., 1999, ApJ, 515, 712.
Routledge D. & Vaneldik J.F., 1988, ApJ, 326, 751.
Saken J.M., Fesen R.A. & Shull J.M., 1992, ApJS, 81, 715.
Sarma A.P., Goss W.M., Green A.J. & Frail D.A., 1997, ApJ, 483, 335.
Seward F.D., 1990, ApJS, 73, 781.
Snowden S.L., Burrows D.N., Sanders W.T., Aschenbach B. & Pfeffermann E., 1995, ApJ, 439, 399.
Sramek R.A., Cowan J.J., Roberts D.A., Goss W.M. & Ekers R.D., 1992, AJ, 104, 704.
Subrahmanyam R., Ekers R.D., Wilson W.E., Goss W.M. & Allen, D.A., 1993, MNRAS, 263, 868.
Tateyama C.E., Strauss F.M. & Kaufmann P., 1991, MNRAS, 249, 716.
Taylor A.R., Wallace B.J. & Goss W.M., 1992, AJ, 103, 931.
Thompson D.J., Djorgovski S. & de Carvalho R.R., 1991, PASP, 103, 487.
Trushkin S.A., 1998, BSAO, 46, 62.
Walker A. & Zealey W.J., 1998, PASA, 15, 79.
Walker A. & Zealey W.J., 2001, MNRAS, 325, 287.
Ward-Thompson D. & Robson E.I., 1991, MNRAS, 248, 670.
Weinberger R., 1995, PASP, 107, 58.
Weinberger R., Tajitsu A., Tamura S. & Yadoumaru Y., 1998, PASP, 110, 722.
Wendker H.I., Higgs L.A. & Landecker T.L., 1991, A&A, 241, 551.
Wendker H.I., Higgs L.A., Landecker T.L. & Ward-Thompson D., 1993, MNRAS, 263, 543.
White R.L. & Becker R.H., 1990, MNRAS, 244, 12P.
Whiteoak J.B.Z. & Green A.J., 1996, A&AS, 118, 329.
Winkler P.F. & Reipurth B., 1992, ApJ, 389, L25.
Winkler P.F., Kirshner R.P., Hughes J.P. & Heathcote S.R. 1989, Natur, 337, 48.
Woermann B., Gaylard M.J. & Otrupcek R., 2001, MNRAS, 325, 1213.
Yoshita K., Niyata E. & Tsunemi H., 1999, AN, 320, 344.
Yoshita K., Niyata E. & Tsunemi H., 2000, PASJ, 52, 867.
Yusef-Zadeh F., Cotton W.D. & Reynolds S.P., 1998, ApJ, 498, L55.
Zanin C. & Kerber F., 2000, A&A, 356, 274.
Zijlstra A.A., 1991, MNRAS, 248, 11P.

| l | b | RA (J2000.0) (h m s) | Dec ($^{\circ}$ $'$) | size /arcmin | type | Flux at 1 GHz/Jy | spectral index | other name(s) |
|------|------|-------------------------|---------------------------|------------------|------|---------------------|-------------------|-----------------------|
| 0.0 | +0.0 | 17 45 44 | -29 00 | 3.5×2.5 | S | 100? | 0.8? | Sgr A East |
| 0.3 | +0.0 | 17 46 15 | -28 38 | 15×8 | S | 22 | 0.6 | |
| 0.9 | +0.1 | 17 47 21 | -28 09 | 8 | C | 18? | varies | |
| 1.0 | -0.1 | 17 48 30 | -28 09 | 8 | S | 15 | 0.6? | |
| 1.4 | -0.1 | 17 49 39 | -27 46 | 10 | S | 2? | ? | |
| 1.9 | +0.3 | 17 48 45 | -27 10 | 1.2 | S | 0.6 | 0.7 | |
| 3.7 | -0.2 | 17 55 26 | -25 50 | 14×11 | S | 2.3 | 0.65 | |
| 3.8 | +0.3 | 17 52 55 | -25 28 | 18 | S? | 4? | ? | |
| 4.2 | -3.5 | 18 08 55 | -27 03 | 28 | S | 3.2? | 0.6? | |
| 4.5 | +6.8 | 17 30 42 | -21 29 | 3 | S | 19 | 0.64 | Kepler, SN1604, 3C358 |
| 4.8 | +6.2 | 17 33 25 | -21 34 | 18 | S | 3 | 0.6 | |
| 5.2 | -2.6 | 18 07 30 | -25 45 | 18 | S | 2.6? | 0.6? | |
| 5.4 | -1.2 | 18 02 10 | -24 54 | 35 | C? | 35? | 0.2? | Milne 56 |
| 5.9 | +3.1 | 17 47 20 | -22 16 | 20 | S | 3.3? | 0.4? | |
| 6.1 | +1.2 | 17 54 55 | -23 05 | 30×26 | F | 4.0? | 0.3? | |
| 6.4 | -0.1 | 18 00 30 | -23 26 | 48 | C | 310 | varies | W28 |
| 6.4 | +4.0 | 17 45 10 | -21 22 | 31 | S | 1.3? | 0.4? | |
| 7.0 | -0.1 | 18 01 50 | -22 54 | 15 | S | 2.5? | 0.5? | |
| 7.7 | -3.7 | 18 17 25 | -24 04 | 22 | S | 11 | 0.32 | 1814-24 |
| 8.7 | -5.0 | 18 24 10 | -23 48 | 26 | S | 4.4 | 0.3 | |
| 8.7 | -0.1 | 18 05 30 | -21 26 | 45 | S? | 80 | 0.5 | (W30) |
| 9.8 | +0.6 | 18 05 08 | -20 14 | 12 | S | 3.9 | 0.5 | |
| 10.0 | -0.3 | 18 08 39 | -20 25 | 8? | ? | 2.9 | 0.8 | |
| 11.2 | -0.3 | 18 11 27 | -19 25 | 4 | C | 22 | 0.50 | |
| 11.4 | -0.1 | 18 10 47 | -19 05 | 8 | S? | 6 | 0.5 | |
| 12.0 | -0.1 | 18 12 11 | -18 37 | 7? | ? | 3.5 | 0.7 | |
| 13.3 | -1.3 | 18 19 20 | -18 00 | 70×40 | S? | ? | ? | |
| 13.5 | +0.2 | 18 14 14 | -17 12 | 5×4 | S | 3.5? | 1.0? | |
| 15.1 | -1.6 | 18 24 00 | -16 34 | 30×24 | S | 5.5? | 0.8? | |
| 15.9 | +0.2 | 18 18 52 | -15 02 | 7×5 | S? | 5 | 0.6? | |
| 16.2 | -2.7 | 18 28 50 | -16 11 | 17 | S | 2 | 0.5 | |
| 16.7 | +0.1 | 18 20 56 | -14 20 | 4 | C | 3.0 | 0.6 | |
| 16.8 | -1.1 | 18 25 20 | -14 46 | $30 \times 24?$ | ? | 2? | ? | |
| 17.4 | -2.3 | 18 30 55 | -14 52 | 24? | S | 4.8? | 0.8? | |
| 17.8 | -2.6 | 18 32 50 | -14 39 | 24 | S | 4.0? | 0.3? | |
| 18.8 | +0.3 | 18 23 58 | -12 23 | 17×11 | S | 33 | 0.4 | Kes 67 |
| 18.9 | -1.1 | 18 29 50 | -12 58 | 33 | C? | 37 | varies | |
| 20.0 | -0.2 | 18 28 07 | -11 35 | 10 | F | 10 | 0.0 | |
| 21.5 | -0.9 | 18 33 33 | -10 35 | 4 | C | 6? | 0.0 | |
| 21.8 | -0.6 | 18 32 45 | -10 08 | 20 | S | 69 | 0.5 | Kes 69 |
| 22.7 | -0.2 | 18 33 15 | -09 13 | 26 | S? | 33 | 0.6 | |
| 23.3 | -0.3 | 18 34 45 | -08 48 | 27 | S | 70 | 0.5 | W41 |
| 23.6 | +0.3 | 18 33 03 | -08 13 | 10? | ? | 8? | 0.3 | |
| 24.7 | -0.6 | 18 38 43 | -07 32 | 15? | S? | 8 | 0.5 | |
| 24.7 | +0.6 | 18 34 10 | -07 05 | 30×15 | C? | 20? | 0.2? | |

| <i>l</i> | <i>b</i> | RA (J2000.0) (h m s) | Dec (° ′) | size /arcmin | type | Flux at 1 GHz/Jy | spectral index | other name(s) |
|----------|----------|-------------------------|--------------|-----------------|------|---------------------|-------------------|-----------------------|
| 27.4 | +0.0 | 18 41 19 | -04 56 | 4 | S | 6 | 0.68 | 4C-04.71 |
| 27.8 | +0.6 | 18 39 50 | -04 24 | 50 × 30 | F | 30 | varies | |
| 28.6 | -0.1 | 18 43 55 | -03 53 | 13 × 9 | S | 3? | ? | |
| 28.8 | +1.5 | 18 39 00 | -02 55 | 100? | S? | ? | 0.4? | |
| 29.6 | +0.1 | 18 44 52 | -02 57 | 5 | S | 1.5? | 0.5? | |
| 29.7 | -0.3 | 18 46 25 | -02 59 | 3 | C | 10 | 0.7 | Kes 75 |
| 30.7 | -2.0 | 18 54 25 | -02 54 | 16 | ? | 0.5? | 0.7? | |
| 30.7 | +1.0 | 18 44 00 | -01 32 | 24 × 18 | S? | 6 | 0.4 | |
| 31.5 | -0.6 | 18 51 10 | -01 31 | 18? | S? | 2? | ? | |
| 31.9 | +0.0 | 18 49 25 | -00 55 | 7 × 5 | S | 24 | 0.55 | 3C391 |
| 32.0 | -4.9 | 19 06 00 | -03 00 | 60? | S? | 22? | 0.5? | 3C396.1 |
| 32.1 | -0.9 | 18 53 10 | -01 08 | 40? | C? | ? | ? | |
| 32.8 | -0.1 | 18 51 25 | -00 08 | 17 | S? | 11? | 0.2? | Kes 78 |
| 33.2 | -0.6 | 18 53 50 | -00 02 | 18 | S | 3.5 | varies | |
| 33.6 | +0.1 | 18 52 48 | +00 41 | 10 | S | 22 | 0.5 | Kes 79, 4C00.70, HC13 |
| 34.7 | -0.4 | 18 56 00 | +01 22 | 35 × 27 | C | 230 | 0.30 | W44, 3C392 |
| 36.6 | -0.7 | 19 00 35 | +02 56 | 25? | S? | ? | ? | |
| 36.6 | +2.6 | 18 48 49 | +04 26 | 17 × 13? | S | 0.7? | 0.5? | |
| 39.2 | -0.3 | 19 04 08 | +05 28 | 8 × 6 | S | 18 | 0.6 | 3C396, HC24, NRAO 593 |
| 39.7 | -2.0 | 19 12 20 | +04 55 | 120 × 60 | ? | 85? | 0.7? | W50, SS433 |
| 40.5 | -0.5 | 19 07 10 | +06 31 | 22 | S | 11 | 0.5 | |
| 41.1 | -0.3 | 19 07 34 | +07 08 | 4.5 × 2.5 | S | 22 | 0.48 | 3C397 |
| 42.8 | +0.6 | 19 07 20 | +09 05 | 24 | S | 3? | 0.5? | |
| 43.3 | -0.2 | 19 11 08 | +09 06 | 4 × 3 | S | 38 | 0.48 | W49B |
| 43.9 | +1.6 | 19 05 50 | +10 30 | 60? | S? | 8.6? | 0.2? | |
| 45.7 | -0.4 | 19 16 25 | +11 09 | 22 | S | 4.2? | 0.4? | |
| 46.8 | -0.3 | 19 18 10 | +12 09 | 17 × 13 | S | 14 | 0.5 | (HC30) |
| 49.2 | -0.7 | 19 23 50 | +14 06 | 30 | S? | 160? | 0.3? | (W51) |
| 53.6 | -2.2 | 19 38 50 | +17 14 | 33 × 28 | S | 8 | 0.75 | 3C400.2, NRAO 611 |
| 54.1 | +0.3 | 19 30 31 | +18 52 | 1.5 | F? | 0.5 | 0.1 | |
| 54.4 | -0.3 | 19 33 20 | +18 56 | 40 | S | 28 | 0.5 | (HC40) |
| 55.0 | +0.3 | 19 32 00 | +19 50 | 20 × 15? | S | 0.5? | 0.5? | |
| 55.7 | +3.4 | 19 21 20 | +21 44 | 23 | S | 1.4 | 0.6 | |
| 57.2 | +0.8 | 19 34 59 | +21 57 | 12? | S? | 1.8? | ? | (4C21.53) |
| 59.5 | +0.1 | 19 42 33 | +23 35 | 5 | S | 3? | ? | |
| 59.8 | +1.2 | 19 38 55 | +24 19 | 20 × 16? | ? | 1.6 | 0.5 | |
| 63.7 | +1.1 | 19 47 52 | +27 45 | 8 | F | 1.8 | 0.3 | |
| 65.1 | +0.6 | 19 54 40 | +28 35 | 90 × 50 | S | 6 | 0.6 | |
| 65.3 | +5.7 | 19 33 00 | +31 10 | 310 × 240 | S? | 52? | 0.6? | |
| 65.7 | +1.2 | 19 52 10 | +29 26 | 18 | ? | 5.1 | 0.6 | DA 495 |
| 67.7 | +1.8 | 19 54 32 | +31 29 | 9 | S | 1.4 | 0.3 | |
| 68.6 | -1.2 | 20 08 40 | +30 37 | 28 × 25? | ? | 0.7? | 0.0? | |
| 69.0 | +2.7 | 19 53 20 | +32 55 | 80? | ? | 120? | varies | CTB 80 |
| 69.7 | +1.0 | 20 02 40 | +32 43 | 16 | S | 1.6 | 0.8 | |
| 73.9 | +0.9 | 20 14 15 | +36 12 | 22? | S? | 9? | 0.3? | |

| l | b | RA (J2000.0) (h m s) | Dec ($^{\circ}$ $'$) | size /arcmin | type | Flux at 1 GHz/Jy | spectral index | other name(s) |
|-------|-------|-------------------------|---------------------------|-----------------|------|---------------------|-------------------|----------------------------|
| 74.0 | -8.5 | 20 51 00 | +30 40 | 230 × 160 | S | 210 | varies | Cygnus Loop |
| 74.9 | +1.2 | 20 16 02 | +37 12 | 8 × 6 | F | 9 | varies | CTB 87 |
| 76.9 | +1.0 | 20 22 20 | +38 43 | 12 × 9 | ? | 2 | 0.6 | |
| 78.2 | +2.1 | 20 20 50 | +40 26 | 60 | S | 340 | 0.5 | DR4, γ Cygni SNR |
| 82.2 | +5.3 | 20 19 00 | +45 30 | 95 × 65 | S | 120? | 0.5? | W63 |
| 84.2 | -0.8 | 20 53 20 | +43 27 | 20 × 16 | S | 11 | 0.5 | |
| 84.9 | +0.5 | 20 50 30 | +44 53 | 6 | S | 0.8 | 0.4 | |
| 85.4 | +0.7 | 20 50 40 | +45 22 | 24 | S | ? | 0.5? | |
| 85.9 | -0.6 | 20 58 40 | +44 53 | 24 | S | ? | 0.5? | |
| 89.0 | +4.7 | 20 45 00 | +50 35 | 120 × 90 | S | 220 | 0.40 | HB21 |
| 93.3 | +6.9 | 20 52 25 | +55 21 | 27 × 20 | S | 9 | 0.54 | DA 530, 4C(T)55.38.1 |
| 93.7 | -0.2 | 21 29 20 | +50 50 | 80 | S | 65 | 0.3 | CTB 104A, DA 551 |
| 94.0 | +1.0 | 21 24 50 | +51 53 | 30 × 25 | S | 15 | 0.44 | 3C434.1 |
| 106.3 | +2.7 | 22 27 30 | +60 50 | 60 × 24 | ? | 6 | 0.6 | |
| 109.1 | -1.0 | 23 01 35 | +58 53 | 28 | S | 20 | 0.50 | CTB 109 |
| 111.7 | -2.1 | 23 23 26 | +58 48 | 5 | S | 2720 | 0.77 | Cassiopeia A, 3C461 |
| 114.3 | +0.3 | 23 37 00 | +61 55 | 90 × 55 | S | 6? | 0.3? | |
| 116.5 | +1.1 | 23 53 40 | +63 15 | 80 × 60 | S | 11? | 0.8? | |
| 116.9 | +0.2 | 23 59 10 | +62 26 | 34 | S | 9? | 0.5? | CTB 1 |
| 119.5 | +10.2 | 00 06 40 | +72 45 | 90? | S | 36 | 0.6 | CTA 1 |
| 120.1 | +1.4 | 00 25 18 | +64 09 | 8 | S | 56 | 0.61 | Tycho, 3C10, SN1572 |
| 126.2 | +1.6 | 01 22 00 | +64 15 | 70 | S? | 7 | varies | |
| 127.1 | +0.5 | 01 28 20 | +63 10 | 45 | S | 13 | 0.6 | R5 |
| 130.7 | +3.1 | 02 05 41 | +64 49 | 9 × 5 | F | 33 | 0.10 | 3C58, SN1181 |
| 132.7 | +1.3 | 02 17 40 | +62 45 | 80 | S | 45 | 0.6 | HB3 |
| 156.2 | +5.7 | 04 58 40 | +51 50 | 110 | S | 5 | 0.5 | |
| 160.9 | +2.6 | 05 01 00 | +46 40 | 140 × 120 | S | 110 | 0.6 | HB9 |
| 166.0 | +4.3 | 05 26 30 | +42 56 | 55 × 35 | S | 7? | 0.4? | VRO 42.05.01 |
| 166.2 | +2.5 | 05 19 00 | +41 55 | 90 × 70 | S | 11 | 0.5 | OA 184 |
| 179.0 | +2.6 | 05 53 40 | +31 05 | 70 | S? | 7 | 0.4 | |
| 180.0 | -1.7 | 05 39 00 | +27 50 | 180 | S | 65 | varies | S147 |
| 182.4 | +4.3 | 06 08 10 | +29 00 | 50 | S | 1.2 | 0.4 | |
| 184.6 | -5.8 | 05 34 31 | +22 01 | 7 × 5 | F | 1040 | 0.30 | Crab Nebula, 3C144, SN1054 |
| 189.1 | +3.0 | 06 17 00 | +22 34 | 45 | C | 160 | 0.36 | IC443, 3C157 |
| 192.8 | -1.1 | 06 09 20 | +17 20 | 78 | S | 20? | 0.6? | PKS 0607+17 |
| 205.5 | +0.5 | 06 39 00 | +06 30 | 220 | S | 160 | 0.5 | Monoceros Nebula |
| 206.9 | +2.3 | 06 48 40 | +06 26 | 60 × 40 | S? | 6 | 0.5 | PKS 0646+06 |
| 260.4 | -3.4 | 08 22 10 | -43 00 | 60 × 50 | S | 130 | 0.5 | Puppis A, MSH 08-44 |
| 261.9 | +5.5 | 09 04 20 | -38 42 | 40 × 30 | S | 10? | 0.4? | |
| 263.9 | -3.3 | 08 34 00 | -45 50 | 255 | C | 1750 | varies | Vela (XYZ) |
| 266.2 | -1.2 | 08 52 00 | -46 20 | 120 | S | 50? | 0.3? | |
| 272.2 | -3.2 | 09 06 50 | -52 02 | 15? | S? | 0.4 | 0.6 | |
| 279.0 | +1.1 | 09 57 40 | -53 15 | 95 | S | 30? | 0.6? | |
| 284.3 | -1.8 | 10 18 15 | -59 00 | 24? | S | 11? | 0.3? | MSH 10-53 |
| 286.5 | -1.2 | 10 35 40 | -59 42 | 26 × 6 | S? | 1.4? | ? | |

| l | b | RA (J2000.0) (h m s) | Dec ($^{\circ}$ $'$) | size /arcmin | type | Flux at 1 GHz/Jy | spectral index | other name(s) |
|-------|-------|-------------------------|---------------------------|-----------------|------|---------------------|-------------------|---------------------|
| 289.7 | -0.3 | 11 01 15 | -60 18 | 18 × 14 | S | 6.2 | 0.2? | |
| 290.1 | -0.8 | 11 03 05 | -60 56 | 19 × 14 | S | 42 | 0.4 | MSH 11-61A |
| 291.0 | -0.1 | 11 11 54 | -60 38 | 15 × 13 | C | 16 | 0.29 | (MSH 11-62) |
| 292.0 | +1.8 | 11 24 36 | -59 16 | 12 × 8 | C | 15 | 0.4 | MSH 11-54 |
| 292.2 | -0.5 | 11 19 20 | -61 28 | 20 × 15 | S | 7? | 0.6? | |
| 293.8 | +0.6 | 11 35 00 | -60 54 | 20 | C | 5? | 0.6? | |
| 294.1 | -0.0 | 11 36 10 | -61 38 | 40 | S | >2? | ? | |
| 296.1 | -0.5 | 11 51 10 | -62 34 | 37 × 25 | S | 8? | 0.6? | |
| 296.5 | +10.0 | 12 09 40 | -52 25 | 90 × 65 | S | 48 | 0.5 | PKS 1209-51/52 |
| 296.8 | -0.3 | 11 58 30 | -62 35 | 20 × 14 | S | 9 | 0.6 | 1156-62 |
| 298.5 | -0.3 | 12 12 40 | -62 52 | 5? | ? | 5? | 0.4? | |
| 298.6 | -0.0 | 12 13 41 | -62 37 | 12 × 9 | S | 5? | 0.3 | |
| 299.2 | -2.9 | 12 15 13 | -65 30 | 18 × 11 | S | 0.5? | ? | |
| 299.6 | -0.5 | 12 21 45 | -63 09 | 13 | S | 1.0? | ? | |
| 301.4 | -1.0 | 12 37 55 | -63 49 | 37 × 23 | S | 2.1? | ? | |
| 302.3 | +0.7 | 12 45 55 | -62 08 | 17 | S | 5? | 0.4? | |
| 304.6 | +0.1 | 13 05 59 | -62 42 | 8 | S | 14 | 0.5 | Kes 17 |
| 308.1 | -0.7 | 13 37 37 | -63 04 | 13 | S | 1.2? | ? | |
| 308.8 | -0.1 | 13 42 30 | -62 23 | 30 × 20? | C? | 15? | 0.4? | |
| 309.2 | -0.6 | 13 46 31 | -62 54 | 15 × 12 | S | 7? | 0.4? | |
| 309.8 | +0.0 | 13 50 30 | -62 05 | 25 × 19 | S | 17 | 0.5 | |
| 310.6 | -0.3 | 13 58 00 | -62 09 | 8 | S | 5? | ? | Kes 20B |
| 310.8 | -0.4 | 14 00 00 | -62 17 | 12 | S | 6? | ? | Kes 20A |
| 311.5 | -0.3 | 14 05 38 | -61 58 | 5 | S | 3? | 0.5 | |
| 312.4 | -0.4 | 14 13 00 | -61 44 | 38 | S | 45 | 0.36 | |
| 315.4 | -2.3 | 14 43 00 | -62 30 | 42 | S | 49 | 0.6 | RCW 86, MSH 14-63 |
| 315.4 | -0.3 | 14 35 55 | -60 36 | 24 × 13 | ? | 8 | 0.4 | |
| 315.9 | -0.0 | 14 38 25 | -60 11 | 25 × 14 | S | 0.8? | ? | |
| 316.3 | -0.0 | 14 41 30 | -60 00 | 29 × 14 | S | 20? | 0.4 | (MSH 14-57) |
| 317.3 | -0.2 | 14 49 40 | -59 46 | 11 | S | 4.7? | ? | |
| 318.2 | +0.1 | 14 54 50 | -59 04 | 40 × 35 | S | >3.9? | ? | |
| 318.9 | +0.4 | 14 58 30 | -58 29 | 30 × 14 | C | 4? | 0.2? | |
| 320.4 | -1.2 | 15 14 30 | -59 08 | 35 | C | 60? | 0.4 | MSH 15-52, RCW 89 |
| 320.6 | -1.6 | 15 17 50 | -59 16 | 60 × 30 | S | ? | ? | |
| 321.9 | -1.1 | 15 23 45 | -58 13 | 28 | S | >3.4? | ? | |
| 321.9 | -0.3 | 15 20 40 | -57 34 | 31 × 23 | S | 13 | 0.3 | |
| 322.5 | -0.1 | 15 23 23 | -57 06 | 15 | C | 1.5 | 0.4 | |
| 323.5 | +0.1 | 15 28 42 | -56 21 | 13 | S | 3? | 0.4? | |
| 326.3 | -1.8 | 15 53 00 | -56 10 | 38 | C | 145 | varies | MSH 15-56 |
| 327.1 | -1.1 | 15 54 25 | -55 09 | 18 | C | 7? | ? | |
| 327.4 | +0.4 | 15 48 20 | -53 49 | 21 | S | 30? | 0.6 | Kes 27 |
| 327.4 | +1.0 | 15 46 48 | -53 20 | 14 | S | 1.9? | ? | |
| 327.6 | +14.6 | 15 02 50 | -41 56 | 30 | S | 19 | 0.6 | SN1006, PKS 1459-41 |
| 328.4 | +0.2 | 15 55 30 | -53 17 | 5 | F | 15 | 0.12 | (MSH 15-57) |
| 329.7 | +0.4 | 16 01 20 | -52 18 | 40 × 33 | S | >34? | ? | |

| l | b | RA (J2000.0) (h m s) | Dec ($^{\circ}$ $'$) | size /arcmin | type | Flux at 1 GHz/Jy | spectral index | other name(s) |
|-------|-------|-------------------------|---------------------------|-----------------|------|---------------------|-------------------|-------------------|
| 330.0 | +15.0 | 15 10 00 | –40 00 | 180? | S | 350? | 0.5? | Lupus Loop |
| 330.2 | +1.0 | 16 01 06 | –51 34 | 11 | S? | 5? | 0.3 | |
| 332.0 | +0.2 | 16 13 17 | –50 53 | 12 | S | 8? | 0.5 | |
| 332.4 | –0.4 | 16 17 33 | –51 02 | 10 | S | 28 | 0.5 | RCW 103 |
| 332.4 | +0.1 | 16 15 17 | –50 42 | 15 | S | 26 | 0.5 | MSH 16–51, Kes 32 |
| 335.2 | +0.1 | 16 27 45 | –48 47 | 21 | S | 16 | 0.5 | |
| 336.7 | +0.5 | 16 32 11 | –47 19 | 14 × 10 | S | 6 | 0.5 | |
| 337.0 | –0.1 | 16 35 57 | –47 36 | 1.5 | S | 1.5 | 0.6? | (CTB 33) |
| 337.2 | –0.7 | 16 39 28 | –47 51 | 6 | S | 2? | 0.7 | |
| 337.3 | +1.0 | 16 32 39 | –46 36 | 15 × 12 | S | 16 | 0.55 | Kes 40 |
| 337.8 | –0.1 | 16 39 01 | –46 59 | 9 × 6 | S | 18 | 0.5 | Kes 41 |
| 338.1 | +0.4 | 16 37 59 | –46 24 | 15? | S | 4? | 0.4 | |
| 338.3 | –0.0 | 16 41 00 | –46 34 | 8 | S | 7? | ? | |
| 338.5 | +0.1 | 16 41 09 | –46 19 | 9 | ? | 12? | ? | |
| 340.4 | +0.4 | 16 46 31 | –44 39 | 10 × 7 | S | 5 | 0.4 | |
| 340.6 | +0.3 | 16 47 41 | –44 34 | 6 | S | 5? | 0.4? | |
| 341.2 | +0.9 | 16 47 35 | –43 47 | 16 × 22 | C | 1.5? | 0.6? | |
| 341.9 | –0.3 | 16 55 01 | –44 01 | 7 | S | 2.5 | 0.5 | |
| 342.0 | –0.2 | 16 54 50 | –43 53 | 12 × 9 | S | 3.5? | 0.4? | |
| 342.1 | +0.9 | 16 50 43 | –43 04 | 10 × 9 | S | 0.5? | ? | |
| 343.0 | –6.0 | 17 25 00 | –46 30 | 250 | S | ? | ? | |
| 343.1 | –2.3 | 17 08 00 | –44 16 | 32? | C? | 8? | 0.5? | |
| 343.1 | –0.7 | 17 00 25 | –43 14 | 27 × 21 | S | 7.8 | 0.55 | |
| 344.7 | –0.1 | 17 03 51 | –41 42 | 10 | C? | 2.5? | 0.5 | |
| 345.7 | –0.2 | 17 07 20 | –40 53 | 6 | S | 0.6? | ? | |
| 346.6 | –0.2 | 17 10 19 | –40 11 | 8 | S | 8? | 0.5? | |
| 347.3 | –0.5 | 17 13 50 | –39 45 | 65 × 55 | S? | ? | ? | |
| 348.5 | –0.0 | 17 15 26 | –38 28 | 10? | S? | 10? | 0.4? | |
| 348.5 | +0.1 | 17 14 06 | –38 32 | 15 | S | 72 | 0.3 | CTB 37A |
| 348.7 | +0.3 | 17 13 55 | –38 11 | 17? | S | 26 | 0.3 | CTB 37B |
| 349.2 | –0.1 | 17 17 15 | –38 04 | 9 × 6 | S | 1.4? | ? | |
| 349.7 | +0.2 | 17 17 59 | –37 26 | 2.5 × 2 | S | 20 | 0.5 | |
| 350.0 | –2.0 | 17 27 50 | –38 32 | 45 | S | 26 | 0.4 | |
| 351.2 | +0.1 | 17 22 27 | –36 11 | 7 | C? | 5? | 0.4 | |
| 351.7 | +0.8 | 17 21 00 | –35 27 | 18 × 14 | S | 10? | ? | |
| 351.9 | –0.9 | 17 28 52 | –36 16 | 12 × 9 | S | 1.8? | ? | |
| 352.7 | –0.1 | 17 27 40 | –35 07 | 8 × 6 | S | 4 | 0.6 | |
| 353.9 | –2.0 | 17 38 55 | –35 11 | 13 | S | 1? | 0.5? | |
| 354.1 | +0.1 | 17 30 28 | –33 46 | 15 × 3? | C? | ? | varies? | |
| 354.8 | –0.8 | 17 36 00 | –33 42 | 19 | S | 2.8? | ? | |
| 355.6 | –0.0 | 17 35 16 | –32 38 | 8 × 6 | S | 3? | ? | |
| 355.9 | –2.5 | 17 45 53 | –33 43 | 13 | S | 8 | 0.5 | |
| 356.2 | +4.5 | 17 19 00 | –29 40 | 25 | S | 4 | 0.7 | |
| 356.3 | –0.3 | 17 37 56 | –32 16 | 11 × 7 | S | 3? | ? | |
| 356.3 | –1.5 | 17 42 35 | –32 52 | 20 × 15 | S | 3? | ? | |

| l | b | RA (J2000.0) (h m s) | Dec ($^{\circ}$ $'$) | size /arcmin | type | Flux at 1 GHz/Jy | spectral index | other name(s) |
|-------|------|-------------------------|---------------------------|-----------------|------|---------------------|-------------------|------------------|
| 357.7 | -0.1 | 17 40 29 | -30 58 | $8 \times 3?$ | ? | 37 | 0.4 | MSH 17-39 |
| 357.7 | +0.3 | 17 38 35 | -30 44 | 24 | S | 10 | 0.4? | |
| 358.0 | +3.8 | 17 26 00 | -28 36 | 38 | S | 1.5? | ? | |
| 359.0 | -0.9 | 17 46 50 | -30 16 | 23 | S | 23 | 0.5 | |
| 359.1 | -0.5 | 17 45 30 | -29 57 | 24 | S | 14 | 0.4? | |
| 359.1 | +0.9 | 17 39 36 | -29 11 | 12×11 | S | 5? | ? | |

Table II

Other names for SNRs

| | | |
|------------------------------|---|---|
| γ Cygni SNR G78.2+2.1 | DR4 G78.2+2.1 | NRAO 593 G39.2-0.3 NRAO 611 G53.6-2.2 |
| 1156-62 G296.8-0.3 | HB3 G132.7+1.3 | |
| 1814-24 G7.7-3.7 | HB9 G160.9+2.6 HB21 G89.0+4.7 | OA 184 G166.2+2.5 |
| 3C10 G120.1+1.4 | | PKS 0607+17 G192.8-1.1 |
| 3C58 G130.7+3.1 | HC13 G33.6+0.1 | PKS 0646+06 G206.9+2.3 |
| 3C144 G184.6-5.8 | HC24 G39.2-0.3 | PKS 1209-51/52 G296.5+10.0 |
| 3C157 G189.1+3.0 | (HC30) G46.8-0.3 | PKS 1459-41 G327.6+14.6 |
| 3C358 G4.5+6.8 | (HC40) G54.4-0.3 | |
| 3C391 G31.9+0.0 | | Puppis A G260.4-3.4 |
| 3C392 G34.7-0.4 | IC443 G189.1+3.0 | |
| 3C396 G39.2-0.3 | | R5 G127.1+0.5 |
| 3C396.1 G32.0-4.9 | Kepler G4.5+6.8 | |
| 3C397 G41.1-0.3 | | RCW 86 G315.4-2.3 |
| 3C400.2 G53.6-2.2 | Kes 17 G304.6+0.1 | RCW 89 G320.4-1.2 |
| 3C434.1 G94.0+1.0 | Kes 20A G310.6-0.3 | RCW 103 G332.4-0.4 |
| 3C461 G111.7-2.1 | Kes 20B G310.8-0.4 Kes 27 G327.4+0.4 | RCW 114 G343.0-6.0 |
| 4C-04.71 G27.4+0.0 | Kes 32 G332.4+0.1 | |
| 4C00.70 G33.6+0.1 | Kes 40 G337.3+1.0 | S147 G180.0-1.7 |
| (4C21.53) G57.2+0.8 | Kes 41 G337.8-0.1 | SN1006 G327.6+14.6 |
| 4C(T)55.38.1 G93.3+6.9 | Kes 67 G18.8+0.3 Kes 69 G21.8-0.6 | SN1054 G184.6-5.8 SN1181 G130.7+3.1 |
| CTA 1 G119.5+10.2 | Kes 75 G29.7-0.3 Kes 78 G32.8-0.1 Kes 79 G33.6+0.1 | SN1572 G120.1+1.4 SN1604 G4.5+6.8 |
| CTB 1 G116.9+0.2 | | SS433 G39.7-2.0 |
| (CTB 33) G337.0-0.1 | | |
| CTB 37A G348.5+0.1 | Lupus Loop G330.0+15.0 | |
| CTB 37B G348.7+0.3 | | Sgr A East G0.0+0.0 |
| CTB 80 G69.0+2.7 | MSH 08-44 G260.4-3.4 | |
| CTB 87 G74.9+1.2 | MSH 10-53 G284.3-1.8 | Tycho G120.1+1.4 |
| CTB 104A G93.7-0.2 | MSH 11-54 G292.0+1.8 | |
| CTB 109 G109.1-1.0 | MSH 11-61A G290.1-0.8 (MSH 11-62) G291.0-0.1 (MSH 14-57) G316.3-0.0 | Vela (XYZ) G263.9-3.3 |
| Cassiopeia A G111.7-2.1 | MSH 14-63 G315.4-2.3 | VRO 42.05.01 G166.0+4.3 |
| Crab Nebula G184.6-5.8 | MSH 15-52 G320.4-1.2 MSH 15-56 G326.3-1.8 | W28 G6.4-0.1 (W30) G8.7-0.1 |
| Cygnus Loop G74.0-8.5 | (MSH 15-57) G328.4+0.2 MSH 16-51 G332.4+0.1 MSH 17-39 G357.7-0.1 | W41 G23.3-0.3 W44 G34.7-0.4 W49B G43.3-0.2 W50 G39.7-2.0 (W51) G49.2-0.7 W63 G82.2+5.3 |
| DA 495 G65.7+1.2 | | |
| DA 530 G93.3+6.9 | | |
| DA 551 G93.7-0.2 | Milne 56 G5.4-1.2 | |
| Monoceros Nebula G205.5+0.5 | | |

Journals

| | |
|-------|--|
| A&A | Astronomy & Astrophysics |
| A&AS | Astronomy & Astrophysics Supplement |
| AJ | Astronomical Journal |
| ApJ | Astrophysical Journal |
| ApJS | Astrophysical Journal Supplement |
| AREp | Astronomy Reports (Astronomicheskii Zhurnal translation) |
| AstL | Astronomy Letters |
| AuJPh | Australian Journal of Physics |
| AuJPA | Australian Journal of Physics Astrophysical Supplement |
| JApA | Journal of Astrophysics & Astronomy |
| JRASC | Journal of the Royal Astronomical Society of Canada |
| MNRAS | Monthly Notices of the Royal Astronomical Society |
| Natur | Nature |
| PASAu | Proceedings of the Astronomical Society of Australia |
| PASJ | Publications of the Astronomical Society of Japan |
| PASP | Publications of the Astronomical Society of the Pacific |
| RMxAA | Review of Mexican Astronomy & Astrophysics |
| Sci | Science |
| SvAL | Soviet Astronomy Letters |

Proceedings

SNRISM is *Supernova Remnants and the Interstellar Medium* (IAU Colloquium 101), eds Roger R.S. & Landecker T.L., (Cambridge University Press), 1988.

Radio Telescopes

| | |
|-------|---|
| 5km | Cambridge 5-km Telescope |
| 6C | Cambridge low frequency northern survey |
| ATCA | Australia Telescope Compact Array |
| BIMA | Berkeley–Illinois–Maryland Array |
| CLFST | Cambridge Low Frequency Synthesis Telescope |
| DRAO | Dominion Radio Astrophysical Observatory |
| FIRST | Fleurs Synthesis Telescope |
| MOST | Molonglo Observatory Synthesis Telescope |
| NRAO | National Radio Astronomy Observatory |
| NRO | Nobeyama Radio Observatory |
| TPT | Clark Lake TPT telescope |
| VLA | Very Large Array |
| VRO | Vermillion River Observatory |
| WSRT | Westerbork Synthesis Radio Telescope |

Satellites

| | |
|--------|---|
| HST | Hubble Space Telescope |
| ISO | Infrared Space Observatory |
| X-ray: | |
| ASCA | Advanced Satellite for Cosmology and Astrophysics |
| EXOSAT | European X-ray Observatory Satellite |
| ROSAT | Röntgensatellit |
| XMM | X-ray Multi-Mirror |