

G0.0+0.0

RA: 17^h45^m44^s
Dec: -29°00'

1-GHz flux/Jy: 100?
Spectral index: 0.8?

Sgr A East
Size/arcmin: 3.5 × 2.5
Type: S

Radio: Non-thermal shell, in complex region, interacting with molecular material to the west.

X-ray: Diffuse emission, centrally peaked.

Point sources: Compact X-ray source.

References:

- Ekers *et al.* 1983, A&A, 122, 143. VLA at 1.4 GHz and 5 GHz (both 5'' × 8''), $S_{1.4 \text{ GHz}} = 77 \text{ Jy}$, $S_{5.0 \text{ GHz}} = 31 \text{ Jy}$.
Pedlar *et al.* 1989, ApJ, 342, 769. VLA at 332 MHz (12''), 1.4 GHz (1''.3 × 2''.5) and 5 GHz (1''.3 × 2''.5).
Mezger *et al.* 1989, A&A, 209, 337. Nearby molecular material.
Ho *et al.* 1991, Nature, 350, 309. VLA of NH₃ emission from surroundings.
Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (17'' × 33'').
Serabyn *et al.* 1992, ApJ, 395, 166. Nearby molecular material.
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 88'').
Yusef-Zadeh & Mehringer 1995, ApJ, 452, L37. VLA of nearby H₂O masers.
Yusef-Zadeh *et al.* 1999, ApJ, 512, 230. OH maser observations.
Coil & Ho 2000, ApJ, 533, 245. NH₃ observations of surroundings.
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24'' × 43'').
Fatuzzo *et al.* 2001, ApJ, 549, 293. Electron-positron lines from the vicinity.
Yusef-Zadeh *et al.* 2001, ApJ, 560, 749. Observations of nearby molecular hydrogen.
Maeda *et al.* 2002, ApJ, 570, 671. Chandra observations.
Sakano *et al.* 2003, AN, 324 (No S1), 197. XMM observations.
Roy & Rao 2004, MNRAS, 349, L25. GMRT at 620 MHz (6''.6 × 11''.4).
Sakano *et al.* 2004, MNRAS, 350, 129. XMM observations.
Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (1'' × 2''.2).
Park *et al.* 2005, ApJ, 631, 964. Chandra observations.
Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.
Koyama *et al.* 2007, PASJ, 59, S237. Suzaku observations.
Lee *et al.* 2008, ApJ, 674, 247. Molecular H₂ observations of surroundings.
Sjouwerman & Pihlström 2008, ApJ, 681, 1287. VLA at 1.7 GHz of OH masers.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G0.3+0.0

RA: 17^h46^m15^s
Dec: -28°38'

1-GHz flux/Jy: 22
Spectral index: 0.6

Size/arcmin: 15 × 8
Type: S

Has been called G0.33+0.04 and G0.4+0.1.

Radio: Bilateral shell, near Galactic Centre.

References:

- Kassim & Frail 1996, MNRAS, 283, L51. VLA at 333 MHz (23'' × 42''), plus review of flux densities and other observations.
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (30'').
Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').

G0.9+0.1

RA: 17^h47^m21^s
Dec: -28°09'

1-GHz flux/Jy: 18?
Spectral index: varies

Size/arcmin: 8
Type: C

Radio: Flat spectrum core within steep spectrum shell.

X-ray: Central core, with non-thermal spectrum.

References:

- Helfand & Becker 1987, ApJ, 314, 203. VLA at 1.4 and 5 GHz, and Einstein observations.
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 91'').
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Mereghetti *et al.* 1998, A&A, 331, L77. X-ray detection.
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').
Sidoli *et al.* 2000, A&A, 361, 719. BeppoSAX observations.
Gaensler *et al.* 2001, ApJ, 556, L107. Chandra observations.
Porquet *et al.* 2003, A&A, 401, 197. XMM observations.
Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.6 × 11''.3).
Aharanion *et al.* 2005, A&A, 432, L25. γ-ray detection.
Dubner *et al.* 2008, A&A, 487, 1033. ATCA and VLA at 1.4 GHz (1''.2 × 2''.5 : $S = 8.3 \pm 0.7$ Jy), 5 GHz (1''.6 × 2''.5) and 8.3 GHz (0''.8 × 1''.5).

G1.0–0.1

RA: 17^h48^m30^s
Dec: -28°09'

1-GHz flux/Jy: 15
Spectral index: 0.6?

Size/arcmin: 8
Type: S

Has been called G1.05–0.1 and G1.05–0.15.

Radio: Incomplete shell, to the S of Sgr D.

X-ray: Possibly detected.

References:

- Downes *et al.* 1979, A&AS, 35, 1. Review of flux densities.
Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (64'' × 100'': $S = 12.3$ Jy).
Liszt 1992, ApJS, 82, 495. VLA at 1.6 GHz (13'' × 23'').
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 91'').
Mehringer *et al.* 1998, ApJ, 493, 274. VLA at 1.6 GHz (15'' × 28'') and 5 GHz, including masers observations.
Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').
Sidoli *et al.* 2001, A&A, 372, 651. BeppoSAX possible detection.
Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G1.4–0.1

RA: 17^h49^m39^s
Dec: -27°46'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 10
Type: S

Radio: Shell, brightest in E.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43'' × 92'': $S = 2$ Jy).
Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (2''.7 × 2''.4 : $S = 4.2 \pm 0.5$).
Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.2 × 12''.2).
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G1.9+0.3

RA: 17^h48^m45^s
Dec: -27°10'

1-GHz flux/Jy: 0.6
Spectral index: 0.6

Size/arcmin: 1.5
Type: S

Radio: Shell, brighter to the N, brightening.

X-ray: Shell, with bright limbs to E and W.

References:

- Green & Gull 1984, Nature, 312, 527. VLA at 5 GHz ($4'' \times 2''$).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 94''$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz ($8''/3 \times 12''$).
 Nord *et al.* 2004, AJ, 128, 1646. VLA at 330 MHz ($7'' \times 12''$).
 Green 2004, BASI, 32, 335. VLA at 1.5 GHz ($7''/2 \times 9''.4$).
 Reynolds *et al.* 2008, ApJ, 680, L41. Chandra observations.
 Green *et al.* 2008, MNRAS, 387, L54. VLA at 4.86 GHz ($4'' \times 10''$), for expansion studies.
 Murphy *et al.* 2008, MNRAS, 389, L23. MOST at 843 MHz for flux increase.

G3.7-0.2

RA: 17^h55^m26^s
Dec: -25°50'

1-GHz flux/Jy: 2.3
Spectral index: 0.65

Size/arcmin: 14 × 11
Type: S

Has been called G003.8-00.3.

Radio: Double arc.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43'' \times 99''$: $S=2.4$ Jy).
 Gaensler 1998, ApJ, 493, 781. VLA at 1.4 GHz ($9'' \times 15''$: $S=1.7 \pm 0.1$ Jy).
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz ($8''/4 \times 11''.4$).

G3.8+0.3

RA: 17^h52^m55^s
Dec: -25°28'

1-GHz flux/Jy: 3?
Spectral index: 0.6

Size/arcmin: 18
Type: S?

Radio: Incomplete shell.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43'' \times 100''$: $S=3.5$ Jy).
 Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz ($27'' \times 17''$: $S=6.0 \pm 0.4$).

G4.2-3.5

RA: 18^h08^m55^s
Dec: -27°03'

1-GHz flux/Jy: 3.2?
Spectral index: 0.6?

Size/arcmin: 28
Type: S

Radio: Elongated shell.

References:

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4''.3$).

G4.5+6.8

RA: 17^h30^m42^s
Dec: -21°29'

1-GHz flux/Jy: 19
Spectral index: 0.64

Kepler, SN1604, 3C358

Size/arcmin: 3
Type: S

This is the remnant of Kepler's SN of AD1604.

Radio: Incomplete shell, brighter to the N.

Optical: Faint filaments.

X-ray: Shell, brighter to the N.

Distance: Optical expansion and proper motion indicates about 2.9 kpc, HI observations suggest 3.4 to 6.4 kpc.

References:

- van den Bergh & Kamper 1977, ApJ, 218, 617. Optical proper motions.
Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
White & Long 1983, ApJ, 264, 196. Einstein observations.
Matsui *et al.* 1984, ApJ, 287, 295. VLA at 1.4 (2''/5 × 3'2) and 5 GHz (3''/2 × 4''/8) and Einstein image (5'').
Dickel *et al.* 1988, ApJ, 330, 254. VLA at 1.4 (1''/2 × 2''/3) and 5 GHz (0''.6 × 1''.0) at two epochs.
Smith *et al.* 1989, ApJ, 347, 925. EXOSAT observations.
Hatsukade *et al.* 1990, PASJ, 42, 279. X-ray spectrum.
Blair *et al.* 1991, ApJ, 366, 484. Optical imaging and spectroscopy.
Bandiera & van den Bergh 1991, ApJ, 374, 186. Optical changes.
van den Bergh 1991, PASP, 103, 194. Optical imaging.
Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Hughes 1999, ApJ, 527, 298. ROSAT and Einstein image comparison for expansion studies.
Reynoso & Goss 1999, AJ, 118, 926. VLA at 1.4 GHz (23'' × 13'') for HI studies.
Kinugasa & Tsunemi 1999, PASJ, 51, 239. ASCA observations.
Gerardy & Fesen 2001, AJ, 121, 2781. IR spectroscopy and imaging.
DeLaney *et al.* 2002, ApJ, 580, 914. VLA at 1.3 to 1.5 GHz and 5 GHz (7''.2) for spectral index studies.
Morgan *et al.* 2003, ApJ, 597, L33. Sub-mm dust observations.
Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
Cassam-Chenai *et al.* 2004, A&A, 414, 545. XMM observations.
Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
Riesgo & López 2005, RMxAA, 41, 57. Optical observations of filament (previously classified as PN, H 2-12).
Blair *et al.* 2007, ApJ, 662, 998. Spitzer IR observations.
Reynolds *et al.* 2007, ApJ, 668, L135. Chandra observations.
Sankrit *et al.* 2008, AJ, 135, 538. HST observations.
Aharonian *et al.* 2008, A&A, 488, 219. γ -ray upper limit.
Enomoto *et al.* 2008, ApJ, 683, 383. γ -ray upper limit.
Katsuda *et al.* 2008, ApJ, 689, 225. Chandra proper motion studies.
Vink *et al.* 2008, ApJ, 689, 231. Chandra proper motion studies.

G4.8+6.2

RA: 17^h33^m25^s
Dec: -21°34'

1-GHz flux/Jy: 3
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G4.5+6.2.

Radio: Faint shell.

References:

- Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (2''.2 × 1''.3 : S = 5.5 ± 1.2 Jy), and NVSS at 1.4 GHz.

G5.2-2.6

RA: 18^h07^m30^s
Dec: -25°45'

1-GHz flux/Jy: 2.6?
Spectral index: 0.6?

Size/arcmin: 18
Type: S

Radio: Poorly resolved shell.

References:

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4''.3).

G5.4–1.2

Milne 56

RA: 18^h02^m10^s
Dec: −24°54'

1-GHz flux/Jy: 35?
Spectral index: 0.2?

Size/arcmin: 35
Type: C?

Part been called G5.3–1.0. Has been suggested that this is not a SNR.

Radio: Incomplete shell, including wide ‘v’ of emission to east with small flat-spectrum source at apex.

Optical: Detected.

X-ray: Pulsar detected, with faint extension.

Distance: HI absorption suggests >4.3 kpc.

Point sources: Pulsar nearby, in flat spectrum source.

References:

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3' : $S=38$ Jy).
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
 Dickey & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'8 : $S=21.9 \pm 2.4$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.
 Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 and 5 GHz.
 Helfand & Becker 1985, Nature, 313, 118. Suggesting it is not a SNR.
 Manchester *et al.* 1985, MNRAS, 212, 975. Pulsar detection.
 Caswell *et al.* 1987, MNRAS, 225, 329. MOST at 843 MHz (42'' \times 110'').
 Frail & Kulkarni 1991, Nature, 352, 785. Pulsar and remnant association.
 Manchester *et al.* 1991, MNRAS, 253, 7P. Pulsar and remnant association.
 Milne *et al.* 1992, MNRAS, 255, 707. Parkes 64-m at 4.75 (4'5 : $S=30.8 \pm 2.1$ Jy) and 8.4 GHz (3' : $S=24 \pm 3$ Jy), including polarisation.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'0 \times 4'9 : $S=38$ Jy).
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz (68'' \times 73''), plus HI absorption.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gaensler & Frail 2000, Nature, 406, 158. Pulsar observations, including proper motion.
 Kaspi *et al.* 2001, ApJ, 562, L163. X-ray detection of pulsar, and upper limit for remnant.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.
 Blazek *et al.* 2006, ApJ, 652, 1523. Proper motion study of pulsar.
 Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion study of pulsar.

G5.5+0.3

RA: 17^h57^m04^s
Dec: −24°00'

1-GHz flux/Jy: 5.5
Spectral index: 0.7

Size/arcmin: 15 \times 12
Type: S

Has been called G5.55+0.32.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S=14.3 \pm 0.3$ Jy), plus other observations.

G5.9+3.1

RA: 17^h47^m20^s
Dec: −22°16'

1-GHz flux/Jy: 3.3?
Spectral index: 0.4?

Size/arcmin: 20
Type: S

Radio: Asymmetric shell.

References:

- Reich *et al.* 1988, in SNRISM, p293. Effelsberg 100-m at 2.7 GHz (4'3).

G6.1+0.5

RA: 17^h57^m29^s
Dec: -23°25'

1-GHz flux/Jy: 4.5
Spectral index: 0.9

Size/arcmin: 18 × 12
Type: S

Has been called G6.10+0.53.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 13.4 \pm 0.2$ Jy), plus other observations.

G6.1+1.2

RA: 17^h54^m55^s
Dec: -23°05'

1-GHz flux/Jy: 4.0?
Spectral index: 0.3?

Size/arcmin: 30 × 26
Type: F

Has been called G6.1+1.15.

Radio: Faint, diffuse emission.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

G6.4–0.1		W28
RA: 18 ^h 00 ^m 30 ^s	1-GHz flux/Jy: 310	Size/arcmin: 48
Dec: −23°26'	Spectral index: varies	Type: C
Has been called G6.6–0.2.		
Radio: Several non-thermal sources in a ring, with flat spectrum core.		
Optical: Diffuse emission.		
X-ray: Diffuse emission from most of the remnant.		
Point sources: Young pulsar near edge of remnant, but not thought to be related.		
Distance: HI observations suggest 1.9 kpc.		
References:		
Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10 GHz (3').		
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical observations.		
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).		
Dopita <i>et al.</i> 1977, ApJ, 214, 179. Some optical line ratios.		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).		
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.		
Andrews <i>et al.</i> 1983, ApJ, 266, 684. VLA at 1.4 GHz (8'') 4.9 GHz (3'') and 15 GHz (1'') of central region only, plus Einstein image of central region.		
Bohigas <i>et al.</i> 1983, RMxAA, 8, 155. Optical spectra.		
Odenwald <i>et al.</i> 1984, AJ, 90, 310. VLA of central component.		
Long <i>et al.</i> 1991, ApJ, 373, 567. Einstein and optical observations.		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'1 × 4'4 : S = 660 Jy).		
Kaspi <i>et al.</i> 1993, ApJ, 409, L57. Pulsar association.		
Frail <i>et al.</i> 1993, Nature, 365, 136. VLA at 327 MHz (smoothed to 65''), plus pulsar association.		
Frail <i>et al.</i> 1994, ApJ, 424, L111. VLA of associated OH masers.		
Esposito <i>et al.</i> 1996, ApJ, 461, 820. Possible associated γ -ray emission.		
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH maser emission.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Claussen <i>et al.</i> 1997, ApJ, 489, 143. VLA of OH masers.		
Claussen <i>et al.</i> 1999, ApJ, 522, 349. High resolution observations of OH masers.		
Arikawa <i>et al.</i> 1999, PASJ, 51, L7. Observations of shocked CO.		
Yusef-Zadeh <i>et al.</i> 2000, ApJ, 540, 842. VLA at 327 MHz (2'6 × 5'5) and 1.48 GHz (40'' × 65'').		
Dubner <i>et al.</i> 2000, AJ, 120, 1933. VLA at 328 MHz (97'' × 52'' : S = 425 ± 40 Jy) and 1415 MHz (88'' × 48'' : S = 246 ± 20 Jy), and comparison with other observations.		
Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.		
Rowell <i>et al.</i> 2000, A&A, 359, 337. Upper limit on high energy γ -rays.		
Douvion <i>et al.</i> 2001, A&A, 373, 281. ISO observations.		
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.		
Velázquez <i>et al.</i> 2002, AJ, 124, 2145. Parkes 64-m at 1.4 GHz (15') for HI.		
Rho & Borkowski 2002, ApJ, 575, 201. ROSAT and ASCA observations.		
Claussen <i>et al.</i> 2002, ApJ, 580, 909. Observations of nearby source.		
Yusef-Zadeh <i>et al.</i> 2003, ApJ, 583, 267. OH observations.		
Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz of associated OH masers.		
Mavromatakis <i>et al.</i> 2004, A&A, 426, 567. Optical observations.		
Reach <i>et al.</i> 2005, ApJ, 618, 297. Molecular lines and near IR observations.		
Hoffman <i>et al.</i> 2005, ApJ, 620, 257. OH maser observations.		
Kawasaki <i>et al.</i> 2005, ApJ, 631, 935. ASCA observations.		
Neufeld <i>et al.</i> 2007, ApJ, 664, 890. Spitzer IR observations.		
Aharonian <i>et al.</i> 2008, A&A, 481, 401. γ -ray observations.		
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.		
Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.		

G6.4+4.0		
RA: 17 ^h 45 ^m 10 ^s	1-GHz flux/Jy: 1.3?	Size/arcmin: 31
Dec: −21°22'	Spectral index: 0.4?	Type: S

Radio: Faint asymmetric shell.

References:

Reich *et al.* 1988, in SNRISM, p293. Effelsberg 100-m at 2.7 GHz (4'3).

G6.5–0.4

RA: 18^h02^m11^s
Dec: −23°34'

1-GHz flux/Jy: 27
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G6.51–0.48, and part has been called G6.67–0.42.

Radio: Shell.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 330 MHz ($2'.6 \times 5'.5$) and 1.4 GHz ($0'.7 \times 1'.1$).
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 60.8 \pm 0.4$ Jy), plus other observations.
Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.

G7.0–0.1

RA: 18^h01^m50^s
Dec: −22°54'

1-GHz flux/Jy: 2.5?
Spectral index: 0.5?

Size/arcmin: 15
Type: S

Has been called G7.06–0.12.

Radio: Double rim, brightest in W, confused by bright HII region M20 in SE.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz ($2'.6 \times 5'.5$) and 1.48 GHz ($40'' \times 65''$).
Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz ($97'' \times 52''$) and 1415 MHz ($88'' \times 48''$).

G7.2+0.2

RA: 18^h01^m07^s
Dec: −22°38'

1-GHz flux/Jy: 2.8
Spectral index: 0.6

Size/arcmin: 12
Type: S

Has been called G7.20+0.20.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 5.2 \pm 0.2$ Jy), plus other observations.

G7.7–3.7

RA: 18^h17^m25^s
Dec: −24°04'

1-GHz flux/Jy: 11
Spectral index: 0.32

1814–24
Size/arcmin: 22
Type: S

Radio: Shell, with high polarisation.

References:

Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'.4$) and 5 GHz ($4'.4$).
Milne *et al.* 1986, MNRAS, 223, 487. MOST at 843 MHz ($44'' \times 108''$) and Parkes 64-m at 8.4 GHz ($3' : S = 4.6 \pm 0.5$ Jy), with polarisation, plus review of flux densities.
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ($70'' \times 35'' : S = 9.9 \pm 0.1$ Jy), including polarisation.

G8.3–0.0

RA: 18^h04^m34^s
Dec: −21°49'

1-GHz flux/Jy: 1.2
Spectral index: 0.6

Size/arcmin: 5 × 4
Type: S

Has been called G8.31–0.09.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 2.3 \pm 0.1$ Jy), plus other observations.
Higashi *et al.* 2008, ApJ, 683, 957. γ-ray detection.

G8.7–5.0

RA: 18^h24^m10^s
Dec: −23°48'

1-GHz flux/Jy: 4.4
Spectral index: 0.3

Size/arcmin: 26
Type: S

Radio: Asymmetric shell.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

G8.7–0.1

RA: 18^h05^m30^s
Dec: −21°26'

1-GHz flux/Jy: 80
Spectral index: 0.5

(W30)
Size/arcmin: 45
Type: S?

Has been called G8.6–0.1.

Radio: Clumpy non-thermal shell, with low-frequency turnover.

X-ray: Northern edge detected.

Point sources: Pulsar inside western edge.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz ($7'.2 \times 9'.7 : S = 190 \pm 50$ Jy).
 Kassim & Weiler 1990, Nature, 343, 146. VLA at 327 MHz ($3'.0 \times 3'.7$).
 Kassim & Weiler 1990, ApJ, 360, 184. VLA at 327 MHz ($2'.8 \times 4'.1 : S = 129 \pm 11$ Jy), and part at 1.4 GHz ($0'.9 \times 1'.8$), plus review of flux densities.
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz ($37'' \times 55''$).
 Finley & Ögelman 1994, ApJ, 434, L25. ROSAT observations, including pulsar.
 Aharanion *et al.* 2005, Science, 307, 1938. γ-ray detection.
 Aharanion *et al.* 2006, ApJ, 636, 777. γ-ray observations.
 Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
 Briskin *et al.* 2006, ApJ, 652, 554. Pulsar proper motion.

G8.9+0.4

RA: 18^h03^m58^s
Dec: −21°03'

1-GHz flux/Jy: 9
Spectral index: 0.6

Size/arcmin: 24
Type: S

Has been called G8.90+0.40.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 18.2 \pm 0.5$ Jy), plus other observations.

G9.7–0.0

RA: 18^h07^m22^s
Dec: −20°35'

1-GHz flux/Jy: 3.7
Spectral index: 0.6

Size/arcmin: 15×11
Type: S

Has been called G9.7–0.1 and G9.70–0.06.

Radio: Shell.

References:

Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 6.5 \pm 0.2$ Jy), plus other observations.

G9.8+0.6

RA: 18^h05^m08^s
Dec: −20°14'

1-GHz flux/Jy: 3.9
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Asymmetric shell.

References:

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2/6).
 Caswell 1983, MNRAS, 204, 833. Molonglo at 408 MHz (3': $S=5.8\pm0.6$ Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41'' \times 63'': $S=3.5\pm0.4$ Jy).
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G9.9−0.8

RA: 18^h10^m41^s
Dec: −20°43'

1-GHz flux/Jy: 6.7
Spectral index: 0.4

Size/arcmin: 12
Type: S

Has been called G9.95−0.81.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=11.0\pm0.3$ Jy), plus other observations.

G10.5−0.0

RA: 18^h09^m08^s
Dec: −19°47'

1-GHz flux/Jy: 0.9
Spectral index: 0.6

Size/arcmin: 6
Type: S

Has been called G10.59−0.04.

Radio: Partial shell.

X-ray: Detected.

References:

- Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=1.4\pm0.1$ Jy), plus other observations.

G11.0−0.0

RA: 18^h10^m04^s
Dec: −19°25'

1-GHz flux/Jy: 1.3
Spectral index: 0.6

Size/arcmin: 11 \times 9
Type: S

Has been called G11.0+0.0 and G11.03−0.05.

Radio: Partial shell.

X-ray: Diffuse emission.

References:

- Bamba *et al.* 2003, ApJ, 589, 253. ASCA observations.
 Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=3.1\pm0.2$ Jy), plus other observations.

G11.1−1.0

RA: 18^h14^m03^s
Dec: −19°46'

1-GHz flux/Jy: 5.8
Spectral index: 0.6

Size/arcmin: 18 \times 12
Type: S

Has been called G11.2−1.1 and G11.17−1.04.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=11.0\pm0.3$ Jy), plus other observations.

G11.1–0.7

RA: 18^h12^m46^s
Dec: −19°38'

1-GHz flux/Jy: 1.0
Spectral index: 0.7

Size/arcmin: 11 × 7
Type: S

Has been called G11.15–0.71.

Radio: Partial shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=2.3 \pm 0.1$ Jy), plus other observations.

G11.1+0.1

RA: 18^h09^m47^s
Dec: −19°12'

1-GHz flux/Jy: 2.3
Spectral index: 0.4

Size/arcmin: 12 × 10
Type: S

Has been called G11.18+0.11.

Radio: Shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=3.5 \pm 0.2$ Jy), plus other observations.

G11.2–0.3

RA: 18^h11^m27^s
Dec: −19°25'

1-GHz flux/Jy: 22
Spectral index: 0.6

Size/arcmin: 4
Type: C

Probably associated with the SN of AD386.

Radio: Symmetrical clumpy shell, with flatter spectrum core.

X-ray: Shell, with hard spectrum centrally brightened region around pulsar.

Point sources: Central pulsar.

Distance: HI absorption indicates 4.4 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Radhakrishnan *et al.* 1972, ApJS, 24, 49. HI absorption.
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20'' × 25'') and Einstein observations, with review of flux densities.
Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz, plus HI absorption, Einstein observations.
Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26.5'': $S=4.04 \pm 0.24$ Jy).
Green *et al.* 1988, MNRAS, 231, 735. VLA at 1.4 and 5 GHz.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.2' × 4.1'': $S=39$ Jy).
Reynolds *et al.* 1994, MNRAS, 271, L1. ROSAT image and spectra.
Vasisht *et al.* 1996, ApJ, 456, L59. ASCA observations.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Torii *et al.* 1997, ApJ, 489, L145. ASCA detection of pulsar.
Torii *et al.* 1999, ApJ, 523, L69. X-ray timing observations of pulsar.
Kothes & Reich 2001, A&A, 372, 627. Effelsberg 100-m at 4.25, 10.45, 14.9 and 32 GHz (2.5', 1.1', 0.86', 0.45': $S=9.6 \pm 0.5, 6.3 \pm 0.4, 5.7 \pm 0.4, 3.8 \pm 0.4$).
Kaspi *et al.* 2001, ApJ, 560, 371. Chandra observations.
Tam *et al.* 2002, ApJ, 572, 202. VLA at 1.4/1.5 GHz (2.6' × 1.8'': $S=16.6 \pm 0.9$ Jy) and 5 GHz (2.1' × 1.5'': $S=8.4 \pm 0.9$ Jy) for spectral studies.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 14.7 GHz.
Tam & Roberts 2003, ApJ, 598, L27. Multi-epoch VLA observations at 1.4/1.5 GHz and 5 GHz, for expansion studies.
Roberts *et al.* 2003, ApJ, 588, 992. Chandra observations.
Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.
Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (18'').
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.
Koo *et al.* 2007, ApJ, 657, 308. IR observations.
Dean *et al.* 2008, MNRAS, 384, L29. Integral observations of pulsar and nebula.

G11.4–0.1

RA: 18^h10^m47^s
Dec: −19°05'

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 8
Type: S?

Radio: Incomplete shell, possibly with central core.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=9.4$ Jy) contaminated by sidelobes of a nearby source, and Parkes 64-m at 5 GHz (4' : $S=2.8$ Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6.8' : $S=2.0 \pm 0.4$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.2' \times 4.1' : $S=18$ Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (44'' \times 63'' : $S=5.1 \pm 0.6$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

G11.8–0.2

RA: 18^h12^m25^s
Dec: −18°44'

1-GHz flux/Jy: 0.7
Spectral index: 0.3

Size/arcmin: 4
Type: S

Has been called G11.89–0.21.

Radio: Shell.

X-ray: Detected.

References:

- Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42' : $S=0.9 \pm 0.1$ Jy), plus other observations.

G12.0–0.1

RA: 18^h12^m11^s
Dec: −18°37'

1-GHz flux/Jy: 3.5
Spectral index: 0.7

Size/arcmin: 7?
Type: ?

Radio: Incomplete shell, defined in E only.

X-ray: Detected.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=6.6$ Jy) and Parkes 64-m at 5 GHz (4' : $S=1.1$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.2' \times 4.1'').
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41'' \times 61'' : $S=0.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

G12.2+0.3

RA: 18^h11^m17^s
Dec: −18°10'

1-GHz flux/Jy: 0.8
Spectral index: 0.7

Size/arcmin: 6 \times 5
Type: S

Has been called G12.26+0.30.

Radio: Partial shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42' : $S=1.5 \pm 0.1$ Jy), plus other observations.

G12.5+0.2

RA: 18^h12^m14^s
Dec: −17°55'

1-GHz flux/Jy: 0.6
Spectral index: 0.4

Size/arcmin: 6 × 5
Type: C?

Has been called G12.58+0.22.

Radio: Diffuse, central brightened.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42" : $S = 0.8 \pm 0.1$ Jy), plus other observations.

G12.7−0.0

RA: 18^h13^m19^s
Dec: −17°54'

1-GHz flux/Jy: 0.8
Spectral index: 0.8

Size/arcmin: 6
Type: S

Has been called G12.72−0.00.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42" : $S = 2.0 \pm 0.1$ Jy), plus other observations.

G12.8−0.0

RA: 18^h13^m37^s
Dec: −17°49'

1-GHz flux/Jy: 0.8
Spectral index: 0.5

Size/arcmin: 3
Type: C?

Has been called G12.82−0.02 and G12.83−0.02.

Radio: Shell.

X-ray: Diffuse, with central source.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.

Brogan *et al.* 2005, ApJ, 629, L105. VLA at 330 MHz (19" × 32"), plus other observations.

Ubertini *et al.* 2005, ApJ, 629, L109. X-ray, γ-ray and other observations.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42" : $S = 1.2 \pm 0.1$ Jy), plus other observations.

Aharonian *et al.* 2006, ApJ, 636, 777. γ-ray detection.

Albert *et al.* 2006, ApJ, 637, L41. γ-ray observations.

Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.

Funk *et al.* 2007, A&A, 470, 249. XMM observations, CO observations of surroundings.

Helfand *et al.* 2007, ApJ, 665, 1297. Chandra observations.

G13.3−1.3

RA: 18^h19^m20^s
Dec: −18°00'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 70 × 40
Type: S?

Radio: Amorphous emission.

X-ray: Elongated emission.

Optical: Filaments in S.

Distance: Absorption indicates 2–4 kpc.

References:

Seward *et al.* 1995, ApJ, 449, 681. ROSAT detection, optical studies and observations of CO.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G13.5+0.2

RA: 18^h14^m14^s
Dec: -17°12'

1-GHz flux/Jy: 3.5?
Spectral index: 1.0?

Size/arcmin: 5 × 4
Type: S

Has been called G13.46+0.16.

Radio: Elongated, incomplete shell.

References:

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ($S=0.65 \pm 0.05$ Jy) and 1.4 GHz ($15'' : S=2.67 \pm 0.5$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G14.1-0.1

RA: 18^h15^m52^s
Dec: -16°34'

1-GHz flux/Jy: 0.5
Spectral index: 0.6

Size/arcmin: 6 × 5
Type: S

Has been called G14.18-0.12.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S=0.9 \pm 0.1$ Jy), plus other observations.

G14.3+0.1

RA: 18^h15^m58^s
Dec: -16°27'

1-GHz flux/Jy: 0.6
Spectral index: 0.4

Size/arcmin: 5 × 4
Type: S

Has been called G14.30+0.14.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S=1.2 \pm 0.1$ Jy), plus other observations.

G15.1-1.6

RA: 18^h24^m00^s
Dec: -16°34'

1-GHz flux/Jy: 5.5?
Spectral index: 0.8?

Size/arcmin: 30 × 24
Type: S

Radio: Elongated, incomplete shell.

Optical: Diffuse shell.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
Boumis *et al.* 2008, A&A, 481, 705. Optical detection.

G15.4+0.1

RA: 18^h18^m02^s
Dec: -15°27'

1-GHz flux/Jy: 5.6
Spectral index: 0.6

Size/arcmin: 15 × 14
Type: S

Has been called G15.42+0.18.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S=10.9 \pm 0.3$ Jy), plus other observations.

G15.9+0.2

RA: 18^h18^m52^s
Dec: −15°02'

1-GHz flux/Jy: 5
Spectral index: 0.6?

Size/arcmin: 7 × 5
Type: S?

Radio: Incomplete shell, with bright concentration to the E.

X-ray: Shell, brighter to S and E.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=7.7$ Jy) and Parkes 64-m at 5 GHz (4' : $S=1.9$ Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 : $S=1.1\pm0.2$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz (58'' × 44'').
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (77'' × 61'': $S=11.2\pm1.0$ Jy), 1.4 GHz (23'' × 14'': $S=3.9\pm0.1$ Jy) and 4.9 GHz (13'' × 16'').
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Reynolds *et al.* 2006, ApJ, 652, L45. Chandra observations.

G16.0−0.5

RA: 18^h21^m56^s
Dec: −15°14'

1-GHz flux/Jy: 2.7
Spectral index: 0.6

Size/arcmin: 15 × 10
Type: S

Has been called G16.05–0.57.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=4.9\pm0.2$ Jy), plus other observations.

G16.2−2.7

RA: 18^h29^m40^s
Dec: −16°08'

1-GHz flux/Jy: 2
Spectral index: 0.5

Size/arcmin: 17
Type: S

Radio: Double rim.

References:

- Trushkin 1999, A&A, 352, L103. Review of radio observations.

G16.4−0.5

RA: 18^h22^m38^s
Dec: −14°55'

1-GHz flux/Jy: 4.6
Spectral index: 0.7

Size/arcmin: 13
Type: S

Has been called G16.41–0.55.

Radio: Partial shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=10.0\pm0.3$ Jy), plus other observations.

G16.7+0.1

RA: 18^h20^m56^s
Dec: −14°20'

1-GHz flux/Jy: 3.0
Spectral index: 0.6

Size/arcmin: 4
Type: C

Has been called G16.73+0.08.

Radio: Asymmetric shell with flat-spectrum core.

X-ray: Non-thermal core.

References:

- Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz (5 : $S=0.95$ Jy) and 1.4 GHz (15'' : $S=2.43$ Jy) and Ooty at 327 MHz ($S=5.13$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.
 Reynoso & Mangum 2000, ApJ, 545, 874. CO observations of surroundings.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Helfand *et al.* 2003, ApJ, 592, 941. XMM observations.
 Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (19'' × 25'').
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G16.8−1.1

RA: 18^h25^m20^s
Dec: −14°46'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 30 × 24?
Type: ?

Has been called G16.85−1.05.

Radio: Overlapping thermal and non-thermal emission, parameters uncertain.

Point sources: Pulsar within boundary of non-thermal emission.

References:

- Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'4), plus other observations.

G17.0−0.0

RA: 18^h21^m57^s
Dec: −14°08'

1-GHz flux/Jy: 0.5
Spectral index: 0.5

Size/arcmin: 5
Type: S

Has been called G17.02−0.04.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S=0.7 \pm 0.1$ Jy), plus other observations.

G17.4−2.3

RA: 18^h30^m55^s
Dec: −14°52'

1-GHz flux/Jy: 4.8?
Spectral index: 0.8?

Size/arcmin: 24?
Type: S

Radio: Incomplete, poorly defined shell.

Optical: Filaments to SE, and diffuse emission.

References:

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2002, A&A, 385, 1042. Optical observations.

G17.4–0.1

RA: 18^h23^m08^s
Dec: −13°46'

1-GHz flux/Jy: 0.4
Spectral index: 0.7

Size/arcmin: 6
Type: S

Has been called G17.48–0.12.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 0.9 \pm 0.1$ Jy), plus other observations.

G17.8–2.6

RA: 18^h32^m50^s
Dec: −14°39'

1-GHz flux/Jy: 4.0?
Spectral index: 0.3?

Size/arcmin: 24
Type: S

Radio: Well defined shell.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4').
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G18.1–0.1

RA: 18^h24^m34^s
Dec: −13°11'

1-GHz flux/Jy: 4.6
Spectral index: 0.5

Size/arcmin: 8
Type: S

Has been called G18.1–0.2 and G18.16–0.16.

Radio: Shell.

X-ray: Detected.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (8').
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 7.6 \pm 0.1$ Jy), plus other observations.

G18.6–0.2

RA: 18^h25^m55^s
Dec: −12°50'

1-GHz flux/Jy: 1.4
Spectral index: 0.4

Size/arcmin: 6
Type: S

Has been called G18.62–0.28.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 1.9 \pm 0.1$ Jy), plus other observations.

G18.8+0.3

Kes 67

RA: 18^h23^m58^s
Dec: -12°23'

1-GHz flux/Jy: 33
Spectral index: 0.4

Size/arcmin: 17×11
Type: S

Has been called G18.9+0.3.

Radio: Incomplete shell, in complex region near the HII region W39.

Distance: Association with molecular cloud and HI absorption suggests 12 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' : $S=17\pm7$ Jy).
Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3' : $S=38$ Jy).
Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
Dicke & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
Milne *et al.* 1989, PASA, 8, 187. Parkes 64-m at 8.4 GHz (3'0 : $S=12.9\pm1.0$ Jy), including polarisation.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (2'9 × 3'5 : $S=55$ Jy).
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (75'' × 55'' : $S=29.9\pm0.3$ Jy).
Dubner *et al.* 1999, AJ, 118, 930. Parkes 64-m at 1.6 GHz (15'') for HI, VLA at 1.6 GHz (17'' × 12'') for OH, plus CO observations.
Dubner *et al.* 2004, A&A, 426, 201. CO observations of environment.
Tian *et al.* 2007, A&A, 474, 541. VGPS at 1.4 GHz (1') including HI, plus CO of surroundings.

G18.9–1.1

RA: 18^h29^m50^s
Dec: -12°58'

1-GHz flux/Jy: 37
Spectral index: varies

Size/arcmin: 33
Type: C?

Has been called G18.95–1.1 and G18.94–1.04.

Radio: Non-thermal, diffuse partially limb-brightened, with central ridge.

X-ray: Partial shell.

Distance: Various observations suggest 2 kpc.

References:

- Fürst *et al.* 1985, Nature, 314, 720. Effelsberg 100-m at 4.75 GHz (2'4 : $S=23.8$ Jy), plus other flux densities.
Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7'2 × 8' : $S=82\pm15$ Jy), plus review of flux densities.
Barnes & Turtle 1988, in SNRISM, p347. Molonglo at 408 MHz (2'9 × 3'1 : $S=58\pm9$ Jy) and Parkes 64-m at 5 GHz (4'4 × 4'1 : $S=23\pm6$ Jy).
Patnaik *et al.* 1988, Nature, 332, 136. Ooty at 327 MHz (0'6 × 1'6).
Fürst *et al.* 1989, A&A, 209, 361. Effelsberg 100-m at 4.75 GHz (2'45 : $S=23.8$ Jy) and VLA at 1.5 and 4.9 GHz (19'' × 14''), and Effelsberg 100-m at 1.4 GHz (9') for HI.
Aschenbach *et al.* 1991, A&A, 246, L32. ROSAT observations.
Fürst *et al.* 1997, A&A, 319, 655. ROSAT observations, and Effelsberg 100-m at 10.6 GHz (1'1).
Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.
Harrus *et al.* 2004, ApJ, 603, 152. ROSAT and ASCA observations.

G19.1+0.2

RA: 18^h24^m56^s
Dec: -12°07'

1-GHz flux/Jy: 10
Spectral index: 0.5

Size/arcmin: 27
Type: S

Has been called G19.15+0.27.

Radio: Partial shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S=17.4\pm0.4$ Jy), plus other observations.

G20.0–0.2

RA: 18^h28^m07^s
Dec: −11°35'

1-GHz flux/Jy: 10
Spectral index: 0.0

Size/arcmin: 10
Type: F

Radio: Faint, filled-centre, polarised.

Point sources: OH source 20.1–0.1 is nearby.

References:

Becker & Helfand 1985, ApJ, 297, L25. VLA at 1.4 and 5 GHz (12'').
Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7'2 × 8': $S=8.5 \pm 2$ Jy), plus review of flux densities.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

G20.4+0.1

RA: 18^h27^m51^s
Dec: −11°00'

1-GHz flux/Jy: 3.1
Spectral index: 0.4

Size/arcmin: 8
Type: S

Has been called G20.47+0.16.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=4.2 \pm 0.1$ Jy), plus other observations.

G21.0–0.4

RA: 18^h31^m12^s
Dec: −10°47'

1-GHz flux/Jy: 1.1
Spectral index: 0.6

Size/arcmin: 9 × 7
Type: S

Has been called G21.04–0.47.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=2.3 \pm 0.2$ Jy), plus other observations.

G21.5–0.9

RA: 18^h33^m33^s
Dec: −10°35'

1-GHz flux/Jy: 6?
Spectral index: 0.0

Size/arcmin: 4
Type: C

Early observations relate to the central core only.

Radio: Filled-centre, with high frequency turnover.

X-ray: Central core, with extended, faint halo.

Point sources: Central pulsar.

Distance: HI absorption indicates 4.6 kpc.

References:

- Wilson & Weiler 1976, A&A, 53, 89. WSRT at 5 GHz ($6'' \times 35''$).
 Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz ($20'' \times 5''$) and 8 GHz ($7'' \times 2''$), plus review of flux densities.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2''.6).
 Becker & Szymkowiak 1981, ApJ, 248, L23. VLA at 5 GHz (8'')., and Einstein observations.
 Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT X-ray spectrum, and VLA HI observations.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz ($26''.5 : S = 5.64 \pm 0.29$ Jy).
 Fürst *et al.* 1988, PASJ, 40, 347. NRO array at 22.3 GHz ($4''.4 \times 7''.3$).
 Salter *et al.* 1989, A&A, 225, 167. Observations at 90.7 (29''.5 : $S = 3.8 \pm 0.4$ Jy) and 141.9 GHz ($S = 2.5 \pm 1.2$ Jy).
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ($S = 3.94 \pm 0.70$ Jy), plus review of flux densities.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3''.3 \times 3''.4 : S = 9$ Jy).
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Slane *et al.* 2000, ApJ, 533, L29. Chandra observations identifying X-ray halo.
 Warwick *et al.* 2001, A&A, 365, L248. XMM observations of X-ray halo.
 Bock *et al.* 2001, ApJ, 561, L203. BIMA at 94 GHz ($8''.6 \times 4''.6$).
 Safi-Harb *et al.* 2001, ApJ, 561, 308. Chandra and other X-ray observations.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 22 GHz (8'') and 32 GHz, including polarisation.
 La Palombara & Mereghetti 2002, A&A, 383, 916. XMM upper limit on pulsations.
 Bocchino *et al.* 2005, A&A, 442, 539. XMM and Chandra observations.
 Gupta *et al.* 2005, Current Science, 89, 853. Pulsar discovery.
 Camilo *et al.* 2006, ApJ, 637, 456. Pulsar discovery.
 Bietenholz & Bartel 2008, MNRAS, 386, 1411. VLA at 4.75 GHz ($0''.53 \times 0''.82$) for expansion studies.
 Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1') including HI.

G21.5–0.1

RA: 18^h30^m50^s
Dec: −10°09'

1-GHz flux/Jy: 0.4
Spectral index: 0.5

Size/arcmin: 5
Type: S

Has been called G21.56–0.10.

Radio: Partial shell.

X-ray: Detected.

References:

- Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.5 \pm 0.1$ Jy), plus other observations.

G21.8–0.6		Kes 69
RA: 18 ^h 32 ^m 45 ^s	1-GHz flux/Jy: 69	Size/arcmin: 20
Dec: −10°08'	Spectral index: 0.5	Type: S
Radio: Incomplete shell.		
X-ray: Detected.		
Distance: HI absorption indicates 5.5 to 7.4 kpc.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Wilson 1972, A&A, 19, 354. H ₂ CO absorption.		
Kundu <i>et al.</i> 1974, AJ, 79, 132. NRAO 140-ft at 5 GHz (6') and 10 GHz (3').		
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft et 2.7 GHz (5'): $S = 42.3 \pm 4.6$ Jy.		
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Seward 1990, ApJS, 73, 781. Einstein observations.		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3 × 3'.5 : $S = 132$ Jy).		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.		
Yusef-Zadeh <i>et al.</i> 2003, ApJ, 585, 319. X-ray observations.		
Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer observations.		
Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.		
Wood <i>et al.</i> 2008, AJ, 135, 2358. VLA at 4.8 GHz, including polarisation.		
Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1') including HI.		

G22.7–0.2		
RA: 18 ^h 33 ^m 15 ^s	1-GHz flux/Jy: 33	Size/arcmin: 26
Dec: −09°13'	Spectral index: 0.6	Type: S?
Radio: Non-thermal ring in complex region, overlapping G23.3–0.3.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3 × 3'.4 : $S = 82$ Jy).		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.		
Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer observations.		

G23.3–0.3		W41
RA: 18 ^h 34 ^m 45 ^s	1-GHz flux/Jy: 70	Size/arcmin: 27
Dec: −08°48'	Spectral index: 0.5	Type: S
Radio: Distorted ring, in complex region, overlapping G22.7–0.2.		
X-ray: Possible extended emission, with compact sources.		
Point sources: Pulsar association suggested.		
Distance: HI and CO observations indicate 4.2 kpc.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3 × 3'.4 : $S = 138$ Jy).		
Gaensler & Johnston 1995, MNRAS, 275, L73. Possible pulsar association.		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.		
Aharonian <i>et al.</i> 2005, Science, 307, 1938. γ -ray detection.		
Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer possible detection.		
Aharonian <i>et al.</i> 2006, ApJ, 636, 777. γ -ray observations.		
Albert <i>et al.</i> 2006, ApJ, 643, L53. γ -ray observations.		
Landi <i>et al.</i> 2006, ApJ, 651, 190. X-ray observations.		
Tian <i>et al.</i> 2007, ApJ, 657, L25. VGPS at 1.4 GHz (1') including HI, plus XMM observations.		
Leahy & Tian 2008, AJ, 135, 167. VGPS at 1.4 GHz (1') including HI, plus CO observations.		

G23.6+0.3

RA: 18^h33^m03^s
Dec: −08°13'

1-GHz flux/Jy: 8?
Spectral index: 0.3

Size/arcmin: 10?
Type: ?

Radio: Not well resolved, in complex region.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G24.7−0.6

RA: 18^h38^m43^s
Dec: −07°32'

1-GHz flux/Jy: 8
Spectral index: 0.5

Size/arcmin: 15?
Type: S?

Radio: Incomplete shell, defined in SW.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=12.3$ Jy) and Parkes 64-m at 5 GHz (4': $S=3.6$ Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8': $S=2.2 \pm 0.5$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (44'' \times 56'': $S=1.9$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

G24.7+0.6

RA: 18^h34^m10^s
Dec: −07°05'

1-GHz flux/Jy: 20?
Spectral index: 0.2?

Size/arcmin: 30 \times 15
Type: C?

Radio: Filled-centre, with faint shell, and a compact HII region to the S.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6).
 Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4.3': $S=19 \pm 3$ Jy) and 4.75 GHz (2'.4': $S=17 \pm 4$ Jy) and NRO 45-m at 10.2 GHz (2'.7': $S=15 \pm 3$ Jy).
 Becker & Helfand 1987, ApJ, 316, 660. VLA at 1.4 GHz (12''), and X-ray upper limit.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

G27.4+0.0

4C–04.71

RA: 18^h41^m19^s
Dec: −04°56'

1-GHz flux/Jy: 6
Spectral index: 0.68

Size/arcmin: 4
Type: S

Early references refer to G27.3–0.1 (Kes 73), a supposed larger remnant.

Radio: Incomplete shell.

X-ray: Diffuse emission, with central low period pulsar.

Point sources: Central AXP.

Distance: HI absorption suggests 7.5 to 9.8 kpc.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5').
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8/4) and 5 GHz (4/4).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6/8 : $S = 2.0 \pm 0.5$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2/6).
 Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3' : $S = 4.4$ Jy).
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz ($60'' \times 45''$: $S = 3.5$ Jy). Revise $S_{408\text{ MHz}} = 10.4$ Jy, and $S_{5\text{ GHz}} = 1.9 \pm 0.2$ Jy.
 Kriss *et al.* 1985, ApJ, 288, 703. Einstein observations, plus VLA at 1.4 and 5 GHz (12'').
 Sanbonmatsu & Helfand 1992, AJ, 104, 2189. VLA at 1.4 GHz for HI absorption.
 Helfand *et al.* 1994, ApJ, 434, 627. ROSAT observations, particularly of central source.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including nearby masers.
 Vasish & Gotthelf 1997, ApJ, 486, L129. ASCA detection of pulsar.
 Gotthelf & Vasish 1997, ApJ, 486, L133. ASCA observations.
 Gotthelf *et al.* 1999, ApJ, 522, L49. X-ray timing observations of pulsar.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Mereghetti *et al.* 2001, MNRAS, 321, 143. Search for optical/IR counterpart to pulsar.
 Kuiper *et al.* 2004, ApJ, 613, 1173. X-ray observations of pulsar.
 Tian & Leany 2008, ApJ, 677, 292. VGPS at 1.4 GHz (1') including HI.

G27.8+0.6

RA: 18^h39^m50^s
Dec: −04°24'

1-GHz flux/Jy: 30
Spectral index: varies

Size/arcmin: 50 × 30
Type: F

Radio: Filled-centre, with spectral turnover.

References:

- Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4/3 : $S = 23 \pm 2$ Jy), and 4.75 GHz (2/4 : $S = 18 \pm 2$ Jy) and NRO 45-m at 10.2 GHz (smoothed to 4/3 : $S = 8.5 \pm 2$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.

G28.6–0.1

RA: 18^h43^m55^s
Dec: −03°53'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 13 × 9
Type: S

Radio: Poorly defined regions of non-thermal emission.

X-ray: Diffuse shell, with thermal and non-thermal emission.

References:

- Helfand *et al.* 1989, ApJ, 341, 151. VLA at 1.4 GHz (15'') and 5 GHz (15'').
 Bamba *et al.* 2001, PASJ, 53, L21. ASCA observations.
 Ueno *et al.* 2003, ApJ, 588, 338. Chandra observations.

G28.8+1.5

RA: 18^h39^m00^s
Dec: −02°55'

1-GHz flux/Jy: ?
Spectral index: 0.4?

Size/arcmin: 100?
Type: S?

Radio: Part of rim detected.

X-ray: Diffuse, Centrally brightened.

References:

- Schwentker 1994, A&A, 286, L47. ROSAT observations.
Song *et al.* 2000, PASJ, 52, 181. ASCA observations.
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

G29.6+0.1

RA: 18^h44^m52^s
Dec: −02°57'

1-GHz flux/Jy: 1.5?
Spectral index: 0.5?

Size/arcmin: 5
Type: S

Radio: Diffuse shell.

Point sources: AXp associated.

References:

- Gaensler *et al.* 1999, ApJ, 526, L37. VLA at 5 GHz (13'') and 8 GHz (8'')
Vasisht *et al.* 2000, ApJ, 542, L49. X-ray observations of AXp.

G29.7−0.3

RA: 18^h46^m25^s
Dec: −02°59'

1-GHz flux/Jy: 10
Spectral index: 0.7

Size/arcmin: 3
Type: C

Kes 75

Has erroneously been called G29.6+0.1.

Radio: Shell with flatter spectrum emission from centre.

X-ray: Thermal shell and non-thermal core, and central pulsar.

Point sources: X-ray pulsar.

Distance: HI absorption indicates possibly < 7.5 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (8'4 : $S = 5 \pm 20$ Jy).
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20'' \times 7'') and 8 GHz (25'' \times 8''), plus review of flux densities.
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Becker *et al.* 1983, ApJ, 268, L93. VLA at 1.4 GHz (3''), plus Einstein observations.
Becker & Helfand 1984, ApJ, 283, 154. VLA at 5 GHz (2''.6), plus HI.
Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26''.5 : $S = 1.02 \pm 0.07$ Jy).
Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3''.4 \times 3''.7 : $S = 27.4$ Jy).
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Blanton & Helfand 1996, ApJ, 470, 961. ASCA observations.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Gothelf *et al.* 2000, ApJ, 542, L37. X-ray pulsar detection.
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Mereghetti *et al.* 2002, ApJ, 574, 873. BeppoSAX observations of pulsar.
Helfand *et al.* 2003, ApJ, 582, 783. Chandra observations.
Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (10'' \times 13'').
Morton *et al.* 2007, ApJ, 667, 219. Spitzer and Chandra observations.
McBride *et al.* 2008, A&A, 477, 249. γ -ray and Chandra observations.
Leahy & Tian 2008, A&A, 480, L25. VGPS at 1.4 GHz (1') including HI.
Kumar & Safi-Harb 2008, ApJ, 678, L43. Chandra observations.
Ng *et al.* 2008, ApJ, 686, 508. Chandra observations.
Gavriil *et al.* 2008, Science, 319, 1802. X-ray observations of pulsar.

G30.7–2.0

RA: 18^h54^m25^s
Dec: −02°54'

1-GHz flux/Jy: 0.5?
Spectral index: 0.7?

Size/arcmin: 16
Type: ?

Radio: Poorly defined.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G30.7+1.0

RA: 18^h44^m00^s
Dec: −01°32'

1-GHz flux/Jy: 6
Spectral index: 0.4

Size/arcmin: 24 × 18
Type: S?

Radio: Non-thermal, highly polarised part shell?

Point sources: Compact source near centre.

References:

Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'.4 : $S = 3.4 \pm 0.4$ Jy), plus other flux densities.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.6 × 3'.9 : $S = 8.6$ Jy).
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

G31.5–0.6

RA: 18^h51^m10^s
Dec: −01°31'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 18?
Type: S?

Has been called G31.55–0.65.

Radio: Distorted shell? near HII region.

Optical: Diffuse, incomplete shell.

References:

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4), plus other flux densities.
 Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

G31.9+0.0		3C391
RA: 18 ^h 49 ^m 25 ^s	1-GHz flux/Jy: 24	Size/arcmin: 7 × 5
Dec: −00°55'	Spectral index: 0.49	Type: S
Radio: Shell, brightest in NW, with low frequency turnover.		
X-ray: Diffuse with central core.		
Distance: HI absorption is seen to the tangent point (8.5 kpc).		
References:		
Radhakrishnan <i>et al.</i> 1972, ApJS, 24, 49. HI absorption.		
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').		
Green <i>et al.</i> 1975, A&A, 44, 187. Effelsberg 100-m at 15 GHz (58'': $S > 1.6$).		
Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20'' × 23'') and 8 GHz (24'' × 9''), plus review of flux densities.		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').		
Goss <i>et al.</i> 1979, A&A, 78, 75. FIRST at 1.4 GHz (54'' × 66'': $S = 20 \pm 2$ Jy) and Effelsberg 100-m at 10.7 GHz (77'': $S = 7.5 \pm 0.8$ Jy).		
Wang & Seward 1984, ApJ, 279, 705. Einstein observations.		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.4' × 3.7': $S = 44.8$ Jy).		
Reynolds & Moffett 1993, AJ, 105, 2226. VLA at 1.4 GHz (6''), including possible associated CO.		
Moffett & Reynolds 1994, ApJ, 425, 668. VLA at 330 MHz (smoothed to 30'': $S = 38.5 \pm 0.5$ Jy) 1.46 GHz (6''.7) and 4.85 GHz (6''.2 × 6''.4), including spectral index and polarisation studies.		
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant, including masers.		
Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.		
Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.		
Rho & Petre 1996, ApJ, 467, 698. ROSAT observations.		
Wilner <i>et al.</i> 1998, AJ, 115, 247. CO observations of surroundings.		
Reach & Rho 1998, ApJ, 507, L93. ISO observations.		
Reach & Rho 1999, ApJ, 511, 836. CO, HCO ⁺ and CS observations of surroundings.		
Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.		
Sugizaki <i>et al.</i> 2001, ApJS, 134, 77. ASCA survey observations.		
Chen & Slane 2001, ApJ, 563, 202. ASCA observations.		
Reach <i>et al.</i> 2002, ApJ, 564, 302. Observations of shocked molecular species.		
Chen <i>et al.</i> 2004, ApJ, 616, 885. Chandra observations.		
Brogan <i>et al.</i> 2005, AJ, 130, 148. VLA at 74 MHz, 330 MHz and 1.5 GHz (70'').		
Kawasaki <i>et al.</i> 2005, ApJ, 631, 935. ASCA observations.		
Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer observations.		
Neufeld <i>et al.</i> 2007, ApJ, 664, 890. Spitzer IR observations.		
Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.		

G32.0−4.9		3C396.1
RA: 19 ^h 06 ^m 00 ^s	1-GHz flux/Jy: 22?	Size/arcmin: 60?
Dec: −03°00'	Spectral index: 0.5?	Type: S?
Radio: Possible large shell?		
References:		
Milne & Hill 1969, AuJPh, 22, 211. Parkes 64-m at 635 MHz (31': $S = 25 \pm 30\%$ Jy), 1410 MHz (15': $S = 19 \pm 15\%$ Jy) and 2650 MHz (8''.4: $S = 8.6 \pm 30\%$ Jy). Fluxes if size is 60', plus review of flux densities.		
Caswell 1970, AuJPh, 23, 105. Revision of low frequency flux densities.		
Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 111 MHz ($S = 105 \pm 30$ Jy).		

G32.1−0.9		
RA: 18 ^h 53 ^m 10 ^s	1-GHz flux/Jy: ?	Size/arcmin: 40?
Dec: −01°08'	Spectral index: ?	Type: C?
Radio: Possible faint shell, not well defined.		
X-ray: Diffuse, with clumps.		
References:		
Folgheraiter <i>et al.</i> 1997, MNRAS, 292, 365. ROSAT and ASCA observations.		

G32.4+0.1

RA: 18^h50^m05^s
Dec: -00°25'

1-GHz flux/Jy: 0.25?
Spectral index: ?

Size/arcmin: 6
Type: S

Has been called G32.45+0.1.

Radio: Shell.

X-ray: Shell.

Distance: X-ray absorption suggests 17 kpc.

References:

Yamaguchi *et al.* 2004, PASJ, 56, 1059. XMM and other observations.
Ueno *et al.* 2005, in XRRC, E4.18. XMM observations.

G32.8-0.1

RA: 18^h51^m25^s
Dec: -00°08'

1-GHz flux/Jy: 11?
Spectral index: 0.2?

Kes 78
Size/arcmin: 17
Type: S?

Part has been called G33.1-0.1.

Radio: Elongated shell?

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5': $S=7.2 \pm 0.5$ Jy).
Dickey & DeNoyer 1975, AJ, 80, 437. Arecibo at 430 MHz ($S=19.0 \pm 15.5$ Jy).
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3': $S=12.8$ Jy) and Parkes 64-m at 5 GHz (4': $S=7.7$ Jy).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.6 \times 3'.8 : $S=31.3$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.

G33.2-0.6

RA: 18^h53^m50^s
Dec: -00°02'

1-GHz flux/Jy: 3.5
Spectral index: varies

Size/arcmin: 18
Type: S

Radio: Incomplete shell.

References:

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
Reich 1982, A&A, 106, 314. Effelsberg 100-m at 2.7 GHz (4'.4 : $S=2.6 \pm 0.3$ Jy) and 4.75 GHz (2'.5 : $S=1.75 \pm 0.2$ Jy).
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (68'' \times 52'' : $S=2.7 \pm 0.3$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G33.6+0.1

RA: 18^h52^m48^s
Dec: +00°41'

1-GHz flux/Jy: 22
Spectral index: 0.5

Kes 79, 4C00.70, HC13

Size/arcmin: 10
Type: S

Has been called G33.7+0.0.

Radio: Shell, with bright central region, in complex region.

X-ray: Multiple shells and filaments.

Point sources: Central X-ray pulsar.

Distance: HI absorption gives about 7.8 kpc.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S = 35.5$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 7.8$ Jy).
 Caswell *et al.* 1975, A&A, 45, 239. HI absorption.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430} MHz = 69 ± 33 Jy.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3' : $S = 6.8 \pm 1.5$ Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8' : $S = 11.4 \pm 1.1$ Jy).
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1'), plus observations of the nearby point source.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source, possibly extragalactic.
 Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source, plus Einstein observations.
 Green 1989, MNRAS, 238, 737. OH absorption.
 Frail & Clifton 1989, ApJ, 336, 854. VLA at 1.4 GHz (1' \times 2'.9'), including HI absorption.
 Velusamy *et al.* 1991, AJ, 102, 676. VLA at 327 MHz (1'), 1.5 (7'' \times 14'') and 5 GHz (7''), including spectral comparison.
 Green & Dewdney 1992, MNRAS, 254, 686. Observations of adjacent molecular material.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.6 \times 3'.8' : $S = 34.8$ Jy).
 Seward & Velusamy 1995, ApJ, 439, 715. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Tsunemi & Enoguchi 2002, PASJ, 54, 735. ASCA observations.
 Seward *et al.* 2003, ApJ, 584, 414. Chandra observations.
 Stanimirović 2003, ApJ, 592, 953. Arecibo OH absorption.
 Sun *et al.* 2004, ApJ, 605, 742. Chandra observations.
 Gotthelf *et al.* 2005, ApJ, 627, 390. XMM pulsar detection.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Halpern *et al.* 2007, ApJ, 665, 1304. X-ray pulsar timing observations.

G34.7–0.4

RA: 18^h56^m00^s
Dec: +01°22'

1-GHz flux/Jy: 230
Spectral index: 0.37

W44, 3C392

Size/arcmin: 35 × 27
Type: C

Has been called G34.6–0.5.

Radio: Distorted shell, brighter to the E, with pulsar and associated nebula.

Optical: Diffuse emission.

X-ray: Centrally concentrated, thermal spectrum, plus pulsar wind nebula.

Point sources: Pulsar within the boundary of the remnant.

Distance: HI absorption indicates 2.8 kpc.

References:

- Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10.7 GHz ($3' : S = 105 \pm 7$ Jy).
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 177 \pm 10$ Jy).
Caswell *et al.* 1975, A&A, 45, 239. HI absorption.
Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 299$ Jy).
Dicke & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 540 \pm 187$ Jy.
Dicke & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
Smith *et al.* 1985, MNRAS, 217, 99. Einstein observations.
Wolszczan *et al.* 1991, ApJ, 372, L99. Pulsar detection.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3.6 \times 3.8' : S = 469$ Jy).
Jones *et al.* 1993, MNRAS, 265, 631. VLA at 1.4 GHz (15''), plus X-ray spectra.
Rho *et al.* 1994, ApJ, 430, 757. Optical and ROSAT observations.
Koo & Heiles 1995, ApJ, 442, 679. HI of surrounding shell.
Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ -ray emission.
Harrus *et al.* 1996, ApJ, 464, L161. ASCA observations.
Frail *et al.* 1996, ApJ, 464, L165. VLA at 1.5 and 8.4 GHz (8''.9 × 7''.8) of pulsar nebula.
Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.
Harrus *et al.* 1997, ApJ, 488, 781. X-ray observations.
Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.
Giacani *et al.* 1997, AJ, 113, 1379. VLA at 1.4 GHz (15''), plus optical images.
Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.
Cox *et al.* 1999, ApJ, 524, 179. Revision of distance.
Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
Petre *et al.* 2002, ApJ, 579, 404. Chandra observations of pulsar and wind nebula.
Mavromatakis *et al.* 2003, A&A, 405, 591. Optical observations.
Shelton *et al.* 2004, ApJ, 611, 906. Chandra observations of part.
Reach *et al.* 2005, ApJ, 618, 297. Molecular line and near-IR observations.
Hoffman *et al.* 2005, ApJ, 627, 803. Observations of OH masers.
Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer IR observations.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
Castelletti *et al.* 2007, A&A, 471, 537. VLA at 74 MHz ($36'' \times 39'' : S = 634 \pm 70$ Jy) and 324 MHz ($13'' : S = 411 \pm 50$ Jy).

G36.6–0.7

RA: 19^h00^m35^s
Dec: +02°56'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 25?
Type: S?

Radio: polarised arc, possibly part of a larger shell?

References:

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4), plus other flux densities.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3.4 \times 3.7' : S = 6.7$ Jy).
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G36.6+2.6

RA: 18^h48^m49^s
Dec: +04°26'

1-GHz flux/Jy: 0.7?
Spectral index: 0.5?

Size/arcmin: 17 × 13?
Type: S

Radio: Poorly resolved shell.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

G39.2–0.3

RA: 19^h04^m08^s
Dec: +05°28'

1-GHz flux/Jy: 18
Spectral index: 0.6

3C396, HC24, NRAO 593

Size/arcmin: 8 × 6
Type: C

Radio: Shell, brighter to W, with faint ‘tail’ to E.

X-ray: Diffuse, brighter to W, with central core.

Point sources: Central X-ray source.

Distance: HI absorption suggests > 7.7 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3').
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430 \text{ MHz}} = 54 \pm 38 \text{ Jy}$.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3': $S = 4.1 \pm 1.0 \text{ Jy}$).
 Caswell *et al.* 1975, A&A, 45, 239. HI absorption.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz ($65'' \times 48''$).
 Becker & Helfand 1987, AJ, 94, 1629. VLA at 1.4 GHz ($12'' : S = 14 \text{ Jy}$) and 5 GHz, plus Einstein observations.
 Patnaik *et al.* 1990, A&A, 232, 467. VLA at 1.5 GHz ($25''$) and 1.4 GHz ($7''.8 \times 7''.5$) and 5 GHz ($25''$) including polarisation, plus Ooty at 327 MHz ($100'' \times 31''$), including review of flux densities.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3''.5 \times 3''.6 : S = 42.5 \text{ Jy}$).
 Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz for spectral index studies.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Harris & Slane 1999, ApJ, 516, 811. ASCA observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Olbert *et al.* 2003, ApJ, 592, L45. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Scaife *et al.* 2007, MNRAS, 377, L69. 33 GHz observations.

G39.7–2.0

RA: 19^h12^m20^s
Dec: +04°55'

1-GHz flux/Jy: 85?
Spectral index: 0.7?

W50, SS433
Size/arcmin: 120 × 60
Type: ?

Eastern part has been called G40.0–3.1. Is this a SNR?

Radio: Elongated shell, containing SS433, adjacent to the HII region S74.

Optical: Faint filaments at the edge of the radio emission.

X-ray: Emission from SS433 and two lobes.

Point sources: SS433 is the compact source in the centre of the W50.

Distance: HI absorption indicates 6.0 ± 0.5 kpc.

References:

- van den Bergh 1980, ApJ, 236, L23. Optical in H α and [SII].
Zealey *et al.* 1980, MNRAS, 192, 731. Optical spectra.
van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source (not SS433).
Sequaist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
Watson *et al.* 1983, ApJ, 273, 688. X-ray observations.
Downes *et al.* 1986, MNRAS, 218, 393. Effelsberg 100-m at 4.75 GHz ($2\farcm4 : S = 34 \pm 4$ Jy), plus previous 1.7 and 2.7 GHz data.
Romney *et al.* 1987, ApJ, 321, 822. VLBI of SS433, including distance.
Elston & Baum 1987, AJ, 94, 1633. Mosaic with VLA at 1.4 GHz ($30''$) of fine structure only.
Kawai *et al.* 1989, PASJ, 41, 491. X-ray observations of SS433.
Band 1989, ApJ, 336, 937. Einstein and EXOSAT observations.
Yamauchi *et al.* 1994, PASJ, 46, L109. X-ray spectral observations.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Brinkmann *et al.* 1996, A&A, 312, 306. ROSAT observations.
Kotami *et al.* 1996, PASJ, 48, 619. X-ray line emission from SS433.
Safi-Harb & Ögelman 1997, ApJ, 483, 868. ROSAT observations.
Dubner *et al.* 1998, AJ, 116, 1842. VLA at 328 MHz ($64'' \times 60'' : S = 160 \pm 20$ Jy), and 1.4 GHz ($56'' \times 54''$), plus NRAO 140-ft at 1.4 GHz ($21'$) for HI observations.
Safi-Harb & Petre 1999, ApJ, 512, 784. X-ray observations.
Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
Brinkmann *et al.* 2007, A&A, 463, 611. XMM observations of E lobe.
Boumis *et al.* 2007, MNRAS, 381, 308. Optical observations.
Lockman *et al.* 2007, MNRAS, 381, 881. HI observations.

G40.5–0.5

RA: 19^h07^m10^s
Dec: +06°31'

1-GHz flux/Jy: 11
Spectral index: 0.5

Size/arcmin: 22
Type: S

Radio: Shell, brightest to the NE.

References:

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2\farcm6$).
Downes *et al.* 1980, A&A, 92, 47. Effelsberg 100-m at 1.7 GHz ($7\farcm6 : S = 9.3 \pm 1.3$ Jy), and 2.7 GHz ($4\farcm4 : S = 7.2 \pm 0.5$ Jy), plus review of flux densities.
Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
Yang *et al.* 2006, ChJAA, 6, 210. CO observations of surroundings.
Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.

G41.1–0.3		3C397
RA: 19 ^h 07 ^m 34 ^s	1-GHz flux/Jy: 22	Size/arcmin: 4.5 × 2.5
Dec: +07°08'	Spectral index: 0.48	Type: S

Radio: 3C397 is two sources: the E is the SNR, the W is a HII region.

X-ray: Brighter to the E and W, with central component.

Distance: Possible limit of > 7.5 kpc for non-thermal component from HI absorption.

References:

- Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 10 GHz (3') and 5 GHz (6').
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5': $S=21.3 \pm 1.2$ Jy for both components).
 Caswell *et al.* 1975, A&A, 45, 239. HI absorption.
 Dickey & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 82 \pm 51$ Jy, also Algonquin 46-m at 10.6 GHz (3': $S=12 \pm 2$ Jy), and Haystack 36-m at 15.5 GHz (2': $S=8.5 \pm 3.0$ Jy).
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3': $S=29.8$ Jy) and Parkes 64-m at 5 GHz (4': $S=8.7$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2').
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (58'' × 52'').
 Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz (8''), plus Einstein observations.
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30'': $S=1.10 \pm 0.19$ Jy).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.5 × 3'.6: $S=46.3$ Jy).
 Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz, for spectral index studies.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Chen *et al.* 1999, ApJ, 520, 737. ASCA and ROSAT observations.
 Dyer & Reynolds 1999, ApJ, 526, 365. VLA at 1.5 GHz (6'.9 × 6'.6) and 4.8 GHz (46''.4 × 5''.6), including polarisation and comparison with ROSAT image.
 Safi-Harb *et al.* 2000, ApJ, 545, 922. ROSAT, ASCA and other X-ray observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Safi-Harb *et al.* 2005, ApJ, 618, 321. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

G42.8+0.6

RA: 19 ^h 07 ^m 20 ^s	1-GHz flux/Jy: 3?	Size/arcmin: 24
Dec: +09°05'	Spectral index: 0.5?	Type: S

Has been called G42.8+0.65.

Radio: Faint shell.

Point sources: Near soft gamma repeater, and young pulsar.

References:

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4: $S=1.5 \pm 0.2$ Jy), plus other flux densities.
 Vasisht *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3'.2 × 3'.4).
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.
 Lorimer & Xilouris 2000, ApJ, 545, 385. Pulsar detection.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50''), and other observations of the region.

G43.3–0.2

W49B

RA: 19^h11^m08^s
Dec: +09°06'

1-GHz flux/Jy: 38
Spectral index: 0.48

Size/arcmin: 4 × 3
Type: S

Radio: Shell, brightest to the SE and W, near the HII region W49A.

X-ray: Centrally brightened, elongated E–W.

Distance: HI absorption indicates 10 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 2.7 GHz (4').
 Downes & Wilson 1974, A&A, 34, 133. Effelsberg 100-m at 10.7 GHz (1.3').
 Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz (58'': $S=9.0 \pm 0.7$ Jy).
 Lockhart & Goss 1978, A&A, 67, 355. HI absorption.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Pye *et al.* 1984, MNRAS, 207, 649. Einstein observations, and VLA at 1.4 and 5 GHz (both 12'').
 Smith *et al.* 1985, ApJ, 296, 469. EXOSAT spectrum.
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30'': $S=6.90 \pm 0.38$ Jy).
 Moffett & Reynolds 1994, ApJ, 437, 705. VLA at 330 MHz (6''.7 × 7''.7: $S=64.4$ Jy), 1.48 GHz (4''.8 × 5''.2: $S=31.8$ Jy) and 4.85 GHz (4''.0 × 4''.1), including polarisation.
 Fujimoto *et al.* 1995, PASJ, 47, L31. ASCA observations.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Hwang *et al.* 2000, ApJ, 532, 970. ROSAT image and ASCA spectroscopy.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Brogan & Troland 2001, ApJ, 550, 799. VLA at 1.4 GHz (27'' × 24'' and 5'') for HI Zeeman splitting.
 Lacey *et al.* 2001, ApJ, 559, 954. VLA at 74 MHz (26'' × 23'': $S=55.6$ Jy) and 326 MHz (6''.6 × 6''.2: $S=56.0$ Jy).
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50''), and other observations of the region.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Miceli *et al.* 2006, A&A, 453, 567. XMM observations.
 Keohane *et al.* 2007, ApJ, 654, 938. IR and Chandra observations.

G43.9+1.6

RA: 19^h05^m50^s
Dec: +10°30'

1-GHz flux/Jy: 8.6?
Spectral index: 0.2?

Size/arcmin: 60?
Type: S?

Radio: Large, poorly defined faint shell.

Point sources: Soft gamma repeater nearby.

References:

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3').
 Vasisht *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3'.2 × 3'.4').
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50''), and other observations of the region.

G45.7–0.4

RA: 19^h16^m25^s
Dec: +11°09'

1-GHz flux/Jy: 4.2?
Spectral index: 0.4?

Size/arcmin: 22
Type: S

Radio: Shell, brightest to the SE, poorly defined to NW.

References:

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4: $S=2.6 \pm 0.3$ Jy), plus other flux densities.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G46.8–0.3

(HC30)

RA: 19^h18^m10^s
Dec: +12°09'

1-GHz flux/Jy: 14
Spectral index: 0.5

Size/arcmin: 17 × 13
Type: S

Has been called G46.6–0.2.

Radio: Shell, two bright arcs to NNW and SSE.

Distance: HI absorption suggests 6.8–8.8 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' : $S=9.8 \pm 0.9$ Jy), and VRO 37-m at 1.7 GHz ($S=14.5 \pm 5.5$ Jy).
Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S=20.3$ Jy) and Parkes 64-m at 5 GHz (4' : $S=7.1$ Jy).
Dicke & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}}=46 \pm 21$ Jy.
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'8" : $S=5.9 \pm 0.6$ Jy).
Sato 1979, ApL, 20, 43. HI observations.
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (58" × 53" : $S=13.3 \pm 0.1$ Jy).

G49.2–0.7

(W51)

RA: 19^h23^m50^s
Dec: +14°06'

1-GHz flux/Jy: 160?
Spectral index: 0.3?

Size/arcmin: 30
Type: S?

Radio: In complex region, parameters uncertain.

X-ray: Elongated east–west.

Optical: Some diffuse emission possibly associated.

Distance: Association with CO gives 6 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Parkes 64-m at 5 GHz (4').
Sato 1973, PASJ, 25, 135. HI absorption.
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5'), $S=51.5 \pm 3.2$ Jy, for the non-thermal component, but probably confused.
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
Seward 1990, ApJS, 73, 781. Einstein observations.
Copetti & Schmidt 1991, MNRAS, 250, 127. CLFST at 151 MHz.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'1 × 3'5).
Subrahmanyam & Goss 1995, MNRAS, 275, 755. VLA at 330 MHz (1'1).
Koo *et al.* 1995, ApJ, 447, 211. ROSAT observations.
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
Green *et al.* 1997, AJ, 114, 2058. OH masers.
Koo & Moon 1997, ApJ, 475, 194. Arecibo (3') and VLA (40" × 42") at 1.4 GHz for HI.
Koo & Moon 1997, ApJ, 485, 263. NRAO 12-m CO and HCO⁺ observations.
Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.
Koo *et al.* 2002, AJ, 123, 1629. ASCA observations.
Koo *et al.* 2005, ApJ, 633, 946. Chandra observations.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.

G53.6–2.2

RA: 19^h38^m50^s
Dec: +17°14'

1-GHz flux/Jy: 8
Spectral index: 0.75

3C400.2, NRAO 611
Size/arcmin: 33 × 28
Type: S

Has been called G53.7–2.2.

Radio: Ring of emission, with extension to NW.

Optical: Filaments and diffuse emission.

X-ray: Centrally brightened, offset to NW.

Distance: Association with HI gives 2.8 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 5.3 \pm 0.6$ Jy).
van den Bergh 1978, ApJS, 38, 119. Optical observations.
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 4.8 \pm 0.3$ Jy).
Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 11.7$ Jy).
Dickey & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 20 \pm 10$ Jy, $S_{318\text{ MHz}} = 20 \pm 3.6$ Jy.
Goss *et al.* 1975, A&A, 43, 459. WSRT at 610 MHz ($1' \times 3' : S = 13.2 \pm 1.6$ Jy).
Sabaddin & d'Odorico 1976, A&A, 49, 119. Optical spectra.
Rosado 1983, RMxAA, 8, 59. Optical spectra.
Blair & Long 1988, PASP, 100, 461. Optical imaging and spectroscopy.
Long *et al.* 1991, ApJ, 373, 567. Einstein and optical observations.
Winkler *et al.* 1993, ApJ, 405, 608. Optical imaging.
Dubner *et al.* 1994, AJ, 108, 207. VLA at 327 MHz (59'') and 1.49 GHz (52''), plus X-rays.
Saken *et al.* 1995, ApJ, 443, 231. ROSAT observations.
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
Giacani *et al.* 1998, A&AS, 133, 61. DRAO at 1.4 GHz for HI studies, including distance.
Yoshita *et al.* 2001, PASJ, 53, 93. ASCA observations, and spectral comparison with ROSAT.
Ambrocio-Cruz *et al.* 2006, RMxAA, 42, 241. Optical imaging and spectroscopy.

G54.1+0.3

RA: 19^h30^m31^s
Dec: +18°52'

1-GHz flux/Jy: 0.5
Spectral index: 0.1

Size/arcmin: 1.5
Type: F?

Radio: Filled-centre.

X-ray: Centrally concentrated, with extensions and diffuse emission.

Point sources: Central pulsar.

Distance: HI absorption suggests 4.5–9 kpc, association with CO suggest 8.2 kpc.

References:

- Green 1985, MNRAS, 216, 691. Radio at 2.7 GHz (7'' × 20'').
Reich *et al.* 1985, A&A, 151, L10. Effelsberg 100-m at 4.75 GHz (2.4' : $S = 0.37 \pm 0.04$ Jy)
Velusamy & Becker 1988, AJ, 95, 1162. VLA at 1.4 (14'': $S = 0.48 \pm 0.03$ Jy), 1.6 (14'': $S = 0.42 \pm 0.03$ Jy) and 5 GHz (5'': $S = 0.33 \pm 0.02$ Jy),
Ooty at 327 MHz ($S = 0.50 \pm 0.08$ Jy), plus review of flux densities.
Seward 1989, AJ, 97, 481. Einstein observations.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Lu *et al.* 2001, A&A, 370, 570. ROSAT and ASCA observations.
Lu *et al.* 2002, ApJ, 568, L49. Chandra observations.
Camilo *et al.* 2002, ApJ, 574, L71. Pulsar detection.
Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.
Leahy *et al.* 2008, AJ, 136, 1477. VGPS at 1.4 GHz (1') including HI.
Koo *et al.* 2008, ApJ, 673, L147. IR observations of surroundings.

G54.4–0.3

RA: 19^h33^m20^s
Dec: +18°56'

1-GHz flux/Jy: 28
Spectral index: 0.5

(HC40)

Size/arcmin: 40
Type: S

Has been called G54.5–0.3.

Radio: Shell, in complex region.

Optical: Faint filaments.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 34.4 \pm 5.0$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Caswell 1985, AJ, 90, 1224. DRAO at 1.4 GHz ($1'.3 \times 2'.6 : S = 18 \pm 4$ Jy).
 Velusamy *et al.* 1986, JApA, 7, 105. WSRT at 609 MHz ($50'' \times 191''$ smoothed to $100'' \times 200''$).
 Junkes *et al.* 1992, A&AS, 96, 1. Surrounding CO.
 Junkes *et al.* 1992, A&A, 261, 289. Nearby IRAS sources.
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.

G55.0+0.3

RA: 19^h32^m00^s
Dec: +19°50'

1-GHz flux/Jy: 0.5?
Spectral index: 0.5?

Size/arcmin: 20 × 15?
Type: S

Has been called G55.2+0.5.

Radio: Faint, partial shell.

Distance: Association with HI features implies 14 kpc.

Point sources: Old pulsar nearby.

References:

- Matthews *et al.* 1998, ApJ, 493, 312. WSRT at 327 MHz ($1'.0 \times 2'.9 : S = 0.98 \pm 0.15$ Jy), DRAO at 1.4 GHz ($1'.0 \times 2'.9 : S = 0.25 \pm 0.12$ Jy), plus HI observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G55.7+3.4

RA: 19^h21^m20^s
Dec: +21°44'

1-GHz flux/Jy: 1.4
Spectral index: 0.6

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Point sources: Old pulsar within the boundary of the remnant.

References:

- Goss *et al.* 1977, A&A, 61, 93. WSRT observations at 610 MHz ($57'' \times 156'' : S = 1.9 \pm 0.2$ Jy) and 1415 MHz ($27'' \times 72'' : S = 1.0 \pm 0.1$ Jy).

G57.2+0.8

(4C21.53)

RA: 19^h34^m59^s
Dec: +21°57'

1-GHz flux/Jy: 1.8?
Spectral index: ?

Size/arcmin: 12?
Type: S?

Radio: Extended non-thermal arc.

Point sources: Near the millisecond pulsar, but not thought to be related.

References:

- Sieber & Seiradakis 1984, A&A, 130, 257. Effelsberg 100-m at 1.4 GHz ($8'.8 : 1.34 \pm 0.1$), 2.7 GHz ($4'.3 : 0.86 \pm 0.1$), plus other surveys of the area.
 Caswell *et al.* 1985, AJ, 90, 488. DRAO at 1.4 GHz ($1' \times 3'$).

G59.5+0.1

RA: 19^h42^m33^s
Dec: +23°35'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 15
Type: S

Has been called G59.6+0.1.

Radio: Incomplete shell.

Optical: Diffuse shell.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz ($1\text{'}0 \times 2\text{'}5 : S = 5.1 \pm 0.2$ Jy), and northern sky survey at 4.9 GHz.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.

G59.8+1.2

RA: 19^h38^m55^s
Dec: +24°19'

1-GHz flux/Jy: 1.6
Spectral index: 0.5

Size/arcmin: 20 × 16?
Type: ?

Has been called G59.7+1.2.

Radio: Poorly defined source.

Optical: Faint diffuse emission and filaments.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
Boumis *et al.* 2005, A&A, 443, 175. Optical observations.

G63.7+1.1

RA: 19^h47^m52^s
Dec: +27°45'

1-GHz flux/Jy: 1.8
Spectral index: 0.3

Size/arcmin: 8
Type: F

Radio: Centrally brightened, with core.

References:

Wallace *et al.* 1997, AJ, 114, 2068. WSRT at 1.4 GHz ($14\text{''} \times 26\text{''} : S = 1.63$ Jy), DRAO at 1.4 GHz (smoothed to 2'), plus review of flux densities and other observations.

G65.1+0.6

RA: 19^h54^m40^s
Dec: +28°35'

1-GHz flux/Jy: 5.5
Spectral index: 0.61

Size/arcmin: 90 × 50
Type: S

Radio: Large, faint shell.

Point sources: Old pulsar nearby.

Distance: Possible association with HI suggests 9 kpc.

References:

Landecker *et al.* 1990, A&A, 232, 207. DRAO at 408 MHz ($3\text{'}5 \times 7\text{'}0 : S = 9.5 \pm 0.1$ Jy), and 1.4 GHz ($1\text{'}0 \times 2\text{'}0 : S = 5.4 \pm 1.0$ Jy).
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Tian & Leahy 2006, A&A, 455, 1053. CGPS at 408 MHz ($2\text{'}8 \times 5\text{'}9 : S = 8.6 \pm 0.8$ Jy) and 1.4 GHz ($0\text{'}8 \times 1\text{'}7 : S = 4.9 \pm 0.5$ Jy) including HI.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3\text{'}$: $S = 9.1 \pm 1.0$ Jy) and 1420 MHz ($\sim 1\text{'}$: $S = 3.9 \pm 0.5$ Jy), including review of flux densities.

G65.3+5.7

RA: 19^h33^m00^s
Dec: +31°10'

1-GHz flux/Jy: 52?
Spectral index: 0.6?

Size/arcmin: 310 × 240
Type: S?

Has been called G65.2+5.7.

Radio: Large, faint ring? near S91 and S94.

Optical: Filamentary ring.

X-ray: Diffuse, centrally brightened.

Distance: Optical proper motions and velocities indicates 0.8 kpc.

References:

- Gull *et al.* 1977, ApJ, 215, L69. Optical plates.
 Reich *et al.* 1979, A&A, 72, 270. Effelsberg 100-m observations at 1.42 GHz (smoothed to 11' : $S = 42.4 \pm 1.6$ Jy), estimate $S_{408\text{ MHz}} = 91 \pm 5$ Jy from previous sky survey.
 Lozinskaya 1981, SVAL, 7, 17. Mean optical velocity.
 Rosado 1981, ApJ, 250, 222. Optical interferometry.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Mavromatakis *et al.* 2002, A&A, 388, 355. Optical observations.
 Boumis *et al.* 2004, A&A, 424, 583. Optical observations.
 Shelton *et al.* 2004, ApJ, 615, 275. ROSAT observations.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.

G65.7+1.2

RA: 19^h52^m10^s
Dec: +29°26'

1-GHz flux/Jy: 5.1
Spectral index: varies

DA 495

Size/arcmin: 22
Type: F

Has mistakenly been called G55.7+1.2.

Radio: Centrally brightened with thick shell?

X-ray: Detected.

Point sources: Compact X-ray source near centre.

Distance: HI polarisation observations suggest 1.5 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' : $S = 2.8 \pm 0.4$ Jy), and VRO 37-m at 1.7 GHz ($S = 4.4 \pm 0.5$ Jy), plus review of flux densities.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 8.7 \pm 4.9$ Jy, $S_{318\text{ MHz}} = 9.7 \pm 2.2$ Jy.
 Landecker & Caswell 1983, AJ, 88, 1810. DRAO at 1.4 GHz (0'.9 × 1'.5 : $S = 4.4 \pm 0.2$ Jy).
 Velusamy *et al.* 1989, JApA, 10, 161. Ooty at 327 MHz (36'' × 64''), WSRT at 610 MHz (62'') and VLA at 1.4 GHz (36'' and 12''), including IRAS imaging.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2004, ApJ, 607, 855. HI polarisation absorption.
 Arzoumanian *et al.* 2004, ApJ, 610, L101. ROSAT and ASCA observations of compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : $S = 6.5 \pm 0.6$ Jy) and 1420 MHz (~1' : $S = 4.0 \pm 0.2$ Jy), including polarisation and review of flux densities.
 Arzoumanian *et al.* 2008, ApJ, 687, 505. Chandra observations.
 Kothes *et al.* 2008, ApJ, 687, 516. CGPS at 408 MHz (2'.9 × 6'.0 : $S = 6.5 \pm 0.5$ Jy) and 1.4 GHz (0'.82 × 1'.75 : $S = 4.0 \pm 0.2$ Jy), Effelsberg 100-m at 4.85 GHz (2'.45 : $S = 1.6 \pm 0.1$ Jy) and 10.55 GHz (smoothed to 2'.45 : $S = 1.1 \pm 0.1$ Jy), plus review of flux densities.

G67.7+1.8

RA: 19^h54^m32^s
Dec: +31°29'

1-GHz flux/Jy: 1.0
Spectral index: 0.5

Size/arcmin: 15 × 12
Type: S

Radio: Double arc shell.

Optical: Filaments in N.

References:

- Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz ($1'9 \times 1'0$: $S=1.9 \pm 0.1$ Jy), and northern sky survey at 4.9 GHz ($S=0.42 \pm 0.05$ Jy).
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=1.1 \pm 0.1$ Jy) and 1420 MHz ($\sim 1'$: $S=0.68 \pm 0.04$ Jy), including polarisation and review of flux densities.
Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.

G68.6–1.2

RA: 20^h08^m40^s
Dec: +30°37'

1-GHz flux/Jy: 0.7?
Spectral index: 0.0?

Size/arcmin: 23
Type: ?

Radio: Faint, poorly defined source.

References:

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4'3$).
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz ($\sim 1'$: $S=0.57 \pm 0.08$ Jy), including review of flux densities

G69.0+2.7

RA: 19^h53^m20^s
Dec: +32°55'

1-GHz flux/Jy: 120?
Spectral index: varies

CTB 80

Size/arcmin: 80?
Type: ?

An association with a SN in AD1408 has been suggested. Has been called G68.8+2.8. Is it a SNR?

Radio: Compact core, flat spectrum plateau, and steeper spectrum extensions, with spectral break?

Optical: Expanding nebulosity near centre, with filaments to the SW and far NE.

X-ray: Diffuse emission with compact source.

Point sources: Pulsar at western edge of core.

References:

- Angerhofer *et al.* 1981, A&A, 94, 313. WSRT at 610 MHz (56'' × 103'') 1.4 GHz (24'' × 44'') and 5 GHz (7'' × 13''), plus optical.
 Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.
 Sofue *et al.* 1983, PASJ, 35, 437. NRO 45-m at 10.2 GHz (2'.7).
 Velusamy & Kundu 1983, JApA, 3, 253. VLA of compact sources.
 Blair *et al.* 1984, ApJ, 282, 161. Optical images and spectra.
 Wang & Seward 1984, ApJ, 285, 607. Einstein observations.
 Strom *et al.* 1984, A&A, 139, 43. Radio observations of flat spectrum component, VLA 5 GHz (1''.7) and 1.4 GHz.
 Mantovani *et al.* 1985, A&A, 145, 50. Bologna at 408 MHz (2'.6 × 4'.9 : $S = 67.5 \pm 10.5$ Jy), Effelsberg 100-m at 1.41 (9' : $S = 62 \pm 9$ Jy), 1.72 (7'.6 : $S = 66 \pm 5$ Jy), 2.7 (4'.5 : $S = 52 \pm 4$ Jy) and 4.75 GHz (2'.4 : $S = 44 \pm 3.3$ Jy), plus review of flux densities.
 Kulkarni *et al.* 1988, Nature, 331, 50. Pulsar detection.
 Angelini *et al.* 1988, ApJ, 330, L43. EXOSAT spectra.
 Fesen *et al.* 1988, Nature, 334, 229. IRAS of surrounding shell.
 Whitehead *et al.* 1989, MNRAS, 237, 1109. Optical of core.
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
 Hester & Kulkarni 1989, ApJ, 340, 362. Optical imaging and spectroscopy.
 Koo *et al.* 1990, ApJ, 364, 178. Large, expanding HI shell.
 Greidanus & Strom 1990, A&A, 240, 376. Optical kinematics of core.
 Koo *et al.* 1993, ApJ, 417, 196. VLA at 1.4 GHz of surrounding HI (1'), plus IRAS.
 Safi-Harb *et al.* 1995, ApJ, 439, 722. ROSAT observations.
 Srinivasan 1997, ApJ, 489, 170. γ -ray observations of pulsar.
 Mavromatakis *et al.* 2001, A&A, 371, 300. Optical observations.
 Butler *et al.* 2002, A&A, 395, 845. HST detection of pulsar.
 Migliazzo *et al.* 2002, ApJ, 567, L141. Pulsar proper motion study.
 Castelletti *et al.* 2003, AJ, 126, 2114. GMRT at 240 and 618 MHz (26'' × 17'' and 10'' × 6'') and VLA at 324 and 1380 MHz (73'' × 63'' and 93'' × 78'').
 Moon *et al.* 2004, ApJ, 610, L33. Chandra and HST observations of core.
 Li *et al.* 2005, ApJ, 628, 931. Chandra observations of pulsar and surroundings.
 Golden *et al.* 2005, ApJ, 635, L153. High resolution radio observations of pulsar and surroundings.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 72 \pm 7$ Jy) and 1420 MHz ($\sim 1' : S = 56 \pm 5$ Jy), including polarisation and review of flux densities.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.
 Albert *et al.* 2007, ApJ, 669, 1143. γ -ray observations.
 Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion of pulsar.

G69.7+1.0

RA: 20^h02^m40^s
Dec: +32°43'

1-GHz flux/Jy: 2.0
Spectral index: 0.7

Size/arcmin: 16 × 14
Type: S

Radio: Poorly resolved source.

X-ray: Detected.

References:

- Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Yoshita *et al.* 2000, PASJ, 52, 867. ROSAT and ASCA observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.2 \pm 0.4$ Jy) and 1420 MHz ($\sim 1' : S = 1.5 \pm 0.1$ Jy), including review of flux densities.

G73.9+0.9

RA: 20^h14^m15^s
Dec: +36°12'

1-GHz flux/Jy: 9
Spectral index: 0.23

Size/arcmin: 27
Type: S?

Radio: Diffuse, centrally brightened to SW.

Optical: Faint shell.

References:

- Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz ($2'4 : S = 6.7 \pm 0.5$ Jy), plus other flux densities.
 Chastenay & Pineault 1988, in SNRISM, p297. DRAO at 408 MHz ($3'5 \times 5'9$) and 1.4 GHz ($1'0 \times 1'7$).
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ($3'4 \times 5'8 : S = 12.7 \pm 1.2$ Jy) and 1.4 GHz ($1'0 \times 1'7 : S = 7.4 \pm 1.0$ Jy).
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Pineault *et al.* 1996, AJ, 112, 201. DRAO at 1.4 GHz (smoothed to $2'$) for H α .
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Mavromatakis 2003, A&A, 398, 153. Optical observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.0 \pm 1.7$ Jy) and 1420 MHz ($\sim 1' : S = 7.6 \pm 0.6$ Jy), including polarisation and review of flux densities.

G74.0–8.5

RA: 20^h51^m00^s
Dec: +30°40'

1-GHz flux/Jy: 210
Spectral index: varies

Cygnus Loop

Size/arcmin: 230 × 160
Type: S

Has been suggested that this is two overlapping remnants.

Radio: Shell, brightest to the NE, with fainter breakout region to S, with spectral variations.

Optical: Large filamentary loop, brightest to the NE, not well defined to the S or W.

X-ray: Shell in soft X-rays.

Point sources: Several compact radio sources within the boundary of the remnant, including CL4, plus X-ray sources in S.

Distance: Optical proper motion and shock velocity gives 0.44 kpc.

References:

- Green 1990, AJ, 100, 1927. DRAO at 408 MHz (3'3 × 6'7) for spectral index study, plus X-ray and optical.
Graham *et al.* 1991, AJ, 101, 175. Shocked molecular H outside rim in NE.
Shull & Clarke 1991, PASP, 103, 811. Optical spectroscopy of nearside filaments.
Greidanus & Strom 1992, A&A, 257, 265. Optical kinematics.
Shull & Hippelien 1992, ApJ, 383, 714. Optical kinematics and proper motion.
Fesen *et al.* 1992, AJ, 104, 719. H α imagery.
Cornett *et al.* 1992, ApJ, 395, L9. UV imagery.
Long *et al.* 1992, ApJ, 400, 214. Optical of Balmer dominated filament.
Arendt *et al.* 1992, ApJ, 400, 562. IRAS observations.
Hester *et al.* 1994, ApJ, 420, 721. H α , [OIII] and other optical observations of Balmer dominated filaments in NE.
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Levenson *et al.* 1997, ApJ, 484, 304. ROSAT observations.
Leahy *et al.* 1997, AJ, 114, 2081. DRAO at 1.4 GHz (1' × 2'), including polarisation.
Miyata *et al.* 1998, PASJ, 50, 257. ASCA observations.
Miyata *et al.* 1998, PASJ, 50, 475. ASCA observations of compact X-ray source in S.
Leahy & Roger 1998, ApJ, 505, 784. DRAO at 1.4 GHz (1'0 × 1'9) and 408 MHz (3'4 × 6'9), for spectral index studies in comparison with other radio observations.
Levenson *et al.* 1998, ApJS, 118, 541. Optical images.
Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density ($S = 1400 \pm 400$ Jy).
Bohigas *et al.* 1999, ApJ, 518, 324. Optical spectroscopy of surroundings.
Levenson *et al.* 1999, ApJ, 526, 874. ROSAT images.
Blair *et al.* 1999, AJ, 118, 942. HST observations, for distance.
Aschenbach & Leahy 1999, A&A, 341, 602. ROSAT image, and comparison with radio.
Sauvageot *et al.* 1999, A&A, 351, 669. [Nev] observations.
Szentygyorgyi *et al.* 2000, ApJ, 529, 279. [Nev] images.
Danforth *et al.* 2000, AJ, 119, 2319. UV, optical and X-ray comparison of selected regions.
Sankrit *et al.* 2000, AJ, 120, 1925. HST far-UV spectra on non-radiative shock.
Miyata *et al.* 2001, ApJ, 550, 1023. ASCA observations of compact X-ray sources.
Miyata & Tsunemi 2001, ApJ, 552, 624. ASCA spectroscopy of regions in N and E.
Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
Danforth *et al.* 2001, AJ, 122, 938. Far-UV spectroscopy, H α and other optical observations of NE region.
Levenson & Graham 2001, ApJ, 559, 948. HST of SE region.
Uyaniker *et al.* 2002, A&A, 389, L61. Effelsberg 100-m at 2.7 GHz (4/3) including polarisation, and comparison with ROSAT data.
Leahy 2002, AJ, 123, 2689. DRAO at 1.4 GHz (4' × 2') for HI.
Patnaude *et al.* 2002, AJ, 124, 2118. Optical and ROSAT observations of region in SW.
Blair *et al.* 2002, ApJS, 140, 367. UV spectroscopy.
Welsh *et al.* 2002, A&A, 391, 705. Optical absorption to background stars.
Sankrit & Blair 2002, ApJ, 565, 297. UV observations of NE filament.
Levenson *et al.* 2002, ApJ, 576, 798. Chandra observations of W edge.
Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5 : $S = 184 \pm 18$ Jy).
Leahy 2004, MNRAS, 351, 385. Chandra observations of SW.
Uyaniker *et al.* 2004, A&A, 426, 909. Effelsberg 100-m at 2.7 GHz (4/3), with comparison with other data for spectral index studies.
Blair *et al.* 2005, AJ, 129, 2268. HST of outer filaments.
Leahy 2005, AJ, 130, 165. DRAO at 1.4 GHz of SE.
Levenson & Graham 2005, ApJ, 622, 366. Chandra observations of know in SE.
Sun *et al.* 2006, A&A, 447, 937. Urumqi 25-m at 4.8 GHz (9'5 : $S = 90 \pm 9$ Jy), with comparisons with other data for spectral index studies.
Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
Seon *et al.* 2006, ApJ, 644, L175. Far UV observations.
Sankrit *et al.* 2007, AJ, 133, 1383. UV observations of part.
Tsunemi *et al.* 2007, ApJ, 671, 1717. XMM observations of NE to SW.
Miyata *et al.* 2007, PASJ, 59, S163. Suzaku observations of NE.
Nemes *et al.* 2008, ApJ, 675, 1293. XMM observations of NE.
Katsuda *et al.* 2008, ApJ, 680, 1198. Chandra observations of NE.
Uchida *et al.* 2008, ApJ, 688, 1102. XMM observations.

G74.9+1.2

CTB 87

RA: 20^h16^m02^s
Dec: +37°12'

1-GHz flux/Jy: 9
Spectral index: varies

Size/arcmin: 8 × 6
Type: F

Radio: Filled-centre, with high polarisation and high frequency turnover.

X-ray: Centrally brightened.

Distance: HI absorption indicates 12 kpc, optical extinction gives 6.1 kpc.

Point sources: Extragalactic compact source is nearby.

References:

- Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430 \text{ MHz}} = 12.2 \pm 9.7 \text{ Jy}$, $S_{318 \text{ MHz}} = 17.7 \pm 5.0 \text{ Jy}$.
 Weiler & Shaver 1978, A&A, 70, 389. WSRT at 610 MHz ($57'' \times 94''$: $S = 9.1 \pm 1.2 \text{ Jy}$), 1.4 ($24'' \times 40''$: $S = 8.7 \pm 1.2 \text{ Jy}$) and 5 GHz ($24'' \times 40''$: $S = 5.6 \pm 1.3 \text{ Jy}$).
 Geldzahler *et al.* 1980, A&A, 84, 237. Effelsberg 100-m at 2.7 GHz ($4'4$: $S = 7.6 \pm 0.5 \text{ Jy}$).
 Wilson 1980, ApJ, 241, L19. Einstein observations.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption of nearby compact source.
 Sequaist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (smoothed to $40''$: $S = 1.47 \pm 0.19 \text{ Jy}$).
 Green & Gull 1989, MNRAS, 237, 555. VLA at 1.4 GHz ($1'2 \times 1'4$) including HI.
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz, plus review of flux densities.
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ($3'4 \times 5'8$: $S = 11.6 \pm 0.4 \text{ Jy}$) and 1.4 GHz ($1'0 \times 1'7$: $S = 7.2 \pm 0.3 \text{ Jy}$).
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz ($3'5 \times 5'2$: $S = 13.3 \pm 0.8 \text{ Jy}$) and Effelsberg 100-m at 4.8 GHz ($S = 7.5 \pm 0.7 \text{ Jy}$).
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.
 Cho *et al.* 1994, AJ, 108, 634. CO of adjacent molecular clouds.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Wallace *et al.* 1997, A&A, 317, 212. DRAO at 408 MHz ($3'4 \times 5'5$) and 1.4 GHz ($1'0 \times 1'6$) including HI (smoothed to $2'$).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2003, ApJ, 588, 852. CGPS at 1.4 GHz ($1'6 \times 1'$) including HI, plus CO observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S = 11.9 \pm 0.9 \text{ Jy}$) and 1420 MHz ($\sim 1'$: $S = 7.1 \pm 1.1 \text{ Jy}$), including polarisation and review of flux densities.

G76.9+1.0

Size/arcmin: 9
Type: ?

RA: 20^h22^m20^s
Dec: +38°43'

1-GHz flux/Jy: 1.2
Spectral index: 0.60

Radio: Diffuse, non-thermal, with low frequency turnover.

References:

- Landecker *et al.* 1993, A&A, 276, 522. VLA at 1.49 GHz ($14''$), 4.86 GHz ($13'' \times 16''$) and 8.55 GHz ($11'' \times 12''$), including polarisation and review of flux densities.
 Landecker *et al.* 1997, A&AS, 123, 199. Miyun at 232-MHz ($3'8 \times 5'4$).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S = 2.3 \pm 0.2 \text{ Jy}$) and 1420 MHz ($\sim 1'$: $S = 1.35 \pm 0.07 \text{ Jy}$), including polarisation and review of flux densities.

G78.2+2.1

RA: 20^h20^m50^s
Dec: +40°26'

1-GHz flux/Jy: 320
Spectral index: 0.51

DR4, γ Cygni SNR
Size/arcmin: 60
Type: S

Has been called G78.1+1.8.

Radio: In complex region (early catalogues refer to other proposed remnants in this region).

Optical: Faint filaments, spectra indicate a SNR superposed on a HII region.

X-ray: Weak emission from the SE of the remnant.

Point sources: γ-ray and X-ray point source in remnant.

References:

- Higgs *et al.* 1977, AJ, 82, 718. DRAO at 1.4 GHz ($2' \times 3'$: $S = 270 \pm 40$ Jy) plus some 10 GHz ($4'$) survey data, reveals true extent of remnant.
d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.
van den Bergh 1978, ApJS, 38, 119. Optical observations.
Landeker *et al.* 1980, A&AS, 39, 133. DRAO HI observations ($2' \times 3'.1'$).
Higgs *et al.* 1983, AJ, 88, 97. CO of surroundings.
Bohigas *et al.* 1983, RMxAA, 8, 155. Optical spectra, find thermal only.
Braun & Strom 1986, A&AS, 63, 345. WSRT HI observations.
Fukui & Tatematsu 1988, in SNRISM, p261. CO observations of the vicinity ($2'.7'$).
Green 1989, MNRAS, 238, 737. OH observations.
Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ($3'.4 \times 5'.8$: $S = 480 \pm 60$ Jy) and 1.4 GHz ($1'.0 \times 1'.7$: $S = 270 \pm 40$ Jy).
Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz ($3'.5 \times 5'.2$: $S = 540 \pm 40$ Jy) and Effelsberg 100-m at 4.8 GHz ($S = 150 \pm 15$ Jy).
Esposito *et al.* 1996, ApJ, 461, 820. Associated γ-ray emission.
Brazier *et al.* 1996, MNRAS, 281, 1033. γ-ray and X-ray point source.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Zhang *et al.* 1997, A&A, 324, 641. Multi-frequency radio comparison.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
Uchiyama *et al.* 2002, ApJ, 571, 866. ASCA observations.
Mavromatakis 2003, A&A, 408, 237. Optical observations.
Bykov 2004, A&A, 427, L21. Hard X-ray observations.
Becker *et al.* 2004, ApJ, 615, 897. Chandra and other observations of compact source.
Weisskopf *et al.* 2006, ApJ, 652, 387. Chandra and other observations of compact source.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S = 500 \pm 35$ Jy) and 1420 MHz ($\sim 1'$: $S = 226 \pm 19$ Jy), including review of flux densities.
Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.
Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.
Ladouceur & Pineault 2008, A&A, 490, 197. CGPS at 408 MHz ($2'.9 \times 4'.5$) and 1.4 GHz ($0'.8 \times 1'.5$).

G82.2+5.3

RA: 20^h19^m00^s
Dec: +45°30'

1-GHz flux/Jy: 120?
Spectral index: 0.5?

Size/arcmin: 95 × 65
Type: S

W63

Has been called G82.5+5.3.

Radio: Shell in the Cygnus X complex.

Optical: In complex region, but spectra indicate SNR filaments.

X-ray: Detected.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5'$: $S = 59.0 \pm 3.5$ Jy).
Sabbadin 1976, A&A, 51, 159. Optical spectra.
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'.8'$). Incomplete mapping.
Rosado & González 1981, RMxAA, 5, 93. Optical spectra.
Seward 1990, ApJS, 73, 781. Einstein observations.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'.5$: $S = 82.5 \pm 5.5$ Jy).
Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation, of part.
Mavromatakis *et al.* 2004, A&A, 415, 1051. ROSAT, ASCA and optical observations.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S = 144 \pm 12$ Jy) and 1420 MHz ($\sim 1'$: $S = 93 \pm 5$ Jy), including review of flux densities.

G83.0–0.3

RA: 20^h46^m55^s
Dec: +42°52'

1-GHz flux/Jy: 1
Spectral index: 0.4

Size/arcmin: 9 × 7
Type: S

Radio: Incomplete shell.

References:

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 1.2 \pm 0.3$ Jy) and 1420 MHz ($\sim 1' : S = 0.8 \pm 0.1$ Jy), including polarisation and review of flux densities.

G84.2–0.8

RA: 20^h53^m20^s
Dec: +43°27'

1-GHz flux/Jy: 11
Spectral index: 0.5

Size/arcmin: 20 × 16
Type: S

Radio: Elongated shell, with a filament aligned with the major axis.

References:

Matthews *et al.* 1977, A&A, 55, 1. WSRT at 610 MHz ($56'' \times 81'' : S = 12.4 \pm 1.5$ Jy) and Effelsberg 100-m at 2.7 GHz ($4'4 : S = 6.8 \pm 1.3$ Jy).
 Matthews & Shaver 1980, A&A, 87, 255. WSRT at 1415 MHz ($23'' \times 32''$), and Effelsberg 100-m at 2.7 GHz ($4'4 : S = 5.6 \pm 0.5$ Jy).
 Feldt & Green 1993, A&A, 274, 421. DRAO at 1.4 GHz ($1' \times 1'5$), including HI, plus CO observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 14.5 \pm 0.5$ Jy) and 1420 MHz ($\sim 1' : S = 7.2 \pm 0.8$ Jy), including review of flux densities.

G85.4+0.7

RA: 20^h50^m40^s
Dec: +45°22'

1-GHz flux/Jy: ?
Spectral index: 0.2

Size/arcmin: 24
Type: S

Radio: Faint, incomplete shell, within larger thermal shell.

X-ray: Centrally brightened.

Distance: HI observations suggest 3.5 kpc.

References:

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($4'4 \times 2'8 : S < 0.45$ Jy) and 1.4 GHz ($1'1 \times 0'8$), plus HI, X-ray and optical data.
 Jackson *et al.* 2008, ApJ, 674, 936. XMM and HI observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 2.9 \pm 0.5$ Jy) and 1420 MHz ($\sim 1' : S = 2.3 \pm 0.2$ Jy), including review of flux densities.

G85.9–0.6

RA: 20^h58^m40^s
Dec: +44°53'

1-GHz flux/Jy: ?
Spectral index: 0.2

Size/arcmin: 24
Type: S

Radio: Faint, incomplete shell.

X-ray: Centrally brightened.

Distance: HI observations suggest 4.8 kpc.

References:

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($4'4 \times 2'8 : S < 0.9$ Jy) and 1.4 GHz ($1'1 \times 0'8$), plus HI, X-ray and optical data.
 Jackson *et al.* 2008, ApJ, 674, 936. XMM and HI observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.0 \pm 1.3$ Jy) and 1420 MHz ($\sim 1' : S = 2.2 \pm 0.8$ Jy), including review of flux densities.

G89.0+4.7

HB21

RA: 20^h45^m00^s
Dec: +50°35'

1-GHz flux/Jy: 220
Spectral index: 0.38

Size/arcmin: 120 × 90
Type: S

Radio: Distorted shell (4C50.52, an extragalactic double, is within the boundary of the remnant).

Optical: Filaments and patches.

X-ray: Centrally brightened.

Distance: Various associations imply 0.8 kpc.

References:

- Hirabayashi & Takahashi 1972, PASJ, 24, 231. 30-m dish at 4.2 GHz (11' : $S = 160 \pm 40$ Jy).
 Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' : $S = 148 \pm 16$ Jy), plus optical filaments.
 Hill 1974, MNRAS, 169, 59. Half-Mile Telescope at 1.4 GHz (3' × 3.9').
 Haslam *et al.* 1975, A&A, 39, 453. Effelsberg 100-m at 2.7 GHz (4'.4)
 Fukui & Tatematsu 1988, in SNRISM, p261. CO observations of the vicinity (2'.7).
 Tatematsu *et al.* 1990, A&A, 237, 189. DRAO at 408 MHz (3'.5 × 4.5) and 1.4 GHz (1'.0 × 1.3), including HI, plus CO observations of adjacent molecular cloud.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Leahy & Aschenbach 1996, A&A, 315, 260. ROSAT observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koo *et al.* 2001, ApJ, 552, 175. NRAO 12-m and other CO observations (27'' and 45'') of eastern part.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'.5 : $S = 228 \pm 5$ Jy).
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1') including polarisation.
 Byun *et al.* 2006, ApJ, 637, 283. CO observations of surroundings.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.
 Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
 Leahy 2006, ApJ, 647, 1125. CGPS at 408 MHz (2'.8 × 3.7) and 1.4 GHz (0'.8 × 1.1).
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 259 \pm 19$ Jy) and 1420 MHz ($\sim 1' : S = 183 \pm 9$ Jy), including polarisation and review of flux densities.
 Mavromatakis *et al.* 2007, A&A, 461, 991. Optical observations.

G93.3+6.9

DA 530, 4C(T)55.38.1

RA: 20^h52^m25^s
Dec: +55°21'

1-GHz flux/Jy: 9
Spectral index: 0.45

Size/arcmin: 27 × 20
Type: C?

Has been called G93.2+6.7.

Radio: Shell, with two bright limbs, highly polarised.

X-ray: Compact central source.

Distance: HI observations suggest 2.2 kpc.

References:

- Roger & Costain 1976, A&A, 51, 151. DRAO at 1.42 GHz (2' × 2'.4 : $S = 6.9$ Jy).
 Haslam *et al.* 1980, A&A, 92, 57. Effelsberg 100-m at 1.72 GHz (7'.6 : $S = 6.47 \pm 0.52$ Jy) and 2.7 GHz (4'.4 : $S = 5.64 \pm 0.64$ Jy), plus review of flux densities.
 Lalitha *et al.* 1984, A&A, 131, 196. Effelsberg 100-m at 4.75 GHz (smoothed to 3' : $S = 4.01 \pm 0.57$ Jy).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Landecker *et al.* 1999, ApJ, 527, 866. DRAO at 408 MHz (3'.5 × 4.3) and 1.4 GHz (1'.0 × 1.2), including polarisation and HI.
 Foster & Routledge 2003, ApJ, 598, 1005. HI for distance.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.5 \pm 0.7$ Jy) and 1420 MHz, including review of flux densities.
 Jiang *et al.* 2007, ApJ, 670, 1142. Chandra observations.
 Bocchino *et al.* 2008, AdSpR, 41, 407. XMM observations.

G93.7–0.2

RA: 21^h29^m20^s
Dec: +50°50'

1-GHz flux/Jy: 65
Spectral index: 0.65

CTB 104A, DA 551

Size/arcmin: 80
Type: S

Has been called G93.6–0.2 and G93.7–0.3.

Radio: Distorted, faint shell.

Distance: Association with HI features suggests 1.5 kpc.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 18.4 \pm 1.0$ Jy).
 Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ($7.6' : S = 53.5 \pm 5.0$ Jy), plus review of flux densities.
 Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 58 \pm 6$ Jy).
 Mantovani *et al.* 1991, A&A, 247, 545. Effelsberg 100-m at 4.75 GHz (smoothed to $3' : S = 33.5 \pm 4.0$ Jy), including polarisation, plus review of flux densities.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Uyaniker *et al.* 2002, ApJ, 565, 1022. CGPS 1.4 GHz ($54'' \times 49''$), including HI, and 408 MHz ($3.7' \times 2.8'$).
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 67 \pm 6$ Jy) and 1420 MHz ($\sim 1' : S = 35 \pm 4$ Jy), including polarisation and review of flux densities.

G94.0+1.0

RA: 21^h24^m50^s
Dec: +51°53'

1-GHz flux/Jy: 13
Spectral index: 0.48

3C434.1

Size/arcmin: 30×25
Type: S

Radio: Incomplete shell, containing HI shell.

Distance: Association with stellar wind bubble implies 5.2 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 6.1 \pm 0.8$ Jy), and VRO 37-m at 1.7 GHz ($S = 11 \pm 3$ Jy).
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 5.8 \pm 0.4$ Jy). Also NRAO 140-ft at 5 GHz ($6'$).
 Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ($7.6' : S = 12.0 \pm 1.3$ Jy), plus review of flux densities.
 Goss *et al.* 1984, A&A, 138, 469. WSRT at 610 MHz (smoothed to $100'' : S = 16 \pm 1.7$ Jy) and Effelsberg 100-m at 4.75 GHz ($2.4' : S = 7.2 \pm 0.5$ Jy).
 Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 16 \pm 3$ Jy).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation.
 Foster *et al.* 2004, A&A, 417, 79. DRAO at 1.4 GHz, including HI.
 Foster 2005, A&A, 441, 1043. CGPS at 408 MHz ($2.8' \times 3.6'$) and 1.4 GHz ($0.8' \times 1.0'$) for spectral index studies, plus other observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 20 \pm 2$ Jy) and 1420 MHz ($\sim 1' : S = 11.3 \pm 1.0$ Jy), including review of flux densities.

G96.0+2.0

RA: 21^h30^m30^s
Dec: +53°59'

1-GHz flux/Jy: 0.3
Spectral index: 0.5

Size/arcmin: 26
Type: S

Radio: Faint, arc in S, poorly defined in N.

Distance: Association for HI indicates 4 kpc.

References:

- Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz ($2.8' \times 3.5'$) and 1.4 GHz ($50'' \times 61''$) including HI.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 0.42 \pm 0.06$ Jy) and 1420 MHz ($\sim 1' : S = 0.24 \pm 0.02$ Jy), including review of flux densities.

G106.3+2.7

RA: 22^h27^m30^s
Dec: +60°50'

1-GHz flux/Jy: 6
Spectral index: 0.6

Size/arcmin: 60 × 24
Type: C?

Incorporates the pulsar wind nebula G106.6+2.9 (the ‘Boomerang’).

Radio: Faint extended source, which brighter ‘head’ to NE.

X-ray: Pulsar and wind nebula.

Point sources: Pulsar.

References:

- Pineault & Joncas 2000, AJ, 120, 3218. DRAO at 408 MHz ($3.9 \times 3.5 : S = 10.5 \pm 0.3$ Jy) and 1.4 GHz ($1.2 \times 1.0 : S = 4.9 \pm 0.6$ Jy), plus HI.
 Halpern *et al.* 2001, ApJ, 547, 323. X-ray and radio observations of the ‘head’.w
 Halpern *et al.* 2001, ApJ, 552, L125. Pulsar detection.
 Kothes *et al.* 2001, ApJ, 560, 236. CGPS at 1.4 GHz, including HI, plus CO and other observations.
 Kothes *et al.* 2004, ApJ, 607, 855. HI polarisation absorption.
 Ng & Romani 2004, ApJ, 601, 479. Chandra detection of pulsar wind nebula.
 Kothes *et al.* 2006, ApJ, 638, 225. Effelsberg 100-m at 4.85 (2.4), 8.35 (1.4), 10.5 (1.2) and 32 GHz (0.45) of pulsar wind nebula, including polarisation.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 8.6 \pm 1.0$ Jy) and 1420 MHz ($\sim 1' : S = 4.8 \pm 0.5$ Jy), including polarisation and review of flux densities.
 Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

G108.2–0.6

RA: 22^h53^m40^s
Dec: +58°50'

1-GHz flux/Jy: 8
Spectral index: 0.5

Size/arcmin: 70 × 54
Type: S

Radio: Faint shell.

Distance: Possible associated HI structures suggest 3.2 kpc.

References:

- Tian *et al.* 2007, A&A, 465, 907. DRAO at 408 MHz ($2.8 \times 3.3 : S = 11.5 \pm 1.2$ Jy) and 1.4 GHz ($1.0 \times 1.2 : S = 6.6 \pm 0.7$ Jy) including HI.

G109.1–1.0

RA: 23^h01^m35^s
Dec: +58°53'

1-GHz flux/Jy: 22
Spectral index: 0.50

CTB 109

Size/arcmin: 28
Type: S

Radio: Semicircular shell, with the Molecular cloud S152 is to the immediate W.

X-ray: Semicircular shell, with pulsar at W edge.

Point sources: Long period X-ray pulsar.

Distance: Association with HII regions implies 3 kpc.

References:

- Hughes *et al.* 1981, ApJ, 246, L127. WSRT at 610 MHz ($1': S = 40 \pm 5$ Jy) shows bad CLEAN artefacts.
 Blair & Kirshner 1981, Nature, 291, 132. Optical spectra.
 Downes 1983, MNRAS, 203, 695. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 13.0 \pm 1.5$ Jy).
 Sofue *et al.* 1983, PASJ, 35, 447. NRO 45-m at 10.2 GHz ($2'.7$).
 Hughes *et al.* 1984, ApJ, 283, 147. WSRT at 610 MHz ($20'' : S = 26 \pm 3$ Jy) 1.4 GHz ($10''$), DRAO at 1.4 GHz ($1' \times 1'.2 : S = 16.8 \pm 2$ Jy) and Algonquin 46-m at 6.5 GHz ($4'.5 : S = 6.7 \pm 1$ Jy).
 Braun & Strom 1986, A&AS, 63, 345. WSRT HI observations.
 Tatematsu *et al.* 1987, PASJ, 39, 755. NRO 45-m at 10 GHz ($2'.7$), plus polarisation.
 Tatematsu *et al.* 1987, A&A, 184, 279. CO observations of the surroundings ($2'.7$).
 Hanson *et al.* 1988, A&A, 195, 114. EXOSAT of pulsar.
 Morini *et al.* 1988, ApJ, 333, 777. EXOSAT observations.
 Koyama *et al.* 1989, PASJ, 41, 461. X-ray observations of pulsar.
 Coe *et al.* 1989, MNRAS, 238, 649. IRAS observations of surroundings.
 Green 1989, MNRAS, 238, 737. OH observations.
 Tatematsu *et al.* 1990, ApJ, 351, 157. CO of surroundings, plus X-ray observations.
 Davies & Coe 1991, MNRAS, 249, 313. Optical and IR observations near pulsar.
 Hurford & Fesen 1995, MNRAS, 277, 549. ROSAT imaging.
 Rho & Petre 1997, ApJ, 484, 828. ROSAT observations.
 Parmar *et al.* 1998, A&A, 330, 175. X-ray observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Patel *et al.* 2001, ApJ, 563, L45. Chandra observations of pulsar.
 Kothes *et al.* 2002, ApJ, 576, 169. CGPS at 1.4 GHz ($68'' \times 59''$), including HI, plus CO observations.
 Sasaki *et al.* 2004, ApJ, 617, 322. XMM observations.
 Sasaki *et al.* 2006, ApJ, 642, L149. CO observations of surroundings.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 26 \pm 3$ Jy) and 1420 MHz ($\sim 1' : S = 17.4 \pm 1.2$ Jy), including polarisation and review of flux densities.

G111.7–2.1

RA: 23^h23^m26^s
Dec: +58°48'

1-GHz flux/Jy: 2720
Spectral index: 0.77

Cassiopeia A, 3C461

Size/arcmin: 5
Type: S

Presumably the remnant of a late 17th century SN.

Radio: Bright shell with compact knots and extended plateau of emission.

Optical: Fast knots and quasi-stationary flocculli, with many filaments at large radii, and NE ‘jet’.

X-ray: Incomplete shell, with hard spectral component, and compact central source.

Distance: Optical expansion, plus proper motions indicate 3.4 kpc.

References:

- Anderson *et al.* 1991, ApJ, 373, 146. VLA at 1.4 and 5 GHz (1'') for spectral variations.
 Anderson *et al.* 1995, ApJ, 441, 300. VLA at 1.4 GHz and 4.8 GHz (both 1''), for polarisation studies.
 Anderson & Rudnick 1995, ApJ, 441, 307. VLA at 1.4 GHz and 4.8 GHz, for proper motion studies.
 Kassim *et al.* 1995, ApJ, 455, L59. VLA at 74 (25'') and 332 MHz (6''), including spectral index comparisons.
 Anderson & Rudnick 1996, ApJ, 456, 234. VLA for spectral index comparisons.
 Schwarz *et al.* 1997, A&AS, 123, 43. WSRT at 1.4 GHz (30'') for HI absorption.
 Reynoso *et al.* 1997, A&A, 317, 203. VLA at 1.4 GHz (11'' × 14'') for HI absorption.
 O’Sullivan & Green 1999, MNRAS, 303, 575. Flux density changes at 13.5, 15.5 and 16.5 GHz.
 Agüeros & Green 1999, MNRAS, 305, 957. CLFST at 151 MHz for bulk expansion studies.
 Mason *et al.* 1999, AJ, 118, 2908. Flux density at 32 GHz ($S = 194 \pm 5$ Jy).
 Reichart & Stephens 2000, ApJ, 537, 904. Flux density changes at 1405 MHz.
 Chakrabarty *et al.* 2001, ApJ, 548, 800. Chandra observations, including central compact source.
 Gotthelf *et al.* 2001, ApJ, 552, L39. Chandra observations, showing outer shock.
 Fesen 2001, ApJS, 133, 161. Optical imaging and spectroscopy.
 Bleeker *et al.* 2001, A&A, 365, L225. XMM observations.
 Maccarone *et al.* 2001, A&A, 368, 267. BeppoSAX observations.
 Douvion *et al.* 2001, A&A, 369, 589. ISO observations.
 Thorstensen *et al.* 2001, AJ, 122, 297. Optical expansion studies.
 Hwang *et al.* 2001, ApJ, 560, L175. Chandra observations of Doppler shifted lines.
 Fesen *et al.* 2001, AJ, 122, 2644. HST observations.
 Willingale *et al.* 2002, A&A, 381, 1039. XMM observations.
 Mereghetti *et al.* 2002, ApJ, 569, 275. XMM observations of central source.
 Reynoso & Goss 2002, ApJ, 575, 871. VLA at 5 GHz (6''.4 × 6''.0) for H₂CO absorption studies.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz, including polarisation.
 Loinard *et al.* 2003, RMxAA, 15, 267. Sub-mm observations.
 Dunne *et al.* 2003, Nature, 424, 285. Sub-mm dust observations.
 Rho *et al.* 2003, ApJ, 592, 299. IR observations.
 Laming & Hwang 2003, ApJ, 597, 347. Chandra spectroscopy.
 DeLaney & Rudnick 2003, ApJ, 589, 818. Chandra expansion studies.
 Jones *et al.* 2003, ApJ, 587, 227. 2.2 micron polarisation.
 Hines *et al.* 2004, ApJS, 154, 290. Spitzer observations.
 DeLaney *et al.* 2004, ApJ, 613, 343. Chandra proper motion studies.
 Morse *et al.* 2004, ApJ, 614, 727. HST proper motion studies.
 Hwang *et al.* 2004, ApJ, 615, L117. Deep Chandra imaging.
 Krause *et al.* 2004, Nature, 432, 596. IR and molecular line observations of surroundings.
 Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
 Wilson & Batrla 2005, A&A, 430, 561. Observational constraints on dust.
 Krause *et al.* 2005, Science, 308, 1604. IR echoes from surroundings.
 Fesen *et al.* 2006, ApJ, 636, 848. Optical and IR limits for central sources.
 Fesen *et al.* 2006, ApJ, 636, 859. HST observations of outlying knots.
 Vinyäkin 2006, ARep, 50, 143. Secular decrease at 36, 81.5 and 151.5 MHz.
 Laming *et al.* 2006, ApJ, 644, 260. Chandra observations, particularly of NE ‘jet’.
 Fesen *et al.* 2006, ApJ, 645, 283. HST observations for expansion studies.
 Renaud *et al.* 2006, ApJ, 647, L41. γ-ray line observations.
 Lazendic *et al.* 2006, ApJ, 651, 250. Chandra spectroscopy of bright knots.
 Ennis *et al.* 2006, ApJ, 652, 376. Spitzer observations.
 Mookerjea *et al.* 2006, MNRAS, 371, 761. Sub-mm [CII] observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.
 Patnaude & Fesen 2007, AJ, 133, 147. X-ray variability of compact features.
 Albert *et al.* 2007, A&A, 474, 937. γ-ray observations.
 Rho *et al.* 2008, ApJ, 673, 271. Spitzer observations.
 Uchiyama & Aharonian 2008, ApJ, 677, L105. X-ray variability of compact features.
 Rest *et al.* 2008, ApJ, 681, L81. SN light echo.
 Hammell & Fesen 2008, ApJS, 179, 195. HST of outer knots.
 Krause *et al.* 2008, Science, 320, 1195. SN light echo spectrum.

G113.0+0.2

RA: 23^h36^m35^s
Dec: +61°22'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 40 × 17?
Type: ?

Radio: Elongated, extent not well defined.

Distance: Association for HI indicates 3.1 kpc.

Point sources: Contains old pulsar.

References:

Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz ($2'.8 \times 3'.1$) and 1.4 GHz ($49'' \times 55''$) including HI.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$) and 1420 MHz ($\sim 1'$).

G114.3+0.3

RA: 23^h37^m00^s
Dec: +61°55'

1-GHz flux/Jy: 5.5
Spectral index: 0.5

Size/arcmin: 90 × 55
Type: S

Radio: Shell, with HII region S165 within the boundary of the remnant.

Optical: Faint emission in centre and to S.

Distance: Association with HI and other features implies 0.7 kpc.

Point sources: Pulsar near centre of remnant.

References:

Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 3.6$ Jy) and $S_{1.4\text{ GHz}} = 4.4$ Jy from 1.4 GHz survey data, plus HI from Maryland-Green Bank survey.
Kulkarni *et al.* 1993, Nature, 362, 135. Pulsar association.
Fürst *et al.* 1993, A&A, 276, 470. Pulsar association.
Becker *et al.* 1996, A&A, 306, 464. ROSAT of pulsar.
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
Mavromatakis *et al.* 2002, A&A, 383, 1011. Optical observations.
Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz ($49'' \times 55''$), including HI ($1'.0 \times 1'.1$).
Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz ($3'.4 \times 3'.9 : S = 12.0 \pm 6.0$ Jy) and 1.4 GHz ($1'.0 \times 1'.1 : S = 9.8 \pm 0.8$ Jy).
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz ($\sim 1' : S = 5.4 \pm 0.8$ Jy), including review of flux densities.

G116.5+1.1

RA: 23^h53^m40^s
Dec: +63°15'

1-GHz flux/Jy: 10
Spectral index: 0.5

Size/arcmin: 80 × 60
Type: S

Radio: Distinct shell, with high polarisation.

Optical: Detected.

Distance: Association with HI features implies 1.6 kpc.

References:

Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 4.7 \pm 0.4$ Jy) and $S_{1.4\text{ GHz}} = 8.0 \pm 0.8$ Jy from 1.4 GHz survey data, plus HI from Maryland-Green Bank survey.
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz ($49'' \times 55''$), including HI ($1'.0 \times 1'.1$).
Mavromatakis *et al.* 2005, A&A, 435, 141. Optical observations.
Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz ($3'.4 \times 3'.8 : S = 15.0 \pm 1.5$ Jy) and 1.4 GHz ($1'.0 \times 1'.1 : S = 10.6 \pm 0.$ Jy).
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 12.5 \pm 1.6$ Jy) and 1420 MHz ($\sim 1' : S = 10.3 \pm 0.70$ Jy), including polarisation and review of flux densities.

G116.9+0.2

CTB 1

RA: 23^h59^m10^s
Dec: +62°26'

1-GHz flux/Jy: 8
Spectral index: 0.61

Size/arcmin: 34
Type: S

Has been called G117.3+0.1 and G116.9+0.1.

Radio: Incomplete shell.

Optical: Filaments on sky survey.

X-ray: Centrally brightened, with NE ‘breakout’.

Point sources: Pulsar to NE.

Distance: Association with HI features implies 1.6 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 3.9 \pm 1.0$ Jy), and VRO 37-m at 1.7 GHz ($S = 5.5 \pm 2.0$ Jy), plus review of flux densities.
van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 4.2 \pm 0.2$ Jy).
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6.8' : S = 3.0 \pm 0.3$ Jy).
Dickey & Willis 1980, A&A, 85, 55. WSRT at 610 MHz ($56'' \times 64''$) and 1.4 GHz ($22'' \times 25''$).
Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz ($4.4' : S = 4.8 \pm 0.4$ Jy) and $S_{1.4\text{ GHz}} = 7.8 \pm 0.8$ Jy from 1.4 GHz survey data, plus HI from Maryland-Green Bank survey.
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
Landecker *et al.* 1982, AJ, 87, 1379. DRAO at 1.42 GHz and HI ($2' \times 2.3' : S = 8.3 \pm 0.5$ Jy), plus review of flux densities.
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
Hailey & Craig 1994, ApJ, 434, 635. Optical spectroscopy.
Hailey & Craig 1995, ApJ, 455, L151. ROSAT of nearby pulsar.
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
Craig *et al.* 1997, ApJ, 488, 307. ROSAT observations.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz ($69''$), including polarisation.
Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz ($49'' \times 55''$), including HI ($1.0' \times 1.1'$).
Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz ($3.4' \times 3.8' : S = 15.0 \pm 1.5$ Jy) and 1.4 GHz ($1.0' \times 1.1' : S = 8.1 \pm 0.4$ Jy).
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.5 \pm 0.8$ Jy) and 1420 MHz ($\sim 1' : S = 7.0 \pm 0.8$ Jy), including review of flux densities.

G119.5+10.2		CTA 1
RA: 00 ^h 06 ^m 40 ^s	1-GHz flux/Jy: 36	Size/arcmin: 90?
Dec: +72°45'	Spectral index: 0.6	Type: S
Has been called G119.5+10.3.		
Radio: Incomplete shell, with ‘breakout’ to NW.		
Optical: Faint diffuse nebulosities.		
X-ray: Centrally brightened.		
Point sources: γ -ray pulsar.		
Distance: Associated H α shell indicates 1.4 kpc.		
References:		
Sieber <i>et al.</i> 1979, A&A, 74, 361. Effelsberg 100-m at 2.7 GHz (4': $S=23.6\pm2.8$ Jy).		
Sieber <i>et al.</i> 1981, A&A, 103, 393. Effelsberg 100-m at 2.7 GHz and 1.7 GHz (7'.6: $S=31.6\pm2.5$ Jy), and 151 MHz (4': $S=62.6\pm6$ Jy).		
Fesen <i>et al.</i> 1981, ApJ, 247, 148. Optical, including spectra.		
Fesen <i>et al.</i> 1983, ApJS, 51, 337. Deep [OIII] imagery.		
Seward 1990, ApJS, 73, 781. Einstein observations.		
Simonetti 1992, ApJ, 386, 170. VLA observations of background sources for rotation measure studies.		
Pineault <i>et al.</i> 1993, AJ, 105, 1060. DRAO at 1.4 GHz (1': $S=34\pm4$ Jy), plus H α and IRAS.		
Seward <i>et al.</i> 1995, ApJ, 453, 284. ROSAT observations.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Slane <i>et al.</i> 1997, ApJ, 485, 221. ASCA and ROSAT observations.		
Pineault <i>et al.</i> 1997, A&A, 324, 1152. DRAO at 408 MHz (3'.5) and 1.4 GHz (1'.0).		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
Brazier <i>et al.</i> 1998, MNRAS, 295, 819. Studies of central, compact γ -ray source.		
Mavromatakis <i>et al.</i> 2000, A&A, 353, 371. Optical imagery.		
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.		
Reich 2002, in NSPS, p1. Effelsberg 100-m at 1.4 GHz (9'.4).		
Slane <i>et al.</i> 2004, ApJ, 601, 1045. XMM observations of central source.		
Halpern <i>et al.</i> 2004, ApJ, 612, 398. Chandra observations of central nebula, plus optical and radio limits for compact source.		
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.		
Abdo <i>et al.</i> 2008, Science, 322, 1218. γ -ray detection of pulsar.		

G120.1+1.4

RA: 00^h25^m18^s
Dec: +64°09'

1-GHz flux/Jy: 56
Spectral index: 0.65

Tycho, 3C10, SN1572

Size/arcmin: 8
Type: S

This is the remnant of the Tycho's SN of AD1572.

Radio: Shell, brightest to the NE.

Optical: Faint filaments/knots to the NNW, NE and E.

X-ray: Shell, brighter to the NE.

Point sources: Faint radio source near centre of the remnant, thought to be extragalactic.

Distance: HI absorption gives 2–5 kpc, optical proper motion and shock velocity gives 2.4 kpc.

References:

- Duin & Strom 1975, A&A, 39, 33. WSRT at 610 MHz ($57'' \times 64''$) and 5 GHz ($7'' \times 8''$).
Klein *et al.* 1979, A&A, 76, 120. Effelsberg 100-m at 10.7 GHz ($1\farcm2 : S = 13.1 \pm 0.8$ Jy), plus review of flux densities.
Strom *et al.* 1982, MNRAS, 200, 473. WSRT at 1415 MHz ($27'' \times 31''$) from 1971 and 1979, for expansion.
Dickel *et al.* 1982, ApJ, 257, 145. Comparison of radio, X-ray and optical observations.
Reid *et al.* 1982, ApJ, 261, 485. Einstein observations.
Seward *et al.* 1983, ApJ, 266, 287. Einstein observations.
Tan & Gull 1985, MNRAS, 216, 949. Radio at 2.7 GHz ($4''$) from 1980 and 1983, and One-Mile Telescope at 1.4 GHz ($23''$) from 1965 and 1980, for expansion.
Albinson *et al.* 1986, MNRAS, 219, 427. HI observations.
Green & Gull 1987, MNRAS, 224, 1055. VLA HI absorption observations towards central radio source.
Smith *et al.* 1988, ApJ, 325, 288. EXOSAT spectrum and image (smoothed to $13''$).
Dickel *et al.* 1991, AJ, 101, 2151. VLA at 1.4 and 5 GHz ($1\farcm5$).
Wood *et al.* 1992, AJ, 103, 1338. VLA at 5 GHz ($1\farcm5$) polarisation studies.
Fink *et al.* 1994, A&A, 283, 635. X-ray spectra.
Vancura *et al.* 1995, ApJ, 441, 680. X-ray spectra and ROSAT image.
Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
Schwarz *et al.* 1995, A&A, 299, 193. WSRT and Effelsberg 100-m at 1.4 GHz ($50''$), and VLA at 1.4 GHz ($13''$), for neutral hydrogen studies.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Hwang & Gotthelf 1997, ApJ, 475, 665. ASCA observations.
Reynoso *et al.* 1997, ApJ, 491, 816. VLA at 1.4 GHz ($1\farcm4 \times 1\farcm5$) from 1984 and 1994 for expansion studies.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Reynoso *et al.* 1999, AJ, 117, 1827. VLA at 1.4 GHz ($13''$), for HI studies.
Katz-Stone *et al.* 2000, ApJ, 529, 453. VLA at 333 MHz and 1.4 GHz ($9\farcm5 \times 7\farcm7$), for spectral index studies.
Ghavamian *et al.* 2000, ApJ, 535, 266. Optical observations of shock precursors.
Hughes 2000, ApJ, 545, L53. ROSAT X-ray expansion.
Decourchelle *et al.* 2001, A&A, 365, L218. XMM observations.
Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
Douvion *et al.* 2001, A&A, 373, 281. ISO observations.
Aharonian *et al.* 2001, A&A, 373, 292. Upper limit on high energy γ -rays.
Hwang *et al.* 2002, ApJ, 581, 1101. Chandra observations.
Lee *et al.* 2004, ApJ, 605, L113. Observations of molecular clouds in vicinity.
Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
Warren *et al.* 2005, ApJ, 634, 376. Chandra observations.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 86 \pm 5$ Jy) and 1420 MHz ($\sim 1' : S = 40.5 \pm 1.5$ Jy), including polarisation and review of flux densities.
Lee *et al.* 2007, ApJ, 659, L133. H α observations.
Cassam-Chenai *et al.* 2007, ApJ, 665, 315. Chandra observations.
Rest *et al.* 2008, ApJ, 681, L81. SN light echo.
Krause *et al.* 2008, Nature, 456, 617. SN light echo spectrum.

G126.2+1.6

RA: 01^h22^m00^s
Dec: +64°15'

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 70
Type: S?

Radio: Poorly defined shell.

Optical: Filaments, mostly in W.

References:

- Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz ($9': S = 6.8 \pm 0.7$ Jy) and 2.7 GHz ($4.4': S = 3.9 \pm 0.4$ Jy).
 Blair *et al.* 1980, ApJ, 242, 592. Optical detection and spectra.
 Rosado 1982, RMxAA, 5, 127. Optical spectra.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz ($4.4'$) and 4.8 GHz ($2.6'$).
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz ($3.5' \times 3.9': S = 12 \pm 2.5$ Jy) and part at 1.4 GHz ($1.1' \times 1.0'$), plus review of flux densities.
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14.5': S = 6.1 \pm 1.6$ Jy).
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
 Tian & Leahy 2006, A&A, 447, 205. CGPS at 408 MHz ($3.4' \times 3.8': S = 9.7 \pm 3.9$ Jy) and 1.4 GHz ($1.0' \times 1.1': S = 6.7 \pm 2.1$ Jy), plus other observations for spectral index studies.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3': S = 5.7 \pm 0.7$ Jy) and 1420 MHz ($\sim 1': S = 6.4 \pm 1.1$ Jy), including review of flux densities.
 Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz ($9.5': S = 2.6 \pm 0.6$ Jy), including polarisation.
 see also: Sun *et al.* 2007, A&A, 469, 1003. Erratum.

G127.1+0.5

RA: 01^h28^m20^s
Dec: +63°10'

1-GHz flux/Jy: 12
Spectral index: 0.45

Size/arcmin: 45
Type: S

Has been called G127.3+0.7.

Radio: Distinct shell, with bright central source.

Point sources: Flat radio spectrum (extragalactic) source at centre of remnant.

Optical: Detected.

Distance: 1.2–1.3 kpc if associated with NGC 559.

References:

- Caswell 1977, MNRAS, 181, 789. Half-Mile Telescope at 1.42 GHz ($3.5' \times 3.9'$), plus other observations of central source.
 Salter *et al.* 1978, A&A, 66, 77. Effelsberg 100-m at 2.7 GHz ($4.4'$), plus 5 and 8.7 GHz of the central source.
 Pauls 1977, A&A, 59, L13. Effelsberg 100-m at 1.4 GHz ($9': S = 8 \pm 1$ Jy).
 Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz ($9': S = 10.8 \pm 1.3$ Jy).
 Pauls *et al.* 1982, A&A, 112, 120. WSRT at 610 MHz ($56'' \times 62''$) and H α absorption to the point source with the VLA.
 Geldzahler & Shaffer 1982, ApJ, 260, L69. Observations of central source.
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz ($4.4'$) and 4.8 GHz ($2.6'$).
 Goss & van Gorkom 1984, JApA, 5, 425. WSRT H α absorption of central source.
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz ($3.5' \times 3.9': S = 17.9 \pm 2.0$ Jy) and 1.4 GHz ($1.13' \times 1.0': S = 10.1 \pm 0.8$ Jy), plus review of flux densities.
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14.5': S = 14.6 \pm 0.8$ Jy).
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact sources.
 Leahy & Tian 2006, A&A, 451, 251. CGPS at 408 MHz ($3.4' \times 3.8': S = 17.1 \pm 1.7$ Jy) and 1.4 GHz ($1.0' \times 1.2': S = 10.0 \pm 0.8$ Jy).
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3': S = 15.9 \pm 1.0$ Jy) and 1420 MHz ($\sim 1': S = 9.7 \pm 0.6$ Jy), including polarisation and review of flux densities.
 Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz ($9.5': S = 6.3 \pm 0.7$ Jy), including polarisation.
 see also: Sun *et al.* 2007, A&A, 469, 1003. Erratum.

G130.7+3.1

RA: 02^h05^m41^s
Dec: +64°49'

1-GHz flux/Jy: 33
Spectral index: 0.07

3C58, SN1181

Size/arcmin: 9 × 5
Type: F

This is the remnant of the SN of AD1181.

Radio: Filled-centre, highly polarised, with high frequency turnover.

Optical: Faint filaments.

X-ray: Centrally brightened, with faint jet.

Point sources: Central pulsar.

Distance: HI absorption indicates 3.2 kpc.

References:

- Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz ($58'' : S = 26.7 \pm 0.5$ Jy).
 Wilson & Weiler 1976, A&A, 49, 357. WSRT at 610 MHz ($58'' \times 64''$), 1.4 GHz ($24'' \times 27''$) and 5 GHz ($7'' \times 8''$).
 van den Berg 1978, ApJ, 220, L9. Optical observations.
 Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.
 Green & Gull 1982, Nature, 299, 606. HI absorption distance.
 Fesen 1983, ApJ, 270, L53. Optical spectra.
 Reynolds & Aller 1985, AJ, 90, 2312. VLA at 1.4 GHz (2'), for limits of shell.
 Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT spectrum.
 Green 1986, MNRAS, 218, 533. CLFST at 151 MHz ($1/2 \times 1/3 : S = 36 \pm 4$ Jy), plus 2.7 GHz (4''), plus Einstein observations for limit on shell.
 Green 1987, MNRAS, 225, 11P. Flux density increase at 408 MHz.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz ($26''/5 : S = 24.2 \pm 1.4$ Jy).
 Reynolds & Aller 1988, ApJ, 327, 845. VLA at 1.4 (2'') and 4.9 GHz (2'').
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ($90'' : S = 15.0 \pm 2.0$ Jy), plus review of flux densities.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Green & Scheuer 1992, MNRAS, 258, 833. IRAS upper limits.
 Roberts *et al.* 1993, A&A, 274, 427. HI absorption.
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.
 Helfand *et al.* 1995, ApJ, 453, 741. ROSAT observations.
 Torii *et al.* 2000, PASJ, 52, 875. ASCA observations.
 Bietenholz *et al.* 2001, ApJ, 560, 772. VLA at 74 MHz ($26'' : S = 33.6$ Jy) and 327 MHz ($8''/2 : S = 33.9$ Jy), for spectral index and expansion studies.
 Bocchino *et al.* 2001, A&A, 369, 1078. XMM observations.
 Murray *et al.* 2002, ApJ, 568, 226. Chandra pulsar detection.
 Camilo *et al.* 2002, ApJ, 571, L41. Pulsar detection in radio.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz ($26''$) for polarised intensity.
 Slane *et al.* 2002, ApJ, 571, L45. Chandra observations.
 Slane *et al.* 2004, ApJ, 616, 403. Deep Chandra imaging.
 Bietenholz 2006, ApJ, 645, 1180. VLA at 1.4 GHz (1''.36) for expansion studies.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 32.2 \pm 2.0$ Jy) and 1420 MHz ($\sim 1' : S = 31.9 \pm 1.0$ Jy), including polarisation and review of flux densities.
 Gotthelf *et al.* 2007, ApJ, 654, 267. XMM observations.
 Slane *et al.* 2008, ApJ, 676, L33. IR observations.
 Fesen *et al.* 2008, ApJS, 174, 379. Optical observations for proper motion studies.
 Shearer & Neustroev 2008, MNRAS, 390, 235. Optical observations of pulsar nebula.
 Shibanov *et al.* 2008, A&A, 486, 273. Optical observations of pulsar nebula.

G132.7+1.3		HB3
RA: 02 ^h 17 ^m 40 ^s	1-GHz flux/Jy: 45	Size/arcmin: 80
Dec: +62°45'	Spectral index: 0.6	Type: S

Has been called G132.4+2.2.

Radio: Faint shell, adjacent to W3/4/5 complex.

Optical: Complete, filamentary shell, shock excited spectra.

X-ray: Partial shell.

Point sources: Pulsar nearby.

Distance: Interaction with surroundings suggests 2.2 kpc.

References:

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 33.8 \pm 7.0$ Jy).
 d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.
 Read 1981, MNRAS, 194, 863. Radio at 151 MHz ($4.4'$) and 1.4 GHz ($2'$) showing HI shell.
 Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
 Fesen & Gull 1983, PASP, 95, 196. Optical image.
 Leahy *et al.* 1985, ApJ, 294, 183. Einstein observations.
 Landecker *et al.* 1987, AJ, 94, 111. DRAO at 408 MHz ($3.5' \times 4' : S = 75 \pm 15$ Jy), plus review of flux densities.
 Routledge *et al.* 1991, A&A, 247, 529. DRAO at 1.4 GHz ($1.0' \times 1.1'$) for HI, plus CO observations.
 Fesen *et al.* 1995, AJ, 110, 2876. Optical imaging and spectroscopy, DRAO at 408 MHz ($3.5' \times 4'$) and 1.4 GHz ($1.0' \times 1.1'$).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14.5' : S = 51.5 \pm 3.5$ Jy).
 Tian & Leahy 2005, A&A, 436, 187. CGPS at 408 MHz ($3.4' \times 3.8'$) and 1.4 GHz ($1.0' \times 1.1'$), for spectral index studies.
 see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 61 \pm 9$ Jy) and 1420 MHz ($\sim 1' : S = 29.4 \pm 2.7$ Jy), including review of flux densities.
 Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
 Green 2007, BASI, 35, 77. Review of radio spectrum.
 Shi *et al.* 2008, A&A, 487, 601. Urumqi 25-m at 4.8 GHz ($9.5'$), plus other survey observations for spectral studies.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

G156.2+5.7

RA: 04 ^h 58 ^m 40 ^s	1-GHz flux/Jy: 5	Size/arcmin: 110
Dec: +51°50'	Spectral index: 0.5	Type: S

Radio: Faint shell, brighter in E and W.

Optical: Filamentary ring and smaller patchy ring.

X-ray: Faint shell.

References:

- Pfeffermann *et al.* 1991, A&A, 246, L28. ROSAT detection.
 Reich *et al.* 1992, A&A, 256, 214. Effelsberg 100-m at 1.4 ($9' : S = 4.2 \pm 1.0$ Jy) and 2.7 GHz ($4.3' : S = 3.0 \pm 1.0$ Jy), plus HI and IRAS.
 Yamauchi *et al.* 1993, PASJ, 45, 795. Hard X-ray observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Yamauchi *et al.* 1999, PASJ, 51, 13. ASCA observations of some regions.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz, including polarisation.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 8.1 \pm 1.3$ Jy), including review of flux densities.
 Xu *et al.* 2007, A&A, 470, 969. Urumqi 25-m at 4.8 GHz ($9.5' : S = 2.5 \pm 0.5$ Jy), including polarisation.
 Gerardy & Fesen 2007, MNRAS, 376, 929. Optical observations.

G160.9+2.6		HB9
RA: 05 ^h 01 ^m 00 ^s Dec: +46°40'	1-GHz flux/Jy: 110 Spectral index: 0.64	Size/arcmin: 140 × 120 Type: S

Has been called G160.5+2.8 and G160.4+2.8.

Radio: Large, filamentary shell.

Optical: Incomplete shell.

X-ray: Centrally brightened.

Point sources: Pulsar within boundary of the remnant, plus several nearby compact radio sources.

Distance: Various observations suggests less than 4 kpc.

References:

- d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.
 Damashek *et al.* 1978, ApJ, 225, L31. Pulsar.
 Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
 Dwarakanath *et al.* 1982, JApA, 3, 207. Radio observations at 34.5 MHz ($26' \times 40' : S = 750 \pm 150$ Jy), plus review of flux densities.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source.
 Sequaist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
 Leahy 1988, ApJ, 322, 917. Einstein observations.
 Leahy & Roger 1991, AJ, 101, 1033. DRAO at 408 MHz ($3'5 \times 4'8$) and 1.4 GHz ($1'0 \times 1'4$), including HI and discussion of distance.
 Yamauchi & Koyama 1993, PASJ, 45, 545. Hard X-ray observations.
 Leahy & Aschenbach 1995, A&A, 293, 853. ROSAT observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Leahy *et al.* 1998, A&A, 339, 601. Miyun at 232 MHz ($3'8 \times 5'2$), CLFST at 151 MHz ($4'2 \times 5'8$) and Effelsberg 100-m at 4.7 GHz ($2'5$) for spectral index studies.
 Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density ($S = 1130 \pm 340$ Jy).
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'5 : S = 91 \pm 3$ Jy).
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 102 \pm 7$ Jy) and 1420 MHz ($\sim 1' : S = 54.0 \pm 2.9$ Jy), including polarisation and review of flux densities.
 Leahy & Tian 2007, A&A, 461, 1013. CGPS at 408 MHz ($2'8 \times 3'9 : S = 117.8 \pm 5.3$ Jy) and 1.4 GHz ($58'' \times 80'' : S = 65.9 \pm 3.4$ Jy).

G166.0+4.3		VRO 42.05.01
RA: 05 ^h 26 ^m 30 ^s Dec: +42°56'	1-GHz flux/Jy: 7 Spectral index: 0.37	Size/arcmin: 55 × 35 Type: S

Radio: Two arcs of strikingly different radii.

Optical: Nearly complete ring.

X-ray: Predominantly in SW.

Distance: HI indicates 4.5 kpc.

References:

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 5.2 \pm 1.0$ Jy).
 Lozinskaya 1979, AuPh, 32, 113. Hα interferometry.
 Landecker *et al.* 1982, ApJ, 261, L41. DRAO at 1.4 GHz ($1'0 \times 1'4$), plus review of flux densities.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Pineault *et al.* 1985, A&A, 151, 52. VLA at 1.4 GHz ($16'' \times 20''$) of part of remnant, and optical observations.
 Pineault *et al.* 1987, ApJ, 315, 580. DRAO and VLA combined at 1.4 GHz ($20''$).
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Braun & Strom 1986, A&AS, 63, 345. WSRT HI Observations.
 Landecker *et al.* 1989, MNRAS, 237, 277. DRAO at 1.4 GHz ($1'0 \times 1'4$), including HI.
 Burrows & Guo 1994, ApJ, 421, L19. ROSAT images and spectra.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Guo & Burrows 1997, ApJ, 480, L51. ASCA observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Leahy & Tian 2005, A&A, 440, 929. CGPS at 408 MHz ($3'4 \times 5'0$) and 1.4 GHz ($1'0 \times 1'4$), for spectral index studies.
 see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 8.1 \pm 0.9$ Jy) and 1420 MHz ($\sim 1' : S = 5.1 \pm 0.4$ Jy), including polarisation and review of flux densities.

G179.0+2.6

RA: 05^h53^m40^s
Dec: +31°05'

1-GHz flux/Jy: 7
Spectral index: 0.4

Size/arcmin: 70
Type: S?

Radio: Thick shell, with background extragalactic sources near centre.

References:

- Fürst & Reich 1986, A&A, 154, 303. Effelsberg 100-m at 1.4 (9'.4), 2.7 (4'.3) and 4.75 GHz (2'.4).
Fürst *et al.* 1989, A&A, 223, 66. Observations of central, extragalactic source.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.

G180.0–1.7

RA: 05^h39^m00^s
Dec: +27°50'

1-GHz flux/Jy: 65
Spectral index: varies

Size/arcmin: 180
Type: S

Radio: Large faint shell, with spectral break.

Optical: Wisp ring.

X-ray: Possible detection.

Point sources: Pulsar within boundary, with faint wind nebula.

Distance: Optical absorption towards stars indicates >0.36 and <0.88 kpc.

References:

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
Sofue *et al.* 1980, PASJ, 32, 1. Effelsberg 100-m at 5 GHz (2'.6) of parts.
Kundu *et al.* 1980, A&A, 92, 225. Effelsberg 100-m at 2.7 GHz (5'.5 : S = 34.9 ± 4 Jy) and 1.6 GHz (10' : S = 60.2 ± 6 Jy).
Angerhofer & Kundu 1981, AJ, 86, 1003. Arecibo at 430 MHz (9' : S = 97 ± 20 Jy).
Fürst *et al.* 1982, A&A, 115, 428. Observations of compact radio sources near the remnant.
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
Fürst & Reich 1986, A&A, 163, 185. Effelsberg 100-m at 1.4, 2.7 and 4.7/5.0 GHz (9'.4, 4'.3 and 2'.4/2'.6).
Sauvageot *et al.* 1990, A&A, 227, 183. EXOSAT possible detection.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Anderson *et al.* 1996, ApJ, 468, L55. Pulsar detection.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'.5 : S = 77 ± 10 Jy).
Romani & Ng 2003, ApJ, 585, L41. Chandra of pulsar.
Kramer *et al.* 2003, ApJ, 593, L31. Pulsar observations.
Sallmen & Welsh 2004, A&A, 426, 555. Optical absorption towards background stars.
Drew *et al.* 2005, MNRAS, 362, 753. Hα imaging.
Ng *et al.* 2007, ApJ, 654, 487. Pulsar and wind nebula observations.
Xiao *et al.* 2008, A&A, 482, 783. Urumqi 25-m at 4.8 GHz (9'.5 : S = 15.4 ± 3.0 Jy) and Effelsberg 100-m at 2.6 GHz (4'.4 : S = 34.6 ± 4.0 Jy).

G182.4+4.3

RA: 06^h08^m10^s
Dec: +29°00'

1-GHz flux/Jy: 1.2
Spectral index: 0.4

Size/arcmin: 50
Type: S

Radio: Incomplete shell.

References:

- Kothes *et al.* 1998, A&A, 331, 661. Effelsberg 100-m at 1.4, 2.7, 4.9 and 10.5 GHz (9'.4 : S = 0.36 ± 0.08 Jy, 4'.4 : S = 0.25 ± 0.04 Jy, 2'.5 : S = 0.20 ± 0.02 Jy and 1'.2 : S = 0.15 ± 0.03 Jy), plus X-ray upper limit.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz and 4.9 GHz (3').

G184.6–5.8

RA: 05^h34^m31^s
Dec: +22°01'

1-GHz flux/Jy: 1040
Spectral index: 0.30

Crab Nebula, 3C144, SN1054

Size/arcmin: 7 × 5
Type: F

This is the remnant of the SN of AD1054.

Radio: Filled-centre, central pulsar, with faint ‘jet’ (or tube) extending from the N edge.

Optical: Strongly polarised filaments, diffuse synchrotron emission, with ‘jet’ faintly visible.

X-ray: Central ‘torus’ around the pulsar.

Point sources: Pulsar powering the remnant.

Distance: Proper motions and radial velocities give 2 kpc.

References:

- Velusamy 1984, Nature, 308, 251. VLA at 1.4 GHz, radio detection of ‘jet’.
 Velusamy 1985, MNRAS, 212, 359. VLA at 1.4 GHz (15'').
 Aller & Reynolds 1985, ApJ, 293, L73. Flux density decrease.
 Fesen & Gull 1986, ApJ, 306, 259. Deep [OIII] imagery of ‘jet’.
 Marcellin *et al.* 1990, A&A, 228, 471. Optical of ‘jet’.
 Fesen & Blair 1990, ApJ, 351, L45. Optical identification of dust in filaments.
 Hester *et al.* 1990, ApJ, 357, 539. Optical and IR images.
 Hickson & van den Bergh 1990, ApJ, 365, 224. Optical polarisation.
 Bietenholz & Kronberg 1990, ApJ, 357, L13. VLA at 1.4 GHz (1''.8).
 Bietenholz & Kronberg 1992, ApJ, 393, 206. VLA at 1.5 and 5 GHz (1''.8) and 1.5 and 14 GHz (6''.5) for spectral studies.
 Fesen & Staker 1993, MNRAS, 263, 69. [OIII] imaging of ‘jet’, and proper motion studies.
 Kassim *et al.* 1993, AJ, 106, 2218. VLA at 74 MHz (20'').
 Véron-Cetty & Woltjer 1993, A&A, 270, 370. Continuum and [OIII] photometry.
 Murdin 1994, MNRAS, 269, 89. Deep Hα plates and spectroscopy of halo.
 Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.
 MacAlpine *et al.* 1994, ApJ, 432, L131. Prominent optical knots.
 Rudy *et al.* 1994, ApJ, 426, 646. IR spectroscopy.
 Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
 Lawrence *et al.* 1995, AJ, 109, 2635. Optical imaging spectroscopy.
 Frail *et al.* 1995, ApJ, 454, L129. VLA at 333 MHz (20'') for limits on shell.
 Bietenholz *et al.* 1997, ApJ, 490, 291. Comparison of VLA observations at 74 MHz, 327 MHz, 1.5 GHz and 5 GHz for spectral index studies.
 Blair *et al.* 1997, ApJS, 109, 473. HST imaging.
 van der Meulen *et al.* 1998, A&A, 330, 321. γ-ray observations.
 Tanimori *et al.* 1998, ApJ, 492, L33. γ-ray observations.
 Nugent 1998, PASP, 110, 831. Optical expansion.
 Hillas *et al.* 1998, ApJ, 503, 744. γ-ray spectrum.
 Sankrit *et al.* 1998, ApJ, 504, 344. HST images.
 Wallace *et al.* 1999, ApJS, 124, 181. DRAO at 1.4 GHz (1''.0 × 2''.8), plus Effelsberg 100-m, for HI studies.
 Greiveldinger & Aschenbach 1999, ApJ, 510, 305. X-ray variability of torus.
 Weisskopf *et al.* 2000, ApJ, 536, L81. Chandra observations.
 Sollerman *et al.* 2000, ApJ, 537, 861. HST observations.
 Aharonian *et al.* 2000, ApJ, 539, 317. High energy γ-ray observations.
 Carramiñana *et al.* 2000, ApJ, 542, 974. Optical spectra of pulsar.
 Oser *et al.* 2001, ApJ, 547, 949. γ-ray observations.
 Willingale *et al.* 2001, A&A, 365, L212. XMM observations.
 Douvion *et al.* 2001, A&A, 373, 281. ISO observations.
 Bietenholz *et al.* 2001, ApJ, 560, 254. Multi-epoch VLA observations, showing variations near the pulsar.
 Bandiera *et al.* 2002, A&A, 386, 1044. 1.3 mm observations.
 Hester *et al.* 2002, ApJ, 577, L49. HST and Chandra multi-epoch observations.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz, including polarisation.
 Atkins *et al.* 2003, ApJ, 595, 803. High energy γ-ray observations.
 Green *et al.* 2004, MNRAS, 355, 1315. Sub-mm and ISO observations.
 Mori *et al.* 2004, ApJ, 609, 186. Chandra observations.
 Čadež *et al.* 2004, ApJ, 609, 797. Optical observations.
 Aharonian *et al.* 2004, ApJ, 614, 897. γ-ray observations.
 Bietenholz *et al.* 2004, ApJ, 615, 794. VLA at 5 GHz (1''.4) and HST multi-epoch observation for proper motion studies.
 Melatos *et al.* 2005, ApJ, 633, 931. Multi-epoch near-IR observations central region.
 Seward *et al.* 2006, ApJ, 636, 873. Chandra observations of scattering halo.
 Temim *et al.* 2006, AJ, 132, 1610. Spitzer observations.
 Aharonian *et al.* 2006, A&A, 457, 899. γ-ray observations.
 Seward *et al.* 2006, ApJ, 652, 1277. Chandra observations.
 MacAlpine *et al.* 2007, AJ, 133, 81. Optical spectroscopy.
 Kaplan *et al.* 2008, ApJ, 677, 1201. HST proper motion of pulsar.
 Rudie *et al.* 2008, MNRAS, 384, 1200. [OIII] observations of ‘jet’, for proper motion.

G189.1+3.0

RA: 06^h17^m00^s
Dec: +22°34'

1-GHz flux/Jy: 160
Spectral index: 0.36

IC443, 3C157

Size/arcmin: 45
Type: C

Radio: Limb-brightened to NE, with faint extension to the E.

Optical: Brightest to the NE, with faint filaments outside the NE boundary.

X-ray: Shell, brightest to the NE, plus compact source with nebula.

Point sources: X-ray source and nebula in S.

Distance: Mean optical velocity suggests 0.7–1.5 kpc, association with S249 gives 1.5–2 kpc.

References:

- Fesen 1984, ApJ, 281, 658. Optical of filament to far NE.
Erickson & Mahoney 1985, ApJ, 290, 596. TPT at 4 frequencies between 31 and 74 MHz ($13' \times 11'$ to $5'4 \times 4'7$), plus review of flux densities.
Braun & Strom 1986, A&A, 164, 193. WSRT at 327 MHz ($72'' \times 185''$) and 1.4 GHz ($17'' \times 43''$), plus HI and IRAS.
Green 1986, MNRAS, 221, 473. CLFST at 151 MHz ($1'.2 \times 3'.1$) and Half-Mile Telescope at 1.4 GHz ($2'.1 \times 5'.4$).
Mufson *et al.* 1986, AJ, 92, 1349. Radio, IR, optical, UV and X-ray comparison, including VLA at 1.6 GHz ($3''.8 \times 3''.3$ and $40''$).
Dickel *et al.* 1989, AJ, 98, 1363. VLA at 1.4 GHz ($1''.1 \times 1''.2$) of NE.
Petre *et al.* 1988, ApJ, 335, 215. Einstein and other X-ray observations.
Burton *et al.* 1990, ApJ, 355, 197. IR observations of shocked O₂.
Wood *et al.* 1991, AJ, 102, 224. VLA at 5 GHz ($3''.6 \times 3''.8$) of northeast, including polarisation.
Wang & Scoville 1992, ApJ, 386, 158. Observations of shocked molecular species.
Turner *et al.* 1992, ApJ, 399, 114. Observations of shocked molecular species.
Dickman *et al.* 1992, ApJ, 400, 203. Observations of shocked molecular species.
van Dishoeck *et al.* 1993, A&A, 279, 541. Observations of shocked molecular species.
Asaoka & Aschenbach 1994, A&A, 284, 573. X-ray, including possible overlapping remnant.
Taubert *et al.* 1994, ApJ, 421, 570. Observations of shocked molecular species.
Esposito *et al.* 1996, ApJ, 461, 820. γ -ray observations.
Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
Keohane *et al.* 1997, ApJ, 484, 350. ASCA observations.
Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.
Claussen *et al.* 1997, ApJ, 522, 349. High resolution observations of OH masers.
Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.
Oliva *et al.* 1999, A&A, 341, L75. ISO observations.
Cesarsky *et al.* 1999, A&A, 348, 945. ISO observations of shocked molecular H₂.
Bocchino & Bykov 2000, A&A, 362, L29. BeppoSAX observations.
Rho *et al.* 2001, ApJ, 547, 885. ISO and 2Mass IR spectroscopy and imaging.
Olbert *et al.* 2001, ApJ, 554, L205. Chandra of compact X-ray source and surrounding nebula.
Bocchino & Bykov 2001, A&A, 376, 248. XMM of compact X-ray source and surrounding nebula.
Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
Kawasaki *et al.* 2002, ApJ, 572, 897. ASCA observations.
Bocchino & Bykov 2003, A&A, 400, 203. XMM observations of compact sources.
Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'.5 : S = 160 \pm 5$ Jy).
Welsh & Sallmen 2003, A&A, 408, 545. Optical absorption studies.
Leahy 2004, AJ, 127, 2277. DRAO at 408 MHz ($3'.3 \times 8'.6$) and 1.4 GHz ($1'.0 \times 2'.6$), for spectral index studies.
see also: Leahy 2004, AJ, 128, 1478. Addendum.
Snell *et al.* 2005, ApJ, 620, 758. Molecular line and IR observations of shocked material.
Bykov *et al.* 2005, ApJ, 624, L41. Chandra observations of compact source.
Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer IR observations.
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
Gaensler *et al.* 2006, ApJ, 648, 1037. Chandra of X-ray source and nebula.
Hewitt *et al.* 2006, ApJ, 652, 1288. GBT at 1.6 and 1.7 GHz ($7'.2$) for OH, and VLA at 330 MHz ($74'' \times 64''$).
Rosado *et al.* 2007, AJ, 133, 89. Observations of shocked H₂.
Albert *et al.* 2007, ApJ, 664, L87. γ -ray observations.
Lee *et al.* 2008, AJ, 135, 796. VLA at 1.4 GHz ($39'' \times 42''$) including HI.
Troja *et al.* 2008, A&A, 485, 777. XMM observations.
Bocchino *et al.* 2008, AdSpR, 41, 396. Integral observations.
Bykov *et al.* 2008, ApJ, 676, 1050. XMM, Chandra and Spitzer observations.

G192.8–1.1		PKS 0607+17
RA: 06 ^h 09 ^m 20 ^s	1-GHz flux/Jy: 20?	Size/arcmin: 78
Dec: +17°20'	Spectral index: 0.6?	Type: S
Has been called G193.3–1.5. Has been regarded as part of the Origem Loop, a supposed larger remnant.		
Radio: In complex region.		
Optical: Encompasses S261 and S254–258.		
References:		
Milne & Dickel 1974, <i>AuJPh</i> , 27, 549. Parkes 64-m at 2.7 GHz ($9' : S = 13 \pm 15\%$ Jy).		
Dickel & DeNoyer 1975, <i>AJ</i> , 80, 437. Arecibo at 318 MHz ($15' : S = 74 \pm 32$ Jy), and $S_{610 \text{ MHz}} = 40$ Jy.		
Caswell 1985, <i>AJ</i> , 90, 1076. DRAO at 1.4 GHz ($1' \times 3'.3 : S = 18 \pm 3$ Jy).		
Biggs & Lyne 1996, <i>MNRAS</i> , 282, 691. Pulsar search.		
Welsh <i>et al.</i> 2001, <i>A&A</i> , 372, 516. Far-UV spectroscopy.		
G205.5+0.5		Monoceros Nebula
RA: 06 ^h 39 ^m 00 ^s	1-GHz flux/Jy: 160	Size/arcmin: 220
Dec: +06°30'	Spectral index: 0.5	Type: S
Radio: In complex region, parts may be HII regions.		
Optical: Large ring, near Rosette nebula.		
X-ray: Possibly detected.		
Distance: Mean optical velocity suggests 0.8 kpc, low frequency radio absorption suggests 1.6 kpc.		
References:		
Milne & Dickel 1974, <i>AuJPh</i> , 27, 549. Parkes 64-m at 2.7 GHz ($9'$).		
Velusamy & Kundu 1974, <i>A&A</i> , 32, 375. NRAO 300-ft at 2.7 GHz ($5'$), part only.		
Dickel & DeNoyer 1975, <i>AJ</i> , 80, 437. Arecibo at 111 MHz ($1' : S = 462 \pm 180$ Jy) and $S_{610 \text{ MHz}} = 245$ Jy.		
Davies <i>et al.</i> 1978, <i>A&AS</i> , 31, 271. Deep optical plates.		
Lozinskaya 1981, <i>SvAL</i> , 7, 17. Mean optical velocity.		
Graham <i>et al.</i> 1982, <i>A&A</i> , 109, 145. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 97.6 \pm 12.5$ Jy), plus review of flux densities.		
Fesen <i>et al.</i> 1985, <i>ApJ</i> , 292, 29. Optical spectra.		
Leahy <i>et al.</i> 1986, <i>MNRAS</i> , 220, 501. Einstein observations.		
Odegard 1986, <i>ApJ</i> , 301, 813. TPT at 20.6, 25.6 and 30.9 MHz ($24', 19'$ and $16'$).		
Esposito <i>et al.</i> 1996, <i>ApJ</i> , 461, 820. Possible associated γ -ray emission.		
Biggs & Lyne 1996, <i>MNRAS</i> , 282, 691. Pulsar search.		
Oliver <i>et al.</i> 1996, <i>A&A</i> , 315, 578. CO observations of some of surroundings.		
Jaffe <i>et al.</i> 1997, <i>ApJ</i> , 484, L129. γ -ray detection.		
Kaplan <i>et al.</i> 2006, <i>ApJS</i> , 163, 344. X-ray upper limit on compact sources.		
Casandjian & Grenier 2008, <i>A&A</i> , 489, 849. γ -ray observations.		
G206.9+2.3		PKS 0646+06
RA: 06 ^h 48 ^m 40 ^s	1-GHz flux/Jy: 6	Size/arcmin: 60×40
Dec: +06°26'	Spectral index: 0.5	Type: S?
Radio: Diffuse source near the Monoceros Nebula.		
Optical: Filaments detected.		
X-ray: Possibly detected.		
References:		
Davies & Meaburn 1978, <i>A&A</i> , 69, 443. Optical observations.		
Nousek <i>et al.</i> 1981, <i>ApJ</i> , 248, 152. HEAO-1 X-ray limit.		
Graham <i>et al.</i> 1982, <i>A&A</i> , 109, 145. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 4.1 \pm 0.6$ Jy), plus review of flux densities.		
Rosado 1982, <i>RMxAA</i> , 5, 127. Optical observations.		
Fesen <i>et al.</i> 1985, <i>ApJ</i> , 292, 29. Optical spectra.		
Leahy <i>et al.</i> 1986, <i>MNRAS</i> , 220, 501. Einstein observations.		
Odegard 1986, <i>ApJ</i> , 301, 813. TPT at 20.6, 25.6 and 30.9 MHz ($24', 19'$ and $16'$).		
Biggs & Lyne 1996, <i>MNRAS</i> , 282, 691. Pulsar search.		
Aharonian <i>et al.</i> 2004, <i>A&A</i> , 417, 973. γ -ray limits.		

G260.4–3.4

RA: 08^h22^m10^s
Dec: −43°00'

1-GHz flux/Jy: 130
Spectral index: 0.5

Puppis A, MSH 08–44
Size/arcmin: 60 × 50
Type: S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Angular shell, brightest to the E, poorly defined to the W.

Optical: Nebulosity and wisps.

X-ray: Brightest to the E.

Point sources: Central possible pulsating X-ray source.

Distance: Association with HI gives 2.2 kpc.

References:

- Green 1971, AuJPh, 24, 773. Molonglo at 408 MHz ($3':S=198\pm20$ Jy).
 Goudis & Meaburn 1978, A&A, 62, 283. H α +[NII] optical image.
 Petre *et al.* 1982, ApJ, 258, 22. Einstein observations.
 Milne *et al.* 1983, MNRAS, 204, 237. FIRST at 1415 MHz ($50''$), large scale emission missing.
 Teske & Petre 1987, ApJ, 318, 370. Coronal optical line emission.
 Dubner & Arnal 1988, A&AS, 75, 363. HI and CO observations of surroundings.
 Arendt *et al.* 1990, ApJ, 350, 266. MOST at 843 MHz ($44''\times65''$), with large scale emission added, plus IR, optical and X-ray observations.
 Dubner *et al.* 1991, AJ, 101, 1466. VLA at 327 MHz ($43''\times73''$) and 1.5 GHz ($43''\times77''$).
 Arendt *et al.* 1991, ApJ, 368, 474. IR observations.
 Milne *et al.* 1993, MNRAS, 261, 366. Parkes 64-m at 4.75 ($4.5':S=59\pm5$ Jy) and 8.4 GHz ($3':S=38\pm4$ Jy), plus polarisation.
 Berthiaume *et al.* 1994, ApJ, 425, 132. X-ray spectroscopy.
 Sutherland & Dopita 1995, ApJ, 439, 365. Spectrophotometry.
 Reynoso *et al.* 1995, AJ, 110, 318. VLA at 1.4 GHz ($90''$) including neutral hydrogen.
 Blair *et al.* 1995, ApJ, 454, L35. Far UV spectroscopy.
 Petre *et al.* 1996, ApJ, 465, L43. ROSAT of central source.
 Bock *et al.* 1998, AJ, 116, 1886. MOST at 843 MHz ($43''\times60''$).
 Pavlov *et al.* 1999, ApJ, 511, L45. Possible pulsation detection from central X-ray source.
 Zavlin *et al.* 1999, ApJ, 525, 959. X-ray observations of central source.
 Bocchino *et al.* 2000, A&A, 359, 316. Optical studies of selected filaments in N.
 Woermann *et al.* 2000, MNRAS, 317, 421. OH observations.
 Gaensler *et al.* 2000, ApJ, 537, L35. Radio limit for nebula around possible pulsar.
 Reynoso *et al.* 2003, MNRAS, 345, 671. ATCA at 1.4 GHz ($90''$) for HI near central X-ray source.
 Hwang *et al.* 2005, ApJ, 635, 355. Chandra observations of E edge.
 Hui & Becker 2006, A&A, 454, 543. XMM and Chandra observations.
 Hui & Becker 2006, A&A, 457, L33. Chandra proper motion study of central source.
 Castelletti *et al.* 2006, A&A, 459, 535. VLA at 1.4 GHz ($16''\times34''$) and 327 MHz ($45''\times90''$) ($S=114\pm8$ Jy).
 Winkler & Petre 2007, ApJ, 670, 635. Chandra proper motion study of central source.
 Paron *et al.* 2008, A&A, 480, 439. CO observations of E.
 Hwang *et al.* 2008, ApJ, 676, 378. Suzaku observations.

G261.9+5.5

RA: 09^h04^m20^s
Dec: −38°42'

1-GHz flux/Jy: 10?
Spectral index: 0.4?

Size/arcmin: 40 × 30
Type: S

Radio: Faint shell with little limb brightening.

References:

- Hill 1967, AuJPh, 20, 297. Parkes 64-m at 2650 MHz ($7.5':S=7$ Jy) also $S_{1410\text{ MHz}}=8$ Jy, $S_{81.5\text{ MHz}}=25$ Jy.
 Colomb & Dubner 1980, A&A, 82, 244. Argentine 30-m dish at 1.4 GHz, for HI possibly associated with remnant.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44''\times71''$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G263.9–3.3		Vela (XYZ)
RA: 08 ^h 34 ^m 00 ^s	1-GHz flux/Jy: 1750	Size/arcmin: 255
Dec: −45°50'	Spectral index: varies	Type: C
This refers to the whole Vela XYZ complex, of which X has at times been classified as a separate (filled-centre) remnant. This remnant is overlapped by G260.4–3.4 and G266.2–1.2.		
Radio: Large shell, with flatter spectrum component (Vela X), and pulsar nebula.		
Optical: Filaments.		
X-ray: Patchy shell, with extensions, central nebula and pulsar.		
Point sources: Pulsar within Vela X, with one-sided ‘jet’.		
Distance: Vela pulsar parallax gives 0.3 kpc, optical spectra and HI studies suggest 0.25 kpc.		
References:		
Milne 1968, <i>AuJPh</i> , 21, 201. Parkes 64-m at various frequencies, including 408 MHz ($48' : S = 2300 \pm 300$ Jy), 635 MHz ($31' : S = 2360 \pm 300$ Jy), 1410 MHz ($14' : S = 1640 \pm 300$ Jy) and 2650 MHz ($7.5' : S = 1400 \pm 250$ Jy), plus discussion of the distance.		
Milne 1980, <i>A&A</i> , 81, 293. Maps of Vela X with Parkes 64-m at 1.66, 2.7 and 5 GHz ($12', 8.4$ and $4.4'$).		
Weiler & Panagia 1980, <i>A&A</i> , 90, 269. Clarification of notation of this region and review previous observations.		
Bignami & Caraveo 1988, <i>ApJ</i> , 325, L5. Pulsar proper motion from optical observations.		
Dwarakanath 1991, <i>JApA</i> , 12, 199. Gauribidanur ‘T’ array at 34.5 MHz ($26' \times 84' : S = 1800$ Jy for Vela X and $S = 3900$ Jy for Vela YZ), plus review of flux densities.		
Bietenholz <i>et al.</i> 1991, <i>ApJ</i> , 376, L41. VLA at 5 GHz in vicinity of pulsar.		
Willmore <i>et al.</i> 1992, <i>MNRAS</i> , 254, 139. Hard X-ray observations.		
Dubner <i>et al.</i> 1992, <i>A&AS</i> , 96, 505. Argentine 30-m at 1.4 GHz ($30' : S = 1800$ Jy) of surrounding HI.		
Strom <i>et al.</i> 1995, <i>Nature</i> , 373, 590. Radio of X-ray extensions.		
Markwardt & Ögelman 1995, <i>Nature</i> , 375, 40. X-ray jet from pulsar.		
Jenkins & Wallerstein 1995, <i>ApJ</i> , 440, 227. Optical absorption of associated neutral carbon cloud.		
Milne 1995, <i>MNRAS</i> , 277, 1435. Parkes 64-m at 8.4 GHz ($3' : S = 1800$ Jy), including polarisation, of Vela X.		
Danks & Sembach 1995, <i>AJ</i> , 109, 2627. Optical spectroscopy of background stars.		
Blair <i>et al.</i> 1995, <i>AJ</i> , 110, 312. UV spectroscopy.		
Duncan <i>et al.</i> 1996, <i>MNRAS</i> , 280, 252. Parkes 64-m at 2.4 GHz ($8.9' : S = 1800$ Jy).		
Whiteoak & Green 1996, <i>A&AS</i> , 118, 329. MOST at 843 MHz ($43'' \times 62''$) of part.		
Green <i>et al.</i> 1997, <i>AJ</i> , 114, 2058. Parkes 64-m and ATCA OH observations.		
Frail <i>et al.</i> 1997, <i>ApJ</i> , 475, 224. VLA at 330 MHz ($1' \times 1.8'$) and comparison with ROSAT observations of Vela X.		
Markwardt & Ögelman 1997, <i>ApJ</i> , 480, L13. ASCA observations of pulsar ‘jet’.		
Raymond <i>et al.</i> 1997, <i>ApJ</i> , 482, 881. UV spectroscopy.		
Yoshikoshi <i>et al.</i> 1997, <i>ApJ</i> , 487, L65. γ-ray observations of pulsar.		
Jenkins <i>et al.</i> 1998, <i>ApJ</i> , 492, L147. UV absorption to background star.		
Dubner <i>et al.</i> 1998, <i>AJ</i> , 116, 813. Parkes 64-m at 1.4 GHz ($15' : S = 1800$ Jy) for HI studies.		
Bock <i>et al.</i> 1998, <i>AJ</i> , 116, 1886. MOST at 843 MHz ($43'' \times 60''$).		
Cha <i>et al.</i> 1999, <i>ApJ</i> , 515, L25. Optical spectra, for distance.		
Lu & Aschenbach 2000, <i>A&A</i> , 362, 1083. ROSAT spatially resolved spectroscopy.		
Chadwick <i>et al.</i> 2000, <i>ApJ</i> , 537, 414. Limit on high energy γ-rays from pulsar.		
Sankrit <i>et al.</i> 2001, <i>ApJ</i> , 549, 416. Far-UV observations of selected region.		
Pavlov <i>et al.</i> 2001, <i>ApJ</i> , 554, L189. Chandra two-epoch observations of pulsar nebula.		
Helfand <i>et al.</i> 2001, <i>ApJ</i> , 556, 380. Chandra observations of pulsar and its nebula.		
Miyata <i>et al.</i> 2001, <i>ApJ</i> , 559, L45. Chandra of X-ray extension.		
Alvarez <i>et al.</i> 2001, <i>A&A</i> , 372, 636. Radio spectral index studies.		
Caraveo <i>et al.</i> 2001, <i>ApJ</i> , 561, 930. HST parallax observations of pulsar.		
Moriguchi <i>et al.</i> 2001, <i>PASJ</i> , 53, 1025. CO observations.		
Dodson <i>et al.</i> 2003, <i>MNRAS</i> , 343, 116. ATCA at 1.4, 2.4, 5.2 and 8.5 GHz ($8.1' \times 6.2', 36'' \times 26'', 12.1' \times 10.5'$ and $11.2' \times 10.6'$) of pulsar nebula.		
Sankrit <i>et al.</i> 2003, <i>ApJ</i> , 589, 242. Optical nebulosity to NE of G266.2–1.2.		
Nichols & Slavin 2004, <i>ApJ</i> , 610, 285. UV absorption toward background sources.		
Hales <i>et al.</i> 2004, <i>ApJ</i> , 613, 977. Vela X at 31 GHz ($4.1'$).		
Mongano <i>et al.</i> 2005, <i>A&A</i> , 436, 917. XMM and other X-ray observations of pulsar nebula.		
Miceli <i>et al.</i> 2005, <i>A&A</i> , 442, 513. XMM observations of N rim.		
Katsuda & Tsunemi 2005, <i>PASJ</i> , 57, 621. XMM observations of E.		
Aharonian <i>et al.</i> 2006, <i>A&A</i> , 448, L43. γ-ray observations.		
McConnell <i>et al.</i> 2006, <i>AJ</i> , 131, 648. ATCA at 4.9 GHz ($12' : S = 1800$ Jy) including polarisation.		
Katsuda & Tsunemi 2006, <i>ApJ</i> , 642, 917. XMM observations of NE.		
Nishikida <i>et al.</i> 2006, <i>ApJ</i> , 644, L171. Far UV observations.		
Miceli <i>et al.</i> 2008, <i>ApJ</i> , 676, 1064. XMM observations.		
LaMassa <i>et al.</i> 2008, <i>ApJ</i> , 689, L121. XMM observations of Vela X.		

G266.2–1.2

RA: 08^h52^m00^s
Dec: −46°20'

1-GHz flux/Jy: 50?
Spectral index: 0.3?

RX J0852.0–4622
Size/arcmin: 120
Type: S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Incomplete shell, confused by the Vela SNR.

Optical: Nebulosity offset to NE.

X-ray: Non-thermal shell, confused by the Vela SNR, with central source, and possible associated pulsar.

Point sources: Central X-ray source, with optical nebula, and possible associated pulsar.

Distance: X-ray data suggest an upper limit of 1 kpc.

References:

- Aschenbach 1998, Nature, 396, 141. ROSAT identification.
 Iyudin *et al.* 1998, Nature, 396, 142. γ-ray observations.
 Combi *et al.* 1999, ApJ, 519, L177. Radio observations.
 Duncan & Green 2000, A&A, 364, 732. Parkes 64-m at 1.4 GHz (14'0), and comparison with other observations.
 Redman *et al.* 2000, ApJ, 543, L153. Optical of nearly nebulosity.
 Tsunemi *et al.* 2000, PASJ, 52, 887. ASCA spectral observations.
 Slane *et al.* 2001, ApJ, 548, 814. ASCA observations.
 Mereghetti *et al.* 2001, ApJ, 548, L213. BeppoSAX observations of central sources.
 Pavlov *et al.* 2001, ApJ, 559, L131. Chandra of central X-ray source.
 Moriguchi *et al.* 2001, PASJ, 53, 1025. CO observations.
 Pellizzoni *et al.* 2002, A&A, 393, L65. Optical observations of central source.
 Redman *et al.* 2002, MNRAS, 336, 1093. Optical nebulosity to NE.
 Kargalstev *et al.* 2002, ApJ, 580, 1060. Chandra observations of central source.
 Sankrit *et al.* 2003, ApJ, 589, 242. Optical nebulosity to NE.
 Redman & Meaburn 2005, MNRAS, 356, 969. Possible pulsar association.
 Iyudin *et al.* 2005, A&A, 429, 225. XMM observations.
 Aharonian *et al.* 2005, A&A, 437, L7. γ-ray observations.
 Katagiri *et al.* 2005, ApJ, 619, L163. γ-ray observations.
 Bamba *et al.* 2005, ApJ, 632, 294. Chandra of NW rim.
 Reynolds *et al.* 2006, A&A, 449, 243. ATCA at 1.38 GHz (32'' × 37'').
 Enomoto *et al.* 2006, ApJ, 652, 1268. γ-ray observations.
 Mignami *et al.* 2007, A&A, 473, 883. Deep optical observations of compact X-ray source.
 Aharonian *et al.* 2007, ApJ, 661, 236. γ-ray observations.
 Katsuda *et al.* 2008, ApJ, 678, L35. XMM proper motion study.
 Katsuda *et al.* 2008, ApJ, 678, 297. XMM observations.

G272.2–3.2

RA: 09^h06^m50^s
Dec: −52°07'

1-GHz flux/Jy: 0.4
Spectral index: 0.6

Size/arcmin: 15?
Type: S?

Radio: Diffuse shell.

X-ray: Centrally brightened.

Optical: Detected.

References:

- Greiner *et al.* 1994, A&A, 286, L35. ROSAT observations, plus optical observations.
 Duncan *et al.* 1997, MNRAS, 289, 97. Parkes 64-m at 1.4 GHz (18' : $S = 0.38 \pm 0.09$ Jy) 2.4 GHz (10'6 : $S = 0.25 \pm 0.04$ Jy) and 4.8 GHz (5'7 : $S = 0.17 \pm 0.02$ Jy), MOST at 843 MHz (45'' × 70'' : $S = 0.45 \pm 0.10$ Jy), and ATCA at 2.4 GHz (37'' × 52''), plus ROSAT observations.
 Harris *et al.* 2001, ApJ, 552, 614. ASCA and ROSAT observations, plus review of earlier observations.

G279.0+1.1

RA: 09^h57^m40^s
Dec: -53°15'

1-GHz flux/Jy: 30?
Spectral index: 0.6?

Size/arcmin: 95
Type: S

Radio: Faint, incomplete shell.

Point sources: Pulsar nearby.

References:

Woermann & Jonas 1988, MNRAS, 234, 971. Hartesbeesthoek 26-m at 1.6 (30' : $S=25.2 \pm 4$ Jy) and 2.3 GHz (20' : $S=20.7 \pm 3$ Jy).
Duncan *et al.* 1995, MNRAS, 277, 319. Parkes 64-m at 1.4 (18' : $S=28 \pm 3$ Jy) and 2.4 GHz (11' : $S=20 \pm 2$ Jy), including polarisation.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 53'') of part.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G284.3-1.8

RA: 10^h18^m15^s
Dec: -59°00'

1-GHz flux/Jy: 11?
Spectral index: 0.3?

MSH 10-53
Size/arcmin: 24?
Type: S

Has been called G284.2-1.8.

Radio: Incomplete, poorly defined shell.

Point sources: Pulsar with wind nebula nearby.

References:

Ruiz & May 1986, ApJ, 309, 667. CO and optical observations.
Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43'' × 50'') and Parkes 64-m at 8.4 GHz (3' : $S=5.4 \pm 0.8$ Jy) including polarisation, plus earlier flux densities.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
Camilo *et al.* 2001, ApJ, 557, L51. Observations of nearby pulsar.
Camilo *et al.* 2004, ApJ, 616, 1118. Chandra observations of pulsar and nebula.

G286.5-1.2

RA: 10^h35^m40^s
Dec: -59°42'

1-GHz flux/Jy: 1.4?
Spectral index: ?

Size/arcmin: 26 × 6
Type: S?

Radio: Double, elongated arc.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 50'') : $S=1.6$ Jy.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G289.7-0.3

RA: 11^h01^m15^s
Dec: -60°18'

1-GHz flux/Jy: 6.2
Spectral index: 0.2?

Size/arcmin: 18 × 14
Type: S

Radio: Incomplete shell.

Point sources: Compact radio source near centre.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 50'') : $S=6.4 \pm 0.5$ Jy), plus Parkes 64-m at 4.5 GHz ($S=7.5 \pm 2.5$ Jy) and 8.55 GHz ($S=3.6 \pm 0.9$ Jy).

G290.1–0.8		MSH 11–61A
RA: 11 ^h 03 ^m 05 ^s	1-GHz flux/Jy: 42	Size/arcmin: 19 × 14
Dec: −60°56'	Spectral index: 0.4	Type: S
Radio: Elongated, clumpy shell.		
Optical: Filaments detected.		
X-ray: Centrally brightened.		
Point sources: Pulsar nearby.		
Distance: HI absorption indicates 7 ± 1 kpc.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4 : $S = 20.2$ Jy).		
Elliott & Malin 1979, MNRAS, 186, 45P. Optical image and spectra.		
Kirshner & Winkler 1979, ApJ, 227, 853. Optical observations.		
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 50'').		
Milne <i>et al.</i> 1989, PASA, 8, 187. MOST at 843 MHz (43'' × 49'': $S = 45 \pm 11$ Jy), and Parkes 64-m at 8.4 GHz (3' : $S = 19.5 \pm 1.0$ Jy), including polarisation.		
Seward 1990, ApJS, 73, 781. Einstein observations.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'': $S = 43$ Jy).		
Rosado <i>et al.</i> 1996, A&A, 315, 243. Optical kinematics.		
Kaspi <i>et al.</i> 1997, ApJ, 485, 820. Pulsar detection.		
Gotthelf & Kaspi 1998, ApJ, 497, L29. ASCA observations of pulsar.		
Slane <i>et al.</i> 2002, ApJ, 564, 284. ASCA observations.		
Filipović <i>et al.</i> 2005, SerAJ, 170, 47. ATCA at 1.4 GHz (21''), plus other observations, including CO of surroundings.		
Reynoso <i>et al.</i> 2006, MNRAS, 369, 416. ATCA at 1.4 GHz (22''.5 × 25''.0) including HI.		

G291.0–0.1		(MSH 11–62)
RA: 11 ^h 11 ^m 54 ^s	1-GHz flux/Jy: 16	Size/arcmin: 15 × 13
Dec: −60°38'	Spectral index: 0.29	Type: C
Radio: Centrally brightened core, with surrounding arcs.		
X-ray: Centrally brightened.		
Point sources: Central compact X-ray source.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Roger <i>et al.</i> 1986, MNRAS, 219, 815. MOST at 843 MHz (43'' × 50'': $S = 17.2 \pm 1.0$ Jy), and Parkes 64-m at 5 and 8.4 GHz (4'.6 and 3' : $S = 10.4 \pm 0.4$ Jy and 9'.1 ± 0.2), with polarisation.		
Wilson 1986, ApJ, 302, 718. Einstein observations.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'': $S = 12.7$ Jy).		
Harrus <i>et al.</i> 1998, ApJ, 499, 273. ASCA observations.		

G292.0+1.8		MSH 11–54
RA: 11 ^h 24 ^m 36 ^s	1-GHz flux/Jy: 15	Size/arcmin: 12 × 8
Dec: −59°16'	Spectral index: 0.4	Type: C

Radio: Centrally brightened source surrounded by a plateau of faint emission.

Optical: Oxygen rich.

X-ray: Ring of emission, with diffuse central nebula and pulsar.

Point sources: Central pulsar.

Distance: HI absorption implies 6.0 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Goss *et al.* 1979, MNRAS, 188, 357. Optical spectra.
 Lockhart *et al.* 1977, MNRAS, 179, 147. Fleurs at 1415 MHz (50'': $S=13.0$ Jy).
 Clark *et al.* 1980, MNRAS, 193, 129. X-ray spectrum.
 Tuohy *et al.* 1982, ApJ, 260, L65. Einstein observations.
 Dopita & Tuohy 1984, ApJ, 282, 135. Optical spectra.
 Braun *et al.* 1986, A&A, 162, 259. MOST at 843 MHz (0.8') and IRAS.
 Hughes & Singh 1994, ApJ, 422, 126. EXOSAT spectrum.
 Sutherland & Dopita 1995, ApJ, 439, 365. Spectrophotometry.
 Hughes *et al.* 2001, ApJ, 559, L153. Chandra observations, including central nebula.
 Camilo *et al.* 2002, ApJ, 567, L71. Pulsar detection.
 Park *et al.* 2002, ApJ, 564, L39. Chandra observations.
 Gonzalez & Safi-Harb 2003, ApJ, 583, L91. Chandra observations.
 Hughes *et al.* 2003, ApJ, 591, L139. Chandra observations of pulsar.
 Gaensler & Wallace 2003, ApJ, 594, 326. ATCA at 1.4, 2.3 and 5.2 GHz (9''.6 × 8''.0, 7''.2 × 6''.2 and 5''.5 × 4''.8 : $S=11.9 \pm 0.1$, 11.4 ± 0.1 and 8.8 ± 0.1), plus HI observations.
 Park *et al.* 2004, ApJ, 602, L33. Chandra observations.
 Vink *et al.* 2004, NuPhS, 132, 62. XMM observations.
 Ghavamian *et al.* 2005, ApJ, 635, 365. Optical imaging spectroscopy.
 Winkler & Long 2006, AJ, 132, 360. Optical observations.
 Park *et al.* 2007, ApJ, 670, L121. Chandra observations.
 Zharikov *et al.* 2008, A&A, 492, 805. Possible optical counterpart to pulsar and nebula.

G292.2–0.5

RA: 11 ^h 19 ^m 20 ^s	1-GHz flux/Jy: 7	Size/arcmin: 20 × 15
Dec: −61°28'	Spectral index: 0.5	Type: S

Radio: Shell.

X-ray: Detected.

Point sources: Central, young pulsar with X-ray nebula.

Distance: HI absorption indicates 8.4 kpc.

References:

- Camilo *et al.* 2000, ApJ, 541, 367. Pulsar detection.
 Crawford *et al.* 2001, ApJ, 554, 152. ATCA at 1.4 GHz (29'' × 25'' : $S=5.6 \times 0.3$ Jy) and 2.5 GHz (21'' × 20'').
 Pivovaroff *et al.* 2001, ApJ, 554, 161. ROSAT and ASCA observations.
 Gonzalez & Safi-Harb 2003, ApJ, 591, L143. Chandra observations of pulsar.
 Caswell *et al.* 2004, MNRAS, 352, 1405. ATCA at 5 GHz (1' : $S=2.8$ Jy), including polarisation, and 1.4 GHz for HI absorption.
 Gonzalez & Safi-Harb 2005, ApJ, 619, 856. Chandra observations.
 Gonzalez *et al.* 2005, ApJ, 630, 489. XMM observations of pulsar.
 Safi-Harb *et al.* 2008, ApJ, 684, 532. Chandra observations of pulsar and nebula.

G293.8+0.6

RA: 11^h35^m00^s
Dec: -60°54'

1-GHz flux/Jy: 5?
Spectral index: 0.6?

Size/arcmin: 20
Type: C

Radio: Central source, with faint extended plateau.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 9.0$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 2.1$ Jy).
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 51''$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 49'' : S = 2.6$ Jy).

G294.1-0.0

RA: 11^h36^m10^s
Dec: -61°38'

1-GHz flux/Jy: >2?
Spectral index: ?

Size/arcmin: 40
Type: S

Radio: Faint shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 49'' : S > 2$).

G296.1-0.5

RA: 11^h51^m10^s
Dec: -62°34'

1-GHz flux/Jy: 8?
Spectral index: 0.6?

Size/arcmin: 37×25
Type: S

Incorporates the previously catalogued remnant G296.1-0.7. Has been called G296.05-0.50.

Radio: Irregular shell, with nearby HII regions.

Optical: Detected.

X-ray: Detected.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 6.9$ Jy) and Parkes 64-m at 5 GHz ($4' : S > 0.74$).
 Longmore *et al.* 1977, MNRAS, 181, 541. Optical spectra.
 van den Berg 1978, ApJS, 38, 119. Optical observations.
 Markert *et al.* 1981, ApJ, 248, L17. Einstein observations.
 Caswell & Barnes 1983, ApJ, 271, L55. Molonglo at 408 MHz ($3' : S = 12.4$ Jy).
 Bignami *et al.* 1986, ApJ, 302, 606. EXOSAT and Einstein observations.
 Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 48'' : S > 2.4$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G296.5+10.0

RA: 12^h09^m40^s
Dec: −52°25'

1-GHz flux/Jy: 48
Spectral index: 0.5

PKS 1209–51/52

Size/arcmin: 90 × 65
Type: S

Has been called G296.5+9.7.

Radio: Shell with two bright limbs.

Optical: Detected.

X-ray: Incomplete shell, with central pulsar.

Point sources: Central pulsar.

References:

- Irvine & Irvine 1974, ApJ, 192, L111. Optical observations.
Danziger & Dennefeld 1976, PASP, 88, 44. Optical spectra.
Dicke & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
Tuohy *et al.* 1979, ApJ, 230, L27. X-ray detection with HEAO-1 A2 experiment.
Ruiz 1983, AJ, 88, 1210. Optical spectra.
Dubner *et al.* 1986, AJ, 91, 343. Argentine 30-m dish at 1.4 GHz (34'), plus H_i.
Kellett *et al.* 1987, MNRAS, 225, 199. EXOSAT of the west of the remnant, including the compact source.
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 56'').
Matsui *et al.* 1988, ApJ, 329, 838. Einstein observations, including compact source.
Roger *et al.* 1988, ApJ, 332, 940. MOST at 843 MHz (44'' × 56'').
Bignami *et al.* 1992, ApJ, 389, L67. Optical in vicinity of X-ray source.
Milne & Haynes 1994, MNRAS, 270, 106. Parkes 64-m at 2.4 GHz (8'3 : S = 33 ± 3 Jy), 4.8 GHz (4'5 : S = 23.3 ± 3 Jy) and 8.4 GHz (3'0 : 18.8 ± 3), including polarisation and review of flux densities.
Mereghetti *et al.* 1996, ApJ, 464, 842. Radio, optical and X-ray observations of central source.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
Vasisht *et al.* 1997, ApJ, 476, L43. ASCA observations of neutron star.
Zavlin *et al.* 1998, A&A, 331, 821. ROSAT and ASCA observations of neutron star.
Giacani *et al.* 2000, AJ, 119, 281. ATCA at 1.4 GHz (2'7 × 4'0) for H_i studies.
Zavlin *et al.* 2000, ApJ, 540, L25. Chandra observations of central pulsar.
Gothelf & Halpern 2007, ApJ, 664, L35. X-ray timing observations of pulsar.

G296.8–0.3

RA: 11^h58^m30^s
Dec: −62°35'

1-GHz flux/Jy: 9
Spectral index: 0.6

1156–62

Size/arcmin: 20 × 14
Type: S

Radio: Shell, brighter to the NW.

Distance: H_i absorption gives 9.6 kpc.

X-ray: Detected.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : S = 15.0 Jy) and Parkes 64-m at 5 GHz (4' : S = 3.2 Jy).
Dicke & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'' : S = 9.2 Jy).
Gaensler *et al.* 1998, MNRAS, 296, 813. ATCA at 1.3 GHz (22'' × 24'' : S = 7.0 ± 0.3 Jy), including polarisation and H_i observations, plus review of flux densities.

G298.5–0.3

RA: 12^h12^m40^s
Dec: −62°52'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 5?
Type: ?

Radio: Not well resolved, may be part of a larger ring?

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'' : S = 1.8 Jy).
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G298.6–0.0

RA: 12^h13^m41^s
Dec: −62°37'

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 12 × 9
Type: S

Has been called G298.6–0.1.

Radio: Incomplete shell, in complex region.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 50'').
Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'': $S=7.4$ Jy).
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G299.2–2.9

RA: 12^h15^m13^s
Dec: −65°30'

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 18 × 11
Type: S

Radio: Faint source.

X-ray: Centrally brightened with shell at higher energies.

Optical: Filaments in W.

References:

Busser *et al.* 1996, A&A, 310, L1. ROSAT detection, plus optical studies.
Slane *et al.* 1996, ApJ, 465, 840. Einstein, IRAS and radio observations.
Bai & Wang 2000, ApJ, 539, 760. ASCA observations.
Park *et al.* 2007, ApJ, 665, 1173. Chandra observations.

G299.6–0.5

RA: 12^h21^m45^s
Dec: −63°09'

1-GHz flux/Jy: 1.0?
Spectral index: ?

Size/arcmin: 13
Type: S

Radio: Faint shell, brightest to E.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'': $S=1.1$ Jy).

G301.4–1.0

RA: 12^h37^m55^s
Dec: −63°49'

1-GHz flux/Jy: 2.1?
Spectral index: ?

Size/arcmin: 37 × 23
Type: S

Radio: Faint, incomplete shell, with possible extension to southwest.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'': $S=2.3$ Jy).

G302.3+0.7

RA: 12^h45^m55^s
Dec: −62°08'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 17
Type: S

Radio: Distorted shell, in complex region, with possibly associated filament.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=7.5$ Jy) and Parkes 64-m at 5 GHz (4': $S=3.0$ Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'': $S=3.2$ Jy).
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

G304.6+0.1			Kes 17
RA: 13 ^h 05 ^m 59 ^s	1-GHz flux/Jy: 14	Size/arcmin: 8	
Dec: -62°42'			
References: Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4'). Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4.4'; S=6.9 Jy). Caswell <i>et al.</i> 1975, A&A, 45, 239. Parkes HI absorption. Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'': S=18 Jy). Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant. Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer observations.			
<hr/>			
G308.1-0.7			
RA: 13 ^h 37 ^m 37 ^s	1-GHz flux/Jy: 1.2?	Size/arcmin: 13	
Dec: -63°04'			
References: Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'': S=1.3 Jy).			
<hr/>			
G308.8-0.1			
RA: 13 ^h 42 ^m 30 ^s	1-GHz flux/Jy: 15?	Size/arcmin: 30 × 20?	
Dec: -62°23'			
References: Clark <i>et al.</i> 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S=16.7 Jy) and Parkes 64-m at 5 GHz (4': S=7.0 Jy). Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4.4'; S=6.5 Jy). Caswell <i>et al.</i> 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50''). Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein. Caswell <i>et al.</i> 1992, ApJ, 399, L151. MOST at 843 MHz (43'' × 49''). Kaspi <i>et al.</i> 1992, ApJ, 399, L155. Pulsar observations. Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations. Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer possible detection.			
<hr/>			
G309.2-0.6			
RA: 13 ^h 46 ^m 31 ^s	1-GHz flux/Jy: 7?	Size/arcmin: 15 × 12	
Dec: -62°54'			
References: Clark <i>et al.</i> 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S=10.0 Jy) and Parkes 64-m at 5 GHz (4': S=3.9 Jy). Caswell <i>et al.</i> 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1.5'). Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 50''). Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 48'': S=6 Jy). Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations. Gaensler <i>et al.</i> 1998, MNRAS, 299, 812. ATCA at 1.3 GHz (24'' × 23'': S=5.2 ± 0.2 Jy). Rakowski <i>et al.</i> 2001, ApJ, 548, 258. ASCA and ROSAT observations. Rakowski <i>et al.</i> 2006, ApJ, 649, L111. Observations of central source. Safi-Harb <i>et al.</i> 2007, ApJ, 659, 407. Observations of central source.			

G309.8+0.0

RA: 13^h50^m30^s
Dec: −62°05'

1-GHz flux/Jy: 17
Spectral index: 0.5

Size/arcmin: 25 × 19
Type: S

Radio: Distorted shell.

Point sources: Steep radio spectrum source near the centre of the remnant.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=26.4$ Jy) and Parkes 64-m at 5 GHz (4' : $S=7.4$ Jy).
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (1').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'' : $S>8.8$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G310.6−0.3

RA: 13^h58^m00^s
Dec: −62°09'

1-GHz flux/Jy: 5?
Spectral index: ?

Kes 20B
Size/arcmin: 8
Type: S

Radio: Asymmetric shell.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'' : $S=5.4$ Jy).
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G310.8−0.4

RA: 14^h00^m00^s
Dec: −62°17'

1-GHz flux/Jy: 6?
Spectral index: ?

Kes 20A
Size/arcmin: 12
Type: S

Radio: Arc in E, in complex region.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'' : $S=6.9$ Jy).
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

G311.5−0.3

RA: 14^h05^m38^s
Dec: −61°58'

1-GHz flux/Jy: 3?
Spectral index: 0.5

Size/arcmin: 5
Type: S

Radio: Shell, not well resolved.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'' : $S=2.9$ Jy).
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

G312.4–0.4

RA: 14^h13^m00^s
Dec: −61°44'

1-GHz flux/Jy: 45
Spectral index: 0.36

Size/arcmin: 38
Type: S

Radio: Irregular, incomplete shell.

Point sources: Nearby γ-ray sources and pulsars.

X-ray: Weak emission in W.

Distance: HI absorption suggests > 6 kpc and possibly > 14 kpc.

References:

- Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3' : $S = 56$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 49'' : $S > 19$), plus Parkes 64-m at 4.5 GHz ($S = 30 \pm 2$ Jy) and 8.55 GHz ($S = 17 \pm 4$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Case & Bhattacharya 1998, ApJ, 521, 246. Nearby γ-ray sources.
 Roberts *et al.* 1999, ApJ, 515, 712. MOST at 843 MHz (43'' × 49'').
 Doherty *et al.* 2003, MNRAS, 339, 1048. ATCA at 1.4 GHz (25'') plus HI absorption, and Chandra observations.

G312.5–3.0

RA: 14^h21^m00^s
Dec: −64°12'

1-GHz flux/Jy: 3.5?
Spectral index: ?

Size/arcmin: 20 × 18
Type: S

Radio: Distorted shell.

References:

- Kane & Vaughan 2003, MNRAS, 344, 625. ATCA at 1.4 GHz (129'' × 116'') and 2.4 GHz (75'' × 67'').

G315.1+2.7

RA: 14^h24^m30^s
Dec: −57°50'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 190 × 150
Type: S

Radio: Poorly defined shell.

Optical: Filaments, brighter in NE.

References:

- Duncan *et al.* 1997, MNRAS, 287, 722. Parkes 64-m at 2.4 GHz (11' : $S = 19 \pm 3$ Jy).
 Stupar *et al.* 2007, MNRAS, 374, 1441. Optical and various radio observations.

G315.4–2.3

RA: 14^h43^m00^s
Dec: −62°30'

1-GHz flux/Jy: 49
Spectral index: 0.6

RCW 86, MSH 14–63

Size/arcmin: 42
Type: S

Possibly the remnant of the SN of AD185?

Radio: Shell, brightest to the SW.

Optical: Bright, radiative filaments, with some faint Balmer dominated filaments.

X-ray: Partial shell, with thermal and non-thermal emission.

Point sources: Several X-ray sources.

Distance: Optical observations imply 2.3 kpc.

References:

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S = 86$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 18.2$ Jy).
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
 Winkler 1978, ApJ, 221, 220. X-ray detection.
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
 Pisarski *et al.* 1984, ApJ, 277, 710. Einstein observations
 Nugent *et al.* 1984, ApJ, 284, 612. X-ray spectrum.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 50'').
 Long & Blair 1990, ApJ, 358, L13. Balmer dominated optical filaments.
 Greidanus & Strom 1990, A&A, 240, 385. IRAS observations.
 Kastra *et al.* 1992, A&A, 264, 654. Ginga X-ray spectra.
 Strom 1994, MNRAS, 268, L5. Historical association
 Chin & Huang 1994, Nature, 371, 398. Questioning of historical association.
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'') : $S > 22$).
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Smith *et al.* 1997, AJ, 114, 2664. Observations of Balmer dominated filaments.
 Vink *et al.* 1997, A&A, 328, 628. ASCA spectroscopy.
 Bocchino *et al.* 2000, A&A, 360, 671. BeppoSAX observations of N and SW.
 Bamba *et al.* 2000, PASJ, 52, 1157. ASCA observations.
 Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
 Dickel *et al.* 2001, ApJ, 546, 447. ATCA at 1.34 GHz (8''), including polarisation.
 Bokowski *et al.* 2001, ApJ, 550, 334. ASCA observations.
 Rho *et al.* 2002, ApJ, 581, 1116. Chandra observations.
 Gvaramadze & Vikhlinin 2003, A&A, 401, 625. Chandra point source search.
 Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
 Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Vink *et al.* 2006, ApJ, 648, L33. Chandra and XMM observations.
 Ueno *et al.* 2007, PASJ, 59, S171. Suzaku observations.

G315.4–0.3

RA: 14^h35^m55^s
Dec: −60°36'

1-GHz flux/Jy: 8
Spectral index: 0.4

Size/arcmin: 24 \times 13
Type: ?

Radio: Irregular non-thermal emission, with HII region superposed in E.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 15.9$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 4.9$ Jy).
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'' : $S = 6.25$ Jy), re-assessment of earlier flux densities.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 49'') : $S = 3.1$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G315.9–0.0

RA: 14^h38^m25^s
Dec: −60°11'

1-GHz flux/Jy: 0.8?
Spectral index: ?

Size/arcmin: 25 × 14
Type: S

Has been called G315.8–0.0.

Radio: Faint, distorted shell, with steep-spectrum ‘jet’?

References:

Kesteven *et al.* 1987, *AuJPh*, 40, 855. MOST at 843 MHz (44'' × 50'').
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 50'': $S=0.9$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m and ATCA OH observations.

G316.3–0.0

(MSH 14–57)

RA: 14^h41^m30^s
Dec: −60°00'

1-GHz flux/Jy: 20?
Spectral index: 0.4

Size/arcmin: 29 × 14
Type: S

Radio: Distorted shell, with possible ‘blowout’.

X-ray: Detected.

Distance: HI absorption data suggests > 7.2 kpc.

References:

Shaver & Goss 1970, *AuJPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1975, *A&A*, 45, 239. Parkes HI absorption.
 Milne & Dickel 1975, *AuJPh*, 28, 209. Parkes 64-m at 5 GHz (4.4': $S=16.7$ Jy).
 Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44'' × 51'').
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 50'': $S=20$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.

G317.3–0.2

RA: 14^h49^m40^s
Dec: −59°46'

1-GHz flux/Jy: 4.7?
Spectral index: ?

Size/arcmin: 11
Type: S

Radio: Incomplete shell.

References:

Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 50'': $S=5.2$ Jy).

G318.2+0.1

RA: 14^h54^m50^s
Dec: −59°04'

1-GHz flux/Jy: >3.9?
Spectral index: ?

Size/arcmin: 40 × 35
Type: S

Radio: Faint shell, with central HII region.

X-ray: Sources within remnant.

References:

Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 50'': $S>4.3$).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Bocchino *et al.* 2001, *A&A*, 367, 629. BeppoSAX and ROSAT observations.

G318.9+0.4

RA: 14^h58^m30^s
Dec: −58°29'

1-GHz flux/Jy: 4?
Spectral index: 0.2?

Size/arcmin: 30 × 14
Type: C

May not be a SNR?

Radio: Complex arcs, with off-centre core.

References:

- Whiteoak 1990, *Nature*, 347, 157. MOST at 843 MHz (43'' × 51'').
 Whiteoak 1993, *ApJ*, 415, 701. MOST at 843 MHz (43'' × 51'') : $S = 4.8 \pm 0.6$ Jy, Parkes 64-m at 4.5 GHz (4'.7 : $S = 3.7 \pm 0.2$ Jy) and 8.4 GHz (2'.8 : $S = 3.0 \pm 0.4$ Jy) including polarisation, and ATCA at 1.4 GHz and 4.8 GHz (11'' × 13'') of core.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 50'') : $S = 4.8$ Jy.
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

G320.4−1.2

RA: 15^h14^m30^s
Dec: −59°08'

1-GHz flux/Jy: 60?
Spectral index: 0.4

MSH 15–52, RCW 89

Size/arcmin: 35
Type: C

Has been suggested as the remnant of the SN of AD185?

Radio: Ragged shell.

Optical: RCW 89 is the Hα emitting region to the NW.

X-ray: Partial shell, central nebula and pulsar and ‘jet’.

Point sources: Radio and X-ray pulsar, with wind nebula.

Distance: HI absorption indicates 5.2 kpc.

References:

- Caswell *et al.* 1975, *A&A*, 45, 239. Parkes HI absorption.
 Dickel & Milne 1976, *AuJPh*, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
 Dopita *et al.* 1977, *ApJ*, 214, 179. Optical spectra.
 Caswell *et al.* 1981, *MNRAS*, 195, 89. FIRST at 1415 MHz (50'').
 Seward & Harnden 1982, *ApJ*, 256, L45. X-ray observations of pulsar.
 Manchester *et al.* 1982, *ApJ*, 262, L31. Radio observations of pulsar.
 Seward *et al.* 1983, *ApJ*, 267, 698. X-ray, Optical and IR.
 van den Bergh & Kamper 1984, *ApJ*, 280, L51. Optical expansion.
 Seward *et al.* 1984, *ApJ*, 281, 650. X-ray observations of pulsar and nebulosity.
 Lortet *et al.* 1987, *A&A*, 180, 65. Optical observations.
 Trussoni *et al.* 1990, *A&A*, 234, 403. EXOSAT observations.
 Asaoka & Koyama 1990, *PASJ*, 42, 625. Ginga X-ray spectrum.
 Arendt 1991, *AJ*, 101, 2160. IRAS observations, including compact source.
 Milne *et al.* 1993, *MNRAS*, 264, 853. Parkes 64-m at 4.8 GHz (4'.5 : $S = 37 \pm 7$ Jy) and 8.4 GHz (3'.0 : $S = 24 \pm 4$ Jy), including polarisation and review of flux densities.
 Strom 1994, *MNRAS*, 268, L5. Historical association.
 Chin & Huang 1994, *Nature*, 371, 398. Questioning of historical association.
 Matz *et al.* 1994, *ApJ*, 434, 288. X-ray observations of pulsar.
 Schaefer 1995, *AJ*, 110, 1793. Questioning of historical association.
 Du Plessis *et al.* 1995, *ApJ*, 453, 746. Hartesbeesthoek 26-m at 2.3, 5 and 8.5 GHz ($S = 42, 35.6, 14.5$ Jy).
 Greiveldinger *et al.* 1995, *ApJ*, 454, 855. ROSAT observations.
 Trassoni *et al.* 1996, *A&A*, 306, 581. ROSAT observations.
 Tamura *et al.* 1996, *PASJ*, 48, L33. ASCA observations.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 50'') : $S = 62$ Jy.
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Marsden *et al.* 1997, *ApJ*, 491, L39. X-ray spectroscopy.
 Brazier & Becker 1997, *MNRAS*, 284, 335. ROSAT observations.
 Gaensler *et al.* 1999, *MNRAS*, 305, 724. ATCA at 1.4 GHz (24'' × 21''), plus HI observations, and 5.3 GHz (15'' × 10'').
 Sako *et al.* 2000, *ApJ*, 537, 422. Possible high energy γ-ray detection of pulsar.
 Mineo *et al.* 2001, *A&A*, 380, 695. BeppoSAX observations.
 Gaensler *et al.* 2002, *ApJ*, 569, 878. Chandra observations of pulsar and nebula.
 Dubner *et al.* 2002, *AJ*, 123, 337. ATCA at 1.4 GHz (4'.0 × 2'.7), plus HI observations.
 Aharonian *et al.* 2005, *A&A*, 435, L17. γ-ray detection.
 Yatsu *et al.* 2005, *ApJ*, 631, 312. Chandra observations of pulsar and jet.
 DeLaney *et al.* 2006, *ApJ*, 640, 929. Chandra and ROSAT multi-epoch observations of pulsar wind nebula.
 Kaplan & Moon 2006, *ApJ*, 644, 1056. Possible IR detection of pulsar.
 Forot *et al.* 2006, *ApJ*, 651, L45. X-ray observations of pulsar and nebula.
 Nakamori *et al.* 2008, *ApJ*, 677, 297. γ-ray observations.

G320.6–1.6

RA: $15^{\text{h}}17^{\text{m}}50^{\text{s}}$
Dec: $-59^{\circ}16'$

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 60×30
Type: S

Radio: Faint shell, overlapping G320.4–1.2 in W.

References:

Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz ($4'5$) and 8.4 GHz ($3'0$), including polarisation.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S > 9.3$).

G321.9–1.1

RA: $15^{\text{h}}23^{\text{m}}45^{\text{s}}$
Dec: $-58^{\circ}13'$

1-GHz flux/Jy: $>3.4?$
Spectral index: ?

Size/arcmin: 28
Type: S

Radio: Faint shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 51'' : S > 3.8$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G321.9–0.3

RA: $15^{\text{h}}20^{\text{m}}40^{\text{s}}$
Dec: $-57^{\circ}34'$

1-GHz flux/Jy: 13
Spectral index: 0.3

Size/arcmin: 31×23
Type: S

Radio: Shell brighter to the W, with Cir X-1 to N.

Point sources: Compact, probably thermal source at S edge.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 18.3$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.8$ Jy).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 52''$).
Stewart *et al.* 1993, MNRAS, 261, 593. ATCA at 1.5 GHz ($21''$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 51'' : S > 8.3$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Mignami *et al.* 2002, A&A, 386, 487. HST proper motion study of Cir X-1.
Tudose *et al.* 2006, MNRAS, 372, 417. MOST at 843 MHz.

G322.5–0.1

RA: $15^{\text{h}}23^{\text{m}}23^{\text{s}}$
Dec: $-57^{\circ}06'$

1-GHz flux/Jy: 1.5
Spectral index: 0.4

Size/arcmin: 15
Type: C

Radio: Shell with central extended source.

Point sources: PN Pe 2-8 within boundary.

References:

Whiteoak 1992, MNRAS, 256, 121. MOST at 843 MHz ($43'' \times 51'' : S = 2.0 \pm 0.3$ Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 51'' : S = 1.7$ Jy), plus Parkes 64-m at 4.5 GHz ($= 0.89 \pm 0.13$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G323.5+0.1

RA: 15^h28^m42^s
Dec: −56°21'

1-GHz flux/Jy: 3?
Spectral index: 0.4?

Size/arcmin: 13
Type: S

Radio: Distorted shell, confused with thermal emission.

Point sources: Compact, probably thermal source near centre.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 4.2$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 1.5$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 52'' : $S = 4.2$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G326.3−1.8

RA: 15^h53^m00^s
Dec: −56°10'

1-GHz flux/Jy: 145
Spectral index: varies

MSH 15–56

Size/arcmin: 38
Type: C

Has been called G326.2–1.7.

Radio: Shell, with elongated, flat-spectrum core.

Optical: Emission around the shell.

X-ray: Shell, with central extended emission.

References:

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3' : $S = 180$ Jy).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz (0'8 : $S > 95$) and Parkes 64-m at 14.7 GHz (2'2 : $S = 69 \pm 8$ Jy).
 van den Berg 1979, ApJ, 227, 497. Optical observations.
 Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.
 Dennefeld 1980, PASP, 92, 603. Optical spectra.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 53'').
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43'' \times 52'' : $S = 153 \pm 40$ Jy), and Parkes 64-m at 8.4 GHz (3' : $S = 68 \pm 5$ Jy), including polarisation.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Kassim *et al.* 1993, ApJ, 419, 733. ROSAT image.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 52'' : $S > 130$).
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Dickel *et al.* 2000, ApJ, 543, 840. ATCA at 1.34 GHz (8'5 \times 6'4 : $S > 60$ Jy), 4.80 GHz (3''8 : $S = 25$ Jy for core only) and 8.64 GHz (3''2 : $S = 15$ Jy for core only).

G327.1−1.1

RA: 15^h54^m25^s
Dec: −55°09'

1-GHz flux/Jy: 7?
Spectral index: ?

Size/arcmin: 18
Type: C

Radio: Shell, with off-centre core.

X-ray: Diffuse, with core.

References:

- Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (8'4 : $S = 10 \pm 15\%$ Jy).
 Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 10.6$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 4.3$ Jy).
 Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 52'' : $S = 7.6$ Jy).
 Seward *et al.* 1996, ApJ, 471, 887. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Sun *et al.* 1999, ApJ, 511, 274. ASCA and ROSAT observations.
 Bocchino & Bandiera 2003, A&A, 398, 195. BeppoSAX observations.

G327.2–0.1

RA: 15^h50^m55^s
Dec: −54°18'

1-GHz flux/Jy: 0.4
Spectral index: ?

Size/arcmin: 5
Type: S

Has been called G327.24–0.13.

Radio: Shell, possibly with central emission.

Point sources: Central pulsar (magnetar).

References:

- Camilo *et al.* 2007, ApJ, 666, L93. Pulsar observations.
 Gelfand & Gaensler 2007, ApJ, 667, 1111. Chandra and XMM observations, and 843 GHz (43'' × 53'': $S = 0.5 \pm 0.1$ Jy) and SGPS at 1.4 GHz ($S = 0.3 \pm 0.1$ Jy).

G327.4+0.4

RA: 15^h48^m20^s
Dec: −53°49'

1-GHz flux/Jy: 30?
Spectral index: 0.6

Size/arcmin: 21
Type: S

Kes 27

Has been called G327.3+0.4 and G327.3+0.5.

Radio: Incomplete, multi-arc shell, brightest to the SE.

X-ray: Diffuse, best defined to E.

Distance: HI absorption indicates 4.3 to 5.4 kpc.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3': $S = 58$ Jy) and Parkes 64-m at 5 GHz (4': $S = 12.4$ Jy).
 Dickey & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
 Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 55'').
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43'' × 53'': $S = 32.2 \pm 6$ Jy), and Parkes 64-m at 8.4 GHz (3': $S = 9.4 \pm 0.8$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 53'': $S = 25$ Jy).
 Seward *et al.* 1996, ApJ, 471, 887. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'0 × 1'8), plus HI.
 Enoguchi *et al.* 2002, PASJ, 54, 229. ASCA observations.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Chen *et al.* 2008, ApJ, 676, 1040. Chandra observations.

G327.4+1.0

RA: 15^h46^m48^s
Dec: −53°20'

1-GHz flux/Jy: 1.9?
Spectral index: ?

Size/arcmin: 14
Type: S

Radio: Asymmetric shell.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 54'': $S = 2.1$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'0 × 1'8), plus HI.

G327.6+14.6		SN1006, PKS 1459–41
RA: 15 ^h 02 ^m 50 ^s	1-GHz flux/Jy: 19	Size/arcmin: 30
Dec: −41°56'	Spectral index: 0.6	Type: S
This is the remnant of the SN of AD1006.		
Radio: Shell, with two bright arcs.		
Optical: Filaments to the NW, with broad H α component.		
X-ray: Thermal shell, with non-thermal limb-brightened arcs.		
Point sources: The background Schweizer–Middleditch star is near the middle of the remnant.		
Distance: Optical spectra and proper motion indicate 2.2 kpc.		
References:		
van den Bergh 1976, ApJ, 208, L17. Optical observations.		
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8.4') and 5 GHz (4.4').		
Becker <i>et al.</i> 1980, ApJ, 240, L33. X-ray spectroscopy.		
Schweizer & Middleditch 1980, ApJ, 241, 1039. Possible stellar remnant.		
Pye <i>et al.</i> 1981, MNRAS, 194, 569. Einstein observations.		
Caswell <i>et al.</i> 1983, MNRAS, 204, 921. FIRST at 1415 MHz (77'').		
Reynolds & Gilmore 1986, AJ, 92, 1138. VLA at 1.37 and 1.67 GHz (16'' × 20'').		
Kirshner <i>et al.</i> 1987, ApJ, 315, L135. Broad H α optical component.		
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 66'').		
Fesen <i>et al.</i> 1988, ApJ, 327, 164. UV absorption spectra of the Schweizer–Middleditch star.		
Roger <i>et al.</i> 1988, ApJ, 332, 940. MOST at 843 MHz (44'' × 66''): $S=17.5 \pm 1.5$ Jy).		
Long <i>et al.</i> 1988, ApJ, 333, 749. Optical proper motion for distance.		
Jones & Pye 1989, MNRAS, 238, 567. EXOSAT observations.		
Leahy <i>et al.</i> 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.		
Wu <i>et al.</i> 1993, ApJ, 416, 247. UV spectra.		
Reynolds & Gilmore 1993, AJ, 106, 272. VLA at 1.37 and 1.67 GHz (24''), including polarisation.		
Moffett <i>et al.</i> 1993, AJ, 106, 1566. VLA at 1.37 and 1.67 GHz (10'' × 14''), for proper motion studies.		
Ozaki <i>et al.</i> 1994, PASJ, 46, 367. X-ray observations.		
Raymond <i>et al.</i> 1995, ApJ, 454, L31. Far UV spectroscopy.		
Koyama <i>et al.</i> 1995, Nature, 378, 255. ASCA observations.		
Willingale <i>et al.</i> 1996, MNRAS, 278, 749. ROSAT observations.		
Blair <i>et al.</i> 1996, ApJ, 468, 871. Optical absorption studies.		
Laming <i>et al.</i> 1996, ApJ, 472, 267. Modelling of optical spectra, including distance.		
Wu <i>et al.</i> 1997, ApJ, 477, L53. Far UV absorption spectra of the Schweizer–Middleditch star.		
Winkler & Long 1997, ApJ, 486, L137. UV absorption spectra of background quasar.		
Winkler & Long 1997, ApJ, 491, 829. ROSAT and optical images.		
Tanimori <i>et al.</i> 1998, ApJ, 497, L25. γ -ray detection.		
Vink <i>et al.</i> 2000, A&A, 354, 931. X-ray spectroscopy.		
Burleigh <i>et al.</i> 2000, A&A, 356, 585. Optical spectroscopy of the Schweizer–Middleditch star.		
Allen <i>et al.</i> 2001, ApJ, 558, 739. ASCA, ROSAT and other X-ray observations.		
Dubner <i>et al.</i> 2002, A&A, 387, 1047. ATCA at 1.4 GHz (4.7' × 3.0') for H α , plus CO observations.		
Ghavamian <i>et al.</i> 2002, ApJ, 572, 888. Optical of filaments in NW.		
Sollerman <i>et al.</i> 2003, A&A, 407, 249. Optical spectroscopy.		
Winkler <i>et al.</i> 2003, ApJ, 585, 324. Optical proper motion studies.		
Vink <i>et al.</i> 2003, ApJ, 587, L31. Chandra observations.		
Bamba <i>et al.</i> 2003, ApJ, 589, 827. Chandra observations of NE.		
Koreck <i>et al.</i> 2004, ApJ, 615, 280. Far-uv observations.		
Aharonian <i>et al.</i> 2005, A&A, 437, 135. γ -ray limit.		
Winkler <i>et al.</i> 2005, ApJ, 624, 189. HST absorption towards background sources.		
Kalemci <i>et al.</i> 2006, ApJ, 640, L55. γ -ray limit upper limit.		
Kalemci <i>et al.</i> 2006, ApJ, 644, 274. Integral observations.		
Acero <i>et al.</i> 2007, A&A, 475, 883. XMM observations.		
Raymond <i>et al.</i> 2007, ApJ, 659, 1257. HST H α observations.		
Hamilton <i>et al.</i> 2007, MNRAS, 381, 771. UV spectroscopy of the Schweizer–Middleditch star.		
Bamba <i>et al.</i> 2008, AdSpR, 41, 411. Suzaku observations.		
Cassam-Chanaï <i>et al.</i> 2008, ApJ, 680, 1180. ATCA and VLA at 1.5 GHz (6'' × 9'') plus Chandra observations.		
Yamaguchi <i>et al.</i> 2008, PASJ, 60, S141. Suzaku observations.		
Bamba 2008, PASJ, 60, S153. Suzaku observations.		

G328.4+0.2		(MSH 15–57)
RA: 15 ^h 55 ^m 30 ^s	1-GHz flux/Jy: 15	Size/arcmin: 5
Dec: −53°17'	Spectral index: 0.0	Type: F

Radio: Amorphous emission, with central bar.

X-ray: Detected at high energies.

Distance: HI absorption indicates > 17.4 kpc.

References:

- Shaver & Goss 1970, *AuJPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1980, *MNRAS*, 190, 881. FIRST at 1415 MHz (50'').
 Wilson 1986, *ApJ*, 302, 718. Lack of detection with Einstein.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 54'': $S = 15$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Gaensler *et al.* 2000, *ApJ*, 542, 380. ATCA at 1.4 GHz (16''.5 × 19''.5: $S = 14.3 \pm 0.1$ Jy) and 4.5 GHz (2''.0 × 1''.5: $S = 12.5 \pm 0.2$ Jy).
 Hughes *et al.* 2000, *ApJ*, 542, 386. ASCA observations.
 McClure-Griffiths *et al.* 2001, *ApJ*, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1''.8 × 2''.0), plus HI.
 Johnston *et al.* 2004, *MNRAS*, 348, L19. ATCA at 19 GHz (6''.1 × 7''.7), including polarisation.
 Gelfand *et al.* 2007, *ApJ*, 663, 468. ATCA at 1.4 GHz (5''.8 × 7''.0: $S = 13.8 \pm 0.4$ Jy), plus XMM observations.

G329.7+0.4

RA: 16 ^h 01 ^m 20 ^s	1-GHz flux/Jy: >34?	Size/arcmin: 40 × 33
Dec: −52°18'	Spectral index: ?	Type: S

Radio: Diffuse shell, in complex region.

References:

- Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 53'': $S > 38$).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, *ApJ*, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2''.0 × 1''.8), plus HI.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer possible detection.

G330.0+15.0

	Lupus Loop
RA: 15 ^h 10 ^m 00 ^s	Size/arcmin: 180?
Dec: −40°00'	Type: S

Radio: Low surface brightness loop with HI shell.

X-ray: Detected, with central source.

Point sources: Central, possibly pulsating, X-ray source.

References:

- Milne 1971, *AuJPh*, 24, 757. Parkes 64-m at 408 MHz (48'), 635 MHz (31') and 1410 MHz (15').
 Milne & Dickel 1974, *AuJPh*, 27, 549. Parkes 64-m at 2.7 GHz (8''.4: $S = 120 \pm 30\%$ Jy).
 Toor 1980, *A&A*, 85, 184. X-ray image and spectrum.
 Colomb & Dubner 1982, *A&A*, 112, 141. Argentine 30-m dish at 1.42 GHz (30'), HI observations.
 Leahy *et al.* 1991, *ApJ*, 374, 218. HEAO-1 X-ray spectra.
 Ozaki *et al.* 1994, *PASJ*, 46, 367. X-ray observations.
 Kaplan *et al.* 2006, *ApJS*, 163, 344. X-ray upper limit on compact sources.
 Shinn *et al.* 2006, *ApJ*, 644, L189. Far UV observations.
 Park *et al.* 2006, *ApJ*, 653, L37. Chandra observations.

G330.2+1.0

RA: 16^h01^m06^s
Dec: -51°34'

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 11
Type: S?

Radio: Clumpy non-thermal emission, possibly a distorted shell.

X-ray: Shell.

Distance: HI absorption indicates > 4.9 kpc.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=8.6$ Jy) and Parkes 64-m at 5 GHz (4' : $S=4.0$ Jy).
 Caswell *et al.* 1983, MNRAS, 204, 915. FIRST at 1415 MHz (52'' \times 47''), and MOST at 843 MHz (43'' \times 55'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 55'') : $S=4.7$ Jy.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'.0 \times 1'.8), plus HI.
 Torii *et al.* 2006, PASJ, 58, L11. ASCA detection.

G332.0+0.2

RA: 16^h13^m17^s
Dec: -50°53'

1-GHz flux/Jy: 8?
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Incomplete shell.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 57'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 55'') : $S=8.9$ Jy.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'.0 \times 1'.8), plus HI.

G332.4–0.4

RA: 16^h17^m33^s
Dec: −51°02'

1-GHz flux/Jy: 28
Spectral index: 0.5

RCW 103

Size/arcmin: 10
Type: S

Radio: Shell, brightest to the S.

Optical: Filaments correspond well to the radio shell, brightest in SE.

X-ray: Brightest to NW, with point source near centre.

Point sources: Central, variable X-ray source, and nearby pulsar.

Distance: HI absorption indicates 3.1 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Tuohy & Garmire 1980, ApJ, 239, L107. Einstein detection of X-ray point source.
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50'').
 Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.
 Ruiz 1983, AJ, 88, 1210. Optical spectra.
 Tuohy *et al.* 1983, ApJ, 268, 778. X-ray observations.
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
 Nugent *et al.* 1984, ApJ, 284, 612. X-ray spectrum (and Einstein image from Tuohy, private communication).
 Meaburn & Allen 1986, MNRAS, 222, 593. Optical spectra.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 57'').
 Oliva *et al.* 1990, A&A, 240, 453. IR spectroscopy.
 Dickel *et al.* 1996, AJ, 111, 340. ATCA at 1.36 (8'') and 2.37 GHz (4''5''), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 55'': $S=34$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Gotthelf *et al.* 1997, ApJ, 487, L175. ASCA observations of compact X-ray source.
 Carter *et al.* 1997, PASP, 109, 990. Optical expansion.
 Torii *et al.* 1998, ApJ, 494, L207. ASCA detection of nearby pulsar.
 Kaspi *et al.* 1998, ApJ, 503, L161. Pulsar observations.
 Gotthelf *et al.* 1999, ApJ, 514, L107. X-ray variability of central source.
 Oliva *et al.* 1999, A&A, 343, 943. ISO spectroscopy.
 Torii *et al.* 2000, ApJ, 534, L71. X-ray timing on pulsar, including glitch.
 Vink 2004, ApJ, 604, 693. Chandra observations.
 Reynolds *et al.* 2004, PASA, 21, 82. ATCA at 1.4 GHz (50''), including HI absorption to central source.
 Russel *et al.* 2005, A&A, 429, 497. H α observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Paron 2006, PASA, 23, 69. CO and HCO⁺ observations of surroundings.
 De Luca 2006, Science, 313, 814. XMM observations of central source.
 De Luca 2007, Ap&SS, 308, 231. XMM observation of periodicity of central source.
 Matsumoto *et al.* 2007, PASJ, 59, S199. Suaku observations.
 De Luca 2008, ApJ, 682, 1185. IR observations of central source.

G332.4+0.1

RA: 16^h15^m20^s
Dec: −50°42'

1-GHz flux/Jy: 26
Spectral index: 0.5

MSH 16–51, Kes 32

Size/arcmin: 15
Type: S

Has been called G332.4+0.2.

Radio: Distorted shell, with thermal jet and plume adjacent.

X-ray: Shell, brightest to NW.

Point sources: Pulsar nearby.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Roger *et al.* 1985, Nature, 316, 44. MOST at 843 MHz (44'' × 57'').
 Caraveo 1993, ApJ, 415, L111. Nearby pulsar.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 56'': $S=29$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Brinkmann *et al.* 1999, A&A, 346, 599. ROSAT image of remnant and nearby pulsar.
 Vink *et al.* 2004, ApJ, 604, 699. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G332.5–5.6

RA: 16^h43^m20^s
Dec: −54°30'

1-GHz flux/Jy: 2?
Spectral index: 0.7?

Size/arcmin: 35
Type: S

Radio: Bipolar shell, with central emission also.

Optical: Patchy filaments.

X-ray: Emission from centre.

References:

- Reynoso & Green 2007, MNRAS, 375, 92. ATCA at 1.4 GHz (40'': $S = 1.90 \pm 0.15$ Jy) and 2.4 GHz (90'': $S = 1.3 \pm 0.2$ Jy) including polarisation, and 843 MHz (43'' \times 53'') plus ROSAT observations.
 Stupar *et al.* 2007, MNRAS, 381, 377. Optical observations, plus ATCA at 1.4 GHz (58'') and 2.4 GHz (95'') including polarisation, and other observations.

G335.2+0.1

RA: 16^h27^m45^s
Dec: −48°47'

1-GHz flux/Jy: 16
Spectral index: 0.5

Size/arcmin: 21
Type: S

Radio: Well defined shell.

Point sources: Old pulsar within remnant boundary.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3'': $S = 27.1$ Jy) and Parkes 64-m at 5 GHz (4'': $S = 8.6$ Jy).
 Kaspi *et al.* 1996, AJ, 111, 2028. Pulsar observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 57'': $S = 16$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G336.7+0.5

RA: 16^h32^m11^s
Dec: −47°19'

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 14 \times 10
Type: S

Radio: Irregular shell.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 60'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 58'': $S = 6.1$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G337.0–0.1

(CTB 33)

RA: 16^h35^m57^s
Dec: −47°36'

1-GHz flux/Jy: 1.5
Spectral index: 0.6?

Size/arcmin: 1.5
Type: S

This entry refers to a small (1.5') SNR, not the larger previously catalogued G337.0–0.1.

Radio: Shell, in a complex region.

Distance: Association with CTB 33 gives 11 kpc.

Point sources: Associated with a soft gamma repeater.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 58'': $S = 21$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Sarma *et al.* 1997, ApJ, 483, 335. ATCA at 1.4 GHz (12'') including H_i, and 8.9 GHz (13'' \times 15'') for recombination lines, clarifying extent of the remnant.
 Woods *et al.* 1999, ApJ, 519, L139. Soft gamma repeater observations.
 Hurley *et al.* 2000, ApJ, 528, L21. ASCA observations of soft gamma repeater.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

G337.2–0.7

RA: 16^h39^m28^s
Dec: −47°51'

1-GHz flux/Jy: 1.5
Spectral index: 0.4

Size/arcmin: 6
Type: S

Radio: Shell, brighter in S.

X-ray: Extended emission.

Distance: HI absorption suggests 2.0 to 9.3 kpc.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 3.8$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 0.70$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 58'' : S = 2.0$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT observations.
 Rakowski *et al.* 2006, ApJ, 646, 982. ATCA observations at 1.3 GHz ($15'' : S = 1.55 \pm 0.05$ Jy) and 5 GHz ($15'' : S = 0.93 \pm 0.02$ Jy), plus HI and Chandra observations.

G337.2+0.1

RA: 16^h35^m55^s
Dec: −47°20'

1-GHz flux/Jy: 1.5?
Spectral index: ?

Size/arcmin: 3 × 2
Type: ?

Radio: Not well defined.

X-ray: Detected.

Distance: Association with HI hole gives 14 kpc.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59'' : S = 1.6$ Jy)
 Combi *et al.* 2005, A&A, 431, L9. ASCA and other observations.
 Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations of nearby source.
 Combi *et al.* 2006, ApJ, 653, L41. XMM observations.

G337.3+1.0

Kes 40

RA: 16^h32^m39^s
Dec: −46°36'

1-GHz flux/Jy: 16
Spectral index: 0.55

Size/arcmin: 15 × 12
Type: S

Radio: Nearly complete shell.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ($3' : S = 24.6$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.2$ Jy).
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8.4') and 5 GHz (4.4').
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 61''$).
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz ($43'' \times 59'' : S = 14.8 \pm 3.0$ Jy), and Parkes 64-m at 8.4 GHz ($3' : S = 5.1 \pm 0.6$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59'' : S = 20$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G337.8–0.1

RA: 16^h39^m01^s
Dec: −46°59'

1-GHz flux/Jy: 18
Spectral index: 0.5

Kes 41

Size/arcmin: 9 × 6
Type: S

Radio: Distorted shell.**X-ray:** Centrally brightened.**Distance:** HI absorption suggests 11 kpc.**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'': $S = 18$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz, for associated OH masers.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including HI.
Combi *et al.* 2008, A&A, 488, L25. XMM observations.

G338.1+0.4

RA: 16^h37^m59^s
Dec: −46°24'

1-GHz flux/Jy: 4?
Spectral index: 0.4

Size/arcmin: 15?
Type: S

Radio: Arc in NE, merging with thermal emission in S.**Optical:** Detected.**X-ray:** Detected.**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'': $S = 3.8$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz of associated OJ masers.

G338.3–0.0

RA: 16^h41^m00^s
Dec: −46°34'

1-GHz flux/Jy: 7?
Spectral index: ?

Size/arcmin: 8
Type: C?

Radio: Irregular shell, in complex region.**X-ray:** Central X-ray source and nebula.**Distance:** HI observations suggest 11 kpc.**References:**

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'': $S = 7.4$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Aharonian *et al.* 2005, Science, 307, 1938. γ-ray detection.
Aharonian *et al.* 2006, ApJ, 636, 777. γ-ray observations.
Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including HI.
Funk *et al.* 2007, ApJ, 662, 517. XMM observations.

G338.5+0.1

RA: 16^h41^m09^s
Dec: −46°19'

1-GHz flux/Jy: 12?
Spectral index: ?

Size/arcmin: 9
Type: ?

Radio: Circle of non-thermal emission in complex region, not well defined.

Distance: HI absorption suggests 11 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 59'': S = 13 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including HI.

G340.4+0.4

RA: 16^h46^m31^s
Dec: −44°39'

1-GHz flux/Jy: 5
Spectral index: 0.4

Size/arcmin: 10 × 7
Type: S

Radio: Distorted shell, elongated east–west.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S = 8.2 Jy) and Parkes 64-m at 5 GHz (4': S = 2.9 Jy).
Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50'').
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (75'' × 48'': S = 9.8 ± 0.9 Jy) and 1.4 GHz (27'' × 9'': S = 3.6 ± 0.1 Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 61'': S = 5.9 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G340.6+0.3

RA: 16^h47^m41^s
Dec: −44°34'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 6
Type: S

Radio: Incomplete shell.

Optical: Possible associated filaments.

Distance: HI absorption suggests 15 kpc.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S = 7.0 Jy) and Parkes 64-m at 5 GHz (4': S = 2.8 Jy).
Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50'').
Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (75'' × 48'': S = 9.2 ± 0.9 Jy) and 1.4 GHz (27'' × 9'': S = 5.8 ± 0.1 Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 61'': S = 4.5 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including HI.

G341.2+0.9

RA: 16^h47^m35^s
Dec: −43°47'

1-GHz flux/Jy: 1.5?
Spectral index: 0.6?

Size/arcmin: 22 × 16
Type: C

Radio: Incomplete shell, with extension to SW.

Point sources: Pulsar in W, with wind nebula.

References:

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (54'' × 116'': S = 3.0 ± 0.1 Jy) and 1.4 GHz (21'' × 25'': S = 12.5 ± 0.05 Jy).
Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4 and 4.9 GHz (25'') of pulsar wind nebula.

G341.9–0.3

RA: 16^h55^m01^s
Dec: −44°01'

1-GHz flux/Jy: 2.5
Spectral index: 0.5

Size/arcmin: 7
Type: S

Radio: Incomplete shell, brightest to NE.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S=7.4$ Jy) and Parkes 64-m at 5 GHz (4' : $S=1.7$ Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50''), revision of previous flux densities.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (35'' × 11'') : $S=2.2 \pm 0.1$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 62'') : $S=2.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.0–0.2

RA: 16^h54^m50^s
Dec: −43°53'

1-GHz flux/Jy: 3.5?
Spectral index: 0.4?

Size/arcmin: 12 × 9
Type: S

Radio: Distorted shell.

References:

- Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50''), estimate $S_{408\text{ MHz}}=5$ Jy, $S_{5\text{ GHz}}=2$ Jy from previous maps.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (35'' × 11'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 62'') : $S=3.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.1+0.9

RA: 16^h50^m43^s
Dec: −43°04'

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 10 × 9
Type: S

Radio: Incomplete shell.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 63'') : $S=0.6$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G343.0–6.0

RA: 17^h25^m00^s
Dec: −46°30'

1-GHz flux/Jy: ?
Spectral index: ?

RCW 114

Size/arcmin: 250
Type: S

Radio: Faint, poorly defined.

Optical: Filamentary shell.

References:

- Walker & Zealey 2001, MNRAS, 325, 287. Optical observations, and review of earlier observations.
 Welsh *et al.* 2003, A&A, 403, 605. Optical spectroscopy.
 Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.

G343.1–2.3

RA: 17^h08^m00^s
Dec: −44°16'

1-GHz flux/Jy: 8?
Spectral index: 0.5?

Size/arcmin: 32?
Type: C?

Radio: Incomplete shell?

X-ray: Pulsar wind nebula.

Point sources: Pulsar near edge, with wind nebula.

References:

- McAdam *et al.* 1993, Nature, 361, 516. MOST at 843 MHz (smoothed to 2').
 Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz ($56'' \times 11''$: $S = 10.6$ Jy) and 1.4 GHz ($22'' \times 27''$) near pulsar.
 Becker *et al.* 1995, A&A, 298, 528. ROSAT of pulsar, and limit for remnant.
 Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4, 4.9 and 8.5 GHz ($25''$) of pulsar wind nebula.
 Gotthelf *et al.* 2002, ApJ, 567, L125. Pulsar x-ray detection.
 Dodson & Golap 2002, MNRAS, 334, L1. ATCA at 1.4 GHz ($70'' \times 47''$) including polarisation, and Chandra observations of pulsar wind nebula.
 Aharonian *et al.* 2005, A&A, 432, L9. γ -ray limit.

G343.1–0.7

RA: 17^h00^m25^s
Dec: −43°14'

1-GHz flux/Jy: 7.8
Spectral index: 0.55

Size/arcmin: 27 × 21
Type: S

Radio: Shell, with smaller thermal shell adjacent.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 63''$: $S = 8.5 \pm 0.6$ Jy), plus Parkes 64-m at 4.5 GHz ($S = 3.9 \pm 0.6$ Jy) and 8.55 GHz ($S = 2.4 \pm 0.5$ Jy)
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G344.7–0.1

RA: 17^h03^m51^s
Dec: −41°42'

1-GHz flux/Jy: 2.5?
Spectral index: 0.5

Size/arcmin: 10
Type: C?

Radio: Asymmetric shell, with possible core.

X-ray: Detected.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 4.7$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 1.3$ Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($30'' \times 43''$: $S = 1.7 \pm 0.1$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 65''$: $S = 2.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Yamauchi *et al.* 2005, PASJ, 57, 459. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

G345.7–0.2

RA: 17^h07^m20^s
Dec: −40°53'

1-GHz flux/Jy: 0.6?
Spectral index: ?

Size/arcmin: 6
Type: S

Radio: Poorly defined diffuse shell.

Point sources: Old pulsar nearby.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 66''$: $S = 0.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G346.6–0.2

RA: 17^h10^m19^s
Dec: −40°11'

1-GHz flux/Jy: 8?
Spectral index: 0.5?

Size/arcmin: 8
Type: S

Radio: Irregular shell.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 14.9$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 4.3$ Jy).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($31'' \times 43'' : S = 8.1 \pm 0.9$ Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 67'' : S = 8.7$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

G347.3–0.5

RA: 17^h13^m50^s
Dec: −39°45'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 65×55
Type: S?

Radio: Faint emission.

X-ray: Non-thermal, limb-brightened to W, with central source.

Point sources: Central X-ray source.

Distance: Association with molecular clouds and X-ray observations imply 1.3 kpc.

References:

- Koyama *et al.* 1997, PASJ, 49, L7. ASCA of NW.
Slane *et al.* 1999, ApJ, 525, 357. ASCA and other observations.
Muraishi *et al.* 2000, A&A, 354, L57. High energy γ -ray detection.
Butt *et al.* 2001, ApJ, 562, L167. Associated γ -ray emission.
Ellison *et al.* 2001, ApJ, 563, 191. ATCA at 1.4 GHz ($46'' \times 36''$), and ASCA observations.
Enomoto *et al.* 2002, Nature, 416, 823. γ -ray observations.
Uchiyama *et al.* 2002, PASJ, 54, L73. ASCA observations.
Uchiyama *et al.* 2003, A&A, 400, 567. Chandra spectroscopy.
Pannuti *et al.* 2003, ApJ, 593, 377. ROSAT and ASCA observations.
Lazendic *et al.* 2003, ApJ, 593, L27. Chandra, XMM and other X-ray observations of central source.
Fukui *et al.* 2003, PASJ, 55, L61. CO observations of surroundings.
Cassam-Chenai *et al.* 2004, A&A, 427, 199. XMM and other observations.
Aharonian *et al.* 2004, Nature, 432, 75. γ -ray detection.
Lazendic *et al.* 2004, ApJ, 602, 271. Chandra observations of parts, and ATCA at 1.4 GHz ($36'' \times 46''$).
Hiraga *et al.* 2005, A&A, 431, 953. XMM observations,
Moriguchi *et al.* 2005, ApJ, 631, 947. CO observations of surroundings.
Aharonian *et al.* 2007, A&A, 464, 235. γ -ray observations.
Tanaka *et al.* 2008, ApJ, 685, 988. Suzaku observations.
Takahashi *et al.* 2008, PASJ, 60, S131. Suzaku observations of SW.
Uchiyama *et al.* 2007, Nature, 449, 576. Chandra multi-epoch observations for study of small-scale variability.
Mignami *et al.* 2008, A&A, 484, 457. Optical and IR observations of central source.

G348.5–0.0

RA: 17^h15^m26^s
Dec: −38°28'

1-GHz flux/Jy: 10?
Spectral index: 0.4?

Size/arcmin: 10?
Type: S?

Radio: Arc, overlapping G348.5+0.1.

References:

- Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz ($46'' \times 53''$), 1.4 GHz ($18'' \times 33''$) and part at 5 GHz ($2''.5 \times 3''.9$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 69'' : S = 10.2$ Jy).
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G348.5+0.1		CTB 37A
RA: 17 ^h 14 ^m 06 ^s	1-GHz flux/Jy: 72	Size/arcmin: 15
Dec: -38°32'	Spectral index: 0.3	Type: S

Radio: Shell, poorly define to S and W, overlapping G348.5–0.0 in E.

Distance: HI absorption indicates 8.0 kpc.

References:

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3' : $S=97$ Jy).
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4 : $S=43$ Jy).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz (0'.8 : $S>50$) and Parkes 64-m at 14.7 GHz (2'.2 : $S=18\pm5$ Jy).
 Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20'' \times 45'').
 Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz (46'' \times 53''), 1.4 GHz (18'' \times 33'') and part at 5 GHz (2''.5 \times 3''.9).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 69'': $S=71$ Jy).
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Reynoso & Mangum 2000, ApJ, 545, 874. CO observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Aharonian *et al.* 2008, A&A, 490, 685. γ -ray observations.

G348.7+0.3		CTB 37B
RA: 17 ^h 13 ^m 55 ^s	1-GHz flux/Jy: 26	Size/arcmin: 17?
Dec: -38°11'	Spectral index: 0.3	Type: S

Radio: Incomplete shell with faint eastern extensions.

X-ray: Diffuse emission.

Distance: HI absorption indicates 8.0 kpc.

References:

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3' : $S=34$ Jy).
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'.4 : $S=32$ Jy).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz (0'.8 : $S>20$) and Parkes 64-m at 14.7 GHz (2'.2 : $S=8\pm3$ Jy).
 Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20'' \times 45'').
 Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz (46'' \times 53'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 69'': $S=33$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.
 Aharonian *et al.* 2008, A&A, 486, 829. γ -ray and Chandra observations.

G349.2–0.1		Size/arcmin: 9 \times 6
RA: 17 ^h 17 ^m 15 ^s	1-GHz flux/Jy: 1.4?	Type: S
Dec: -38°04'	Spectral index: ?	

Radio: Elongated shell, adjacent to bright HII region.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 70'': $S=1.6$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

G349.7+0.2

RA: 17^h17^m59^s
Dec: −37°26'

1-GHz flux/Jy: 20
Spectral index: 0.5

Size/arcmin: 2.5 × 2
Type: S

Radio: Incomplete clumpy shell, with enhancement to the S.

Distance: HI absorption indicates 14.8 kpc, association with OH features gives 22 kpc.

X-ray: Irregular shell, brighter to S and E.

References:

- Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3': $S=31.0$ Jy) and Parkes 64-m at 5 GHz (4': $S=9.1$ Jy), no maps.
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz (3''.4 × 14''.5).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 71'': $S=22$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Reynoso & Mangum 2001, AJ, 121, 347. CO observations of the vicinity.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Slane *et al.* 2002, ApJ, 580, 904. ASCA observations.
 Dubner *et al.* 2004, A&A, 426, 201. CO observations of surroundings.
 Lazendic *et al.* 2005, ApJ, 618, 733. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G350.0−2.0

RA: 17^h27^m50^s
Dec: −38°32'

1-GHz flux/Jy: 26
Spectral index: 0.4

Size/arcmin: 45
Type: S

Incorporates the previously catalogued G350.0–1.8 in the NW.

Radio: Shell, brightest in NW.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=49.5$ Jy) and Parkes 64-m at 5 GHz (4': $S=13.6$ Jy).
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4''.4).
 Gaensler 1998, ApJ, 493, 781. VLA and Parkes 64-m at 1.4 GHz (18'' × 21'': $S=22.3 \pm 0.3$ Jy), clarifying extent of remnant.

G350.1−0.3

RA: 17^h17^m40^s
Dec: −37°24'

1-GHz flux/Jy: 6?
Spectral index: 0.8?

Size/arcmin: 4?
Type: ?

Radio: Several clumps of emission.

X-ray: Diffuse emission, with compact source.

Distance: HI absorption indicates 4.5 to 10.7 kpc, possible interaction with molecular cloud indicates 4.5 kpc.

Point sources: X-ray source.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=10.7$) and Parkes 64-m at 5 GHz (4': $S=1.7$).
 Salter *et al.* 1986, A&A, 162, 217. VLA at 1.5 (4'' × 10''.4) and 4.8 GHz (15'' × 35'') and 15 GHz (5'' × 2'').
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA observations.
 Gaensler *et al.* 2008, ApJ, 680, L37. VLA at 4.8 GHz (11''.4 × 5''.6) plus XMM observations.

G351.2+0.1

RA: 17^h22^m27^s
Dec: -36°11'

1-GHz flux/Jy: 5?
Spectral index: 0.4

Size/arcmin: 7
Type: C?

Has been called G351.3+0.2.

Radio: Distorted shell, with possible flat-spectrum core.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=8.1$ Jy) and Parkes 64-m at 5 GHz (4' : $S=3.1$ Jy).
 Becker & Helfand 1988, AJ, 95, 883. VLA at 5 GHz (15''), and at 15 GHz of core.
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (32'' × 36'': $S=4.8 \pm 0.2$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 73'': $S=5.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G351.7+0.8

RA: 17^h21^m00^s
Dec: -35°27'

1-GHz flux/Jy: 10
Spectral index: 0.5?

Size/arcmin: 18 × 14
Type: S

Radio: Elongated shell, adjacent to bright HII region.

Point sources: Pulsar nearby.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 74'': $S=11$ Jy).
 Tian *et al.* 2007, MNRAS, 378, 1283. SGPS at 1.4 GHz (100'': $S=8.4 \pm 0.7$ Jy) including HI.

G351.9-0.9

RA: 17^h28^m52^s
Dec: -36°16'

1-GHz flux/Jy: 1.8?
Spectral index: ?

Size/arcmin: 12 × 9
Type: S

Radio: Asymmetric shell.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 73'': $S=2.0$ Jy).

G352.7-0.1

RA: 17^h27^m40^s
Dec: -35°07'

1-GHz flux/Jy: 4
Spectral index: 0.6

Size/arcmin: 8 × 6
Type: S

Radio: Distorted shell.

X-ray: Detected.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S=9.6$ Jy) and Parkes 64-m at 5 GHz (4' : $S=2.3$ Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (1'1).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (34'': $S=3.4 \pm 0.4$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 75'': $S=4.4$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Kinugasa *et al.* 1998, PASJ, 50, 249. ASCA observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

G353.6–0.7

RA: $17^{\text{h}}32^{\text{m}}00^{\text{s}}$
Dec: $-34^{\circ}44'$

1-GHz flux/Jy: 2.5?
Spectral index: ?

Size/arcmin: 30
Type: S

Radio: Shell, brighter to S.

X-ray: Detected.

References:

Tian *et al.* 2008, ApJ, 679, L85. SGPS at 1.4 GHz ($100''$), plus 843 MHz ($43''$) and X-ray observations.

G353.9–2.0

RA: $17^{\text{h}}38^{\text{m}}55^{\text{s}}$
Dec: $-35^{\circ}11'$

1-GHz flux/Jy: 1?
Spectral index: 0.5?

Size/arcmin: 13
Type: S

Radio: Shell, with central double source.

References:

Green 2001, MNRAS, 326, 283. VLA at 327 MHz ($3.0' \times 2.7'$) and 1.4 GHz ($42'' \times 36''$), plus 8.4 GHz ($8.4'' \times 6.1''$) of central source only.

G354.1+0.1

RA: $17^{\text{h}}30^{\text{m}}28^{\text{s}}$
Dec: $-33^{\circ}46'$

1-GHz flux/Jy: ?
Spectral index: varies

Size/arcmin: 15×3 ?
Type: C?

Is this a SNR?

Radio: Elongated N–S.

Point sources: Pulsar at S tip.

References:

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz ($47'' \times 99''$) and 1.4 GHz ($8.8'' \times 21''$).

G354.8–0.8

RA: $17^{\text{h}}36^{\text{m}}00^{\text{s}}$
Dec: $-33^{\circ}42'$

1-GHz flux/Jy: 2.8?
Spectral index: ?

Size/arcmin: 19
Type: S

Radio: Distorted shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 78''$: $S=3.1$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G355.4+0.7

RA: $17^{\text{h}}31^{\text{m}}20^{\text{s}}$
Dec: $-32^{\circ}26'$

1-GHz flux/Jy: 5?
Spectral index: ?

Size/arcmin: 25
Type: S

Radio: Faint, incomplete shell.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 77''$).

Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1': S=8.9 \pm 1.3$ Jy).

G355.6–0.0

RA: 17^h35^m16^s
Dec: −32°38'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 8 × 6
Type: S

Radio: Well defined shell.

X-ray: Detected.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43'' × 80'': $S=2.6$ Jy).
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1': $S=3.3 \pm 0.5$ Jy).

G355.9–2.5

RA: 17^h45^m53^s
Dec: −33°43'

1-GHz flux/Jy: 8
Spectral index: 0.5

Size/arcmin: 13
Type: S

Radio: Distorted shell, brightest to SE.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=12.3$ Jy) and Parkes 64-m at 5 GHz (4': $S=3.4$ Jy).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (32'' × 34'': $S=5.0 \pm 0.3$ Jy).
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 77'').
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G356.2+4.5

RA: 17^h19^m00^s
Dec: −29°40'

1-GHz flux/Jy: 4
Spectral index: 0.7

Size/arcmin: 25
Type: S

Has been called G356.2+4.4.

Radio: Faint shell.

References:

Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (3' × 1.5': $S=8.1 \pm 1.7$ Jy), and NVSS at 1.4 GHz.

G356.3–0.3

RA: 17^h37^m56^s
Dec: −32°16'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 11 × 7
Type: S

Radio: Elongated shell, brighter in N.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43'' × 81'': $S=2.6$ Jy).
Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (0'.8 × 1'.7: $S=5.7 \pm 0.2$).

G356.3–1.5

RA: 17^h42^m35^s
Dec: −32°52'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 20 × 15
Type: S

Radio: Double arc.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43'' × 79'': $S=2.8$ Jy).

G357.7–0.1

RA: 17^h40^m29^s
Dec: −30°58'

1-GHz flux/Jy: 37
Spectral index: 0.4

MSH 17–39

Size/arcmin: 8 × 3?
Type: ?

Has been suggested that this is not a SNR.

Radio: Multiple arcs and filaments, with compact HII region at W edge.

X-ray: Detected.

Distance: HI absorption suggests beyond Galactic Centre.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ($3' : S = 54.2$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 18.5$ Jy).
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ($4'.4 : S = 14.6$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'.6$).
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz ($50''$).
 Weiler & Panagia 1980, A&A, 90, 269. Effelsberg 100-m at 9 GHz ($1'.5$) (private communication from Baker).
 Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz ($3''.8 \times 10''.9$) and 5 GHz ($12'' \times 26''$).
 Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 GHz and 5 GHz.
 Helfand & Becker 1985, Nature, 313, 118. Suggest it is not a SNR.
 Shaver *et al.* 1985, A&A, 147, L23. Observations of peripheral compact source.
 Caswell *et al.* 1989, PASA, 8, 184. MOST at 843 MHz ($43'' \times 83''$).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 84''$).
 Stewart *et al.* 1994, ApJ, 432, L39. ATCA at 4.79 and 5.84 GHz ($12'' \times 22''$) and Effelsberg 100-m at 10.6 GHz ($1'$), including polarisation.
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz ($43'' \times 24''$).
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Brogan & Goss 2003, AJ, 125, 272. VLA at 1.4 GHz ($13''.6 \times 11''.4$), including HI, and 8.3 GHz ($14''.3 \times 6''.8$) recombination line observation of HII region.
 Gaensler *et al.* 2003, ApJ, 594, L35. Chandra detection.
 Lazendic *et al.* 2003, AN, 324 (No S1), 157. Molecular line observations.
 Burton *et al.* 2004, MNRAS, 348, 638. IR and radio observations of HII region.
 Lazendic *et al.* 2004, MNRAS, 354, 393. IF and molecular line observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G357.7+0.3

RA: 17^h38^m35^s
Dec: −30°44'

1-GHz flux/Jy: 10
Spectral index: 0.4?

Size/arcmin: 24
Type: S

Radio: Non-thermal shell in complex region.

References:

- Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz ($4'.3 : S = 7 \pm 1.5$ Jy), $S_{5\text{ GHz}} = 5.5 \pm 1.5$ Jy from surveys.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 84''$).
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G358.0+3.8

RA: 17^h26^m00^s
Dec: −28°36'

1-GHz flux/Jy: 1.5?
Spectral index: ?

Size/arcmin: 38
Type: S

Radio: Faint shell.

References:

- Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz ($2'.2 \times 1'.3 : S = 2.5 \pm 1.3$ Jy), and NVSS at 1.4 GHz.

G358.1+0.1

RA: 17^h37^m00^s
Dec: −29°59'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 20
Type: S

Radio: Faint shell.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 77'').
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1': $S=6.0 \pm 2.5$ Jy).

G358.5−0.9

RA: 17^h46^m10^s
Dec: −30°40'

1-GHz flux/Jy: 4?
Spectral index: ?

Size/arcmin: 17
Type: S

Radio: Shell, brighter to NE.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 77'').
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1': $S=8.0 \pm 2.5$ Jy).

G359.0−0.9

RA: 17^h46^m50^s
Dec: −30°16'

1-GHz flux/Jy: 23
Spectral index: 0.5

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

X-ray: Partial shell.

References:

Reich *et al.* 1988, in SNRISM, p293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 86'').
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.4 × 12''.8) of part.

G359.1−0.5

RA: 17^h45^m30^s
Dec: −29°57'

1-GHz flux/Jy: 14
Spectral index: 0.4?

Size/arcmin: 24
Type: S

Radio: Non-thermal shell in complex region, crossed by the ‘snake’.

X-ray: Centrally brightened.

Point sources: Several compact radio sources near centre, OH masers around edge.

References:

Downes *et al.* 1979, A&AS, 35, 1. From observations by Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6: $S=13$ Jy).
 Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz (4'.3: $S=10 \pm 1.5$ Jy) and 4.8 GHz (2'.4: $S=8.1 \pm 0.5$ Jy).
 Uchida *et al.* 1992, ApJ, 398, 128. VLA at 1.5 GHz (11'' × 10''), and observations of nearby molecular material.
 Uchida *et al.* 1992, AJ, 104, 1533. VLA at 1.4 GHz.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43'' × 85'').
 Yusef-Zadeh *et al.* 1995, Science, 270, 1801. VLA at 1.4 GHz (33'' × 31''), and 1.7 GHz for OH survey.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43'' × 24'').
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.
 Lazendic *et al.* 2002, MNRAS, 331, 537. Observations of shocked molecular gas where the ‘snake’ crosses the remnant.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.4 × 12''.8).
 Aharonian *et al.* 2008, A&A, 483, 509. γ -ray observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

G359.1+0.9

RA: 17^h39^m36^s
Dec: −29°11'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 12 × 11
Type: S

Radio: Shell, brightest in E.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43'' \times 88''$: $S=4.3$ Jy).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1'$: $S=4.3 \pm 1.0$ Jy).
 Law *et al.* 2008, ApJS, 177, 515. VLA at 1.4 GHz ($10''.9 \times 15''.9$: $S=1.3 \pm 0.5$ Jy).