

**A Catalogue of
Galactic Supernova Remnants
(2009 March version)**

D. A. Green

*Cavendish Laboratory
19 J. J. Thomson Avenue
Cambridge CB3 0HE
UNITED KINGDOM*

E-mail: D.A.Green@mrao.cam.ac.uk

Please cite the following paper for the summary data from this catalogue:

- Green D. A., 2009, *Bulletin of the Astronomical Society of India*, 37, 45.

If you make use of the detailed version of the catalogue, then please also cite:

- Green D. A., 2009, 'A *Catalogue of Galactic Supernova Remnants (2009 March version)*', Cavendish Laboratory, Cambridge, UK (available on the World-Wide-Web at "<http://www.mrao.cam.ac.uk/surveys/snrs/>").

1. The Catalogue Format

This catalogue of Galactic supernova remnants (SNRs) is an updated version of those presented in detail in Green (1984, 1988) and in summary form in Green (1991, 1996, 2004) – hereafter Versions I, II, III, IV and V respectively – and on the World-Wide-Web, in versions of 1995 July, 1996 August, 1998 September, 2000 August, 2001 December, 2004 January and 2006 April. (Version IV, although published in 1996, was produced in 1993, and a detailed version of this was made available on the World-Wide-Web in 1993 November. The summary data from the 2001 December version of the catalogue was also published as an Appendix in Stephenson & Green 2002.)

This, the 2009 March version of the catalogue, contains 274 SNRs (which is 9 more than in the previous, 2006 April, version: 10 new remnants have been added, and 1 has been removed, see below), with over a thousand references in the detailed listings, plus notes on many possible or probable remnants.

For each remnant in the catalogue the following parameters are given.

- **Galactic Coordinates** of the source centroid, quoted to the nearest tenth of a degree as is conventional. (Note: in this catalogue additional leading zeros are not used.)
- **Other Names** that are commonly used for the remnant. These are given in parentheses if the remnant is only a part of the source. For some remnants, notably the Crab Nebula, not all common names are given.
- **Right Ascension** and **Declination** of the source centroid. The accuracy of the quoted values depends on the size of the remnant; for small remnants they are to the nearest few seconds of time and the nearest minute of arc respectively, whereas for larger remnants they are rounded to coarser values, but are in every case sufficient to specify a point within the boundary of the remnant. These coordinates are usually deduced from radio maps rather than from X-ray or optical observations, and are for J2000.0.

- **Angular Size** of the remnant, in arcminutes, usually taken from the highest resolution radio map available. The boundary of most remnants approximates reasonably well to a circle or an ellipse. A single value is quoted for the angular size of the more nearly circular remnants, which is the diameter of a circle with an area equal to that of the remnant. For elongated remnants the product of two values is quoted, and these are the major and minor axes of the remnant boundary modelled as an ellipse. In a few cases an ellipse is not a satisfactory description of the boundary of the object (refer to the description of the individual object given in its catalogue entry), although an angular size is still quoted for information. For ‘filled-centre’ remnants the size quoted is for the largest extent of the observed radio emission, not, as at times has been used by others, the half-width of the centrally brightened peak.
- **Flux Density** of the remnant at 1 GHz in jansky. This is *not* a measured value, but is deduced from the observed radio-frequency spectrum of the source. The frequency of 1 GHz is chosen because flux density measurements at frequencies both above and below this value are usually available.
- **Spectral Index** of the integrated radio emission from the remnant, α (here defined in the sense, $S \propto \nu^{-\alpha}$, where S is the flux density at a frequency ν), either a value that is quoted in the literature, or one deduced from the available integrated flux densities of the remnant. For several SNRs a simple power law is not adequate to describe their radio spectra, either because there is evidence that the integrated spectrum is curved or the spectral index varies across the face of the remnant. In these cases the spectral index is given as ‘varies’ (refer to the description of the remnant and appropriate references in the detailed catalogue entry for more information). In some cases, for example where the remnant is highly confused with thermal emission, the spectral index is given as ‘?’ since no value can be deduced with any confidence.
- **Type** of the SNR: ‘S’ or ‘F’ if the remnant shows a ‘shell’ or ‘filled-centre’ structure, or ‘C’ if it shows ‘composite’ (or ‘combination’) radio structure with a combination of shell and filled-centre characteristics; or ‘S?’, ‘F?’ or ‘C?’, respectively, if there is some uncertainty; or ‘?’ in several cases where an object is conventionally regarded as an SNR even though its nature is poorly known or not well-understood. Until recently only a few remnants were classified as composite remnants, as available observations were only able to identify the more obvious pulsar-powered, flatter radio spectrum filled-centre components within shells. However, in recent years improved observations – particularly in X-rays with the *Chandra* satellite – have identified many faint, pulsar powered nebulae in what until then had been identified as pure shell remnants. (Note: the term ‘composite’ has been used in a different sense, by some authors, to describe SNRs with shell radio and centrally-brightened X-ray morphologies. An alternative term used to describe such remnants is ‘mixed morphology’, see Rho & Petre 1998.)

In the detailed listings, for each remnant, notes on a variety of topics are given. First, it is noted if other Galactic coordinates have at times been used to label it (usually before good observations have revealed the full extent of the object), if the SNR is thought to be the remnant of a historical SN, or if the nature of the source as an SNR has been questioned (in which case an appropriate reference is usually given later in the entry). Brief descriptions of the remnant from the available radio, optical and X-ray observations as applicable are then given, together with notes on available distance determinations, and any point sources or pulsars in or near the object (although they may not necessarily be related to the remnant). Finally, appropriate references to observations are given for each remnant, complete with journal, volume, page, and a short description of what information each paper contains (for radio observations these include the telescopes used, the observing frequencies and resolutions, together with any flux density determinations). These references are *not* complete, but cover representative and recent observations of the remnant – up to the first the end of 2008 in this version of the catalogue – and they should themselves include references to earlier work. The references do not generally include large observational surveys – of particular interest in this respect are: the Effelsberg 100-m survey at 2.7 GHz of the Galactic plane $358^\circ < l < 240^\circ$, $|b| < 5^\circ$ by Reich *et al.* (1990) and Fürst *et al.* (1990a); reviews of the radio spectra of some SNRs by Kassim (1989), Kovalenko, Pynzar’ & Udal’tsov (1994) and Trushkin (1998); the Parkes 64-m survey at 2.4 GHz of the Galactic plane $238^\circ < l < 365^\circ$, $|b| < 5^\circ$ by Duncan *et al.* (1995) and Duncan *et al.* (1997); the Molonglo Galactic plane survey at 843 MHz of $245^\circ < l < 355^\circ$, $|b| < 1.5^\circ$ by Green *et al.* (1999); the survey of $345^\circ < l < 255^\circ$, $|b| < 5^\circ$ at 8.35 and 14.35 GHz by Langston *et al.* (2000); MAGPIS, see White, Becker & Helfand (2005) and Helfand *et al.* (2006); the VLA Galactic Plane Survey, see Stil *et al.* (2006); surveys of IRAS observations of SNRs and their immediate surroundings by Arendt (1989) and by Saken, Fesen & Shull (1992); the survey of HI emission towards SNRs by Koo & Heiles (1991); the *SPITZER* survey of inner galaxy SNRs by Reach *et al.* (2006); and the catalogue by Fesen & Hurford (1996) of UV/optical/infra-red lines identified in SNRs.

A summary of the data available for all 274 remnants in the catalogue is given in Table I. The other names for SNRs are listed in Table II, and the abbreviations for journals, proceedings and telescopes are listed in Table III. The detailed listings for each SNR are given in Table IV.

2. Revisions and Notes

2.1 Objects no longer thought to be SNRs

The following objects, which were listed in Version I of the catalogue were removed because they were no longer thought to be remnants, or were poorly observed (see Version II for references and further details): G2.4+1.4 (see also Gray 1994a; Goss & Lozinskaya 1995; Polcaro *et al.* 1995), G41.9–4.1 (=CTB 73, PKS 1920+06), G47.6+6.1 (=CTB 63), G53.9+0.3 (part of HC40), G93.4+1.8 (=NRAO 655), G123.2+2.9, G194.7+0.4 (the Origen Loop), G287.8–0.5 (see below), G322.3–1.2 (=Kes 24) and G343.0–6.0 (but see below). G358.4–1.9, which was listed in Version IV of the catalogue, was removed, as following the discussion of Gray (1994a), as it is not clear that this is a SNR. G240.9–0.9, G299.0+0.2 and G328.0+0.3, which were listed in 1995 July version of the catalogue, were removed from the 1996 August version, following the improved observations of Duncan *et al.* (1996) and Whiteoak & Green (1996). For the 1998 September revision of the catalogue G350.0–1.8 was incorporated into G350.0–2.0, and G337.0–0.1 refers to a smaller remnant than that previously catalogued with the same name. G112.0+1.2, G117.4+5.0, G152.2–1.2 and G211.7–1.1 – which were reported as SNRs by Bonsignori-Facondi & Tomasi (1979) – were removed from the 2001 December version of the catalogue, as the first three of these are not confirmed as SNRs from the ongoing *Canadian Galactic Plane Survey* (Roland Kothes, private communication; but see below for further discussion of another proposed remnant, G213.0–0.6). G10.0–0.3, which was regarded as a remnant – possibly associated with a soft-gamma repeater – was removed from the 2004 January version of the catalogue, as it is now thought to be radio nebula powered by a stellar wind (see Gaensler *et al.* 2001, Corbel & Eikenberry 2004, and references therein). G166.2+2.5 (=OA 184) was removed from the 2006 April version of the catalogue, as it was identified as an HII region by Foster *et al.* (2006).

G84.9+0.5 was removed from this version of the catalogue, as it was identified as an HII region by Foster *et al.* (2007) (see also Kothes *et al.* 2006).

The following objects, which have been reported as SNRs, but have not been included in any of the versions of the SNR catalogue, have subsequently been shown not to be SNRs.

- G70.7+1.2, which was reported as a SNR by Reich *et al.* (1985), but this has not been confirmed by later observations (see Green 1986; de Muizon *et al.* 1988; Becker & Fesen 1988; Caswell 1988; Bally *et al.* 1989; Phillips, Onello & Kulkarni 1993; Onello *et al.* 1995; Cameron & Kulkarni 2007).
- G81.6+1.0 a possible SNR in W75 reported by Ward-Thompson & Robson (1991). From the published data (see the observations in Wendker, Higgs & Landecker 1991) it was noted in Version IV of the catalogue that this is thermal source not a SNR, because of its thermal radio spectrum, and high infrared-to-radio emission (see also the subsequent discussion by Wendker *et al.* 1993).
- Green & Gull (1984) suggested G227.1+1.0 as a very young SNR, but subsequent observations (Channan *et al.* 1986; Green & Gull 1986) have shown that this is most likely an extragalactic source, not an SNR.
- A candidate SNR, G274.7–2.8, identified by Helfand & Channan (1989), has been shown not to be a SNR by Caswell & Stewart (1991).
- G159.6–18.5, was suggested as a SN by Pauls & Schwartz (1989), from IRAS and other observations, but is probably an HII region (see Andersson *et al.* 2000).
- G25.5+0.2, which was reported as a very young SNR by Cowan *et al.* (1989), although this identification was not certain (see White & Becker 1990; Green 1990; Zijlstra 1991). Sramek *et al.* (1992) report the detection of recombination lines from this source (also see Subrahmanyam *et al.* 1993). Becklin *et al.* (1994) identify G25.5+0.2 as a ring nebula around a luminous blue star. See also Clark, Steele & Langer (2000), and Phillips & Ramos-Larios (2008) who identified G25.5+0.2 as a possible symbiotic outflow.
- Several of the possible SNRs listed by Gorham (1990) – following up SNR candidates suggested by Kassim (1988) – have been shown not to be SNRs by Gorham, Kulkarni & Prince (1993).
- G203.2–12.3, a optical ring about 3 arcmin in diameter, was reported as a possible SNR by Winkler & Reipurth (1992), but was shown to be a Herbig–Haro object (HH 311) by Reipurth, Bally & Devine (1997).

- G359.87+0.18 was reported as a possible young SNR near the Galactic Centre by Yusef-Zadeh, Cotton & Reynolds (1998), but was shown to be a radio galaxy by Lazio *et al.* (1999).
- G104.7+2.8, a possible SNR suggested by Green & Joncas (1994), which instead appears to be an HII region, based on the improved observations by Kothes *et al.* (2006).
- G106.6+2.9, a small remnant proposed by Halpern *et al.* (2001), is incorporated into the larger catalogued remnant G106.3+2.7.
- Leahy, Tian & Wang (2008) proposed that a large radio shell, G53.9+0.2, as a possible SNR. As noted above, this feature was included, as G53.9+0.3 (part of HC40), in Version I of the catalogue, but was subsequently removed, following the discussions of Caswell (1985) who concluded it was a thermal source (see also Velusamy, Goss & Arnal 1986) – results which Leahy *et al.* ignored.

Some entries in the catalogue have been renamed, due to improved observations revealing a larger true extent for the object (previously G5.3–1.0 is now G5.4–1.2; G193.3–1.5 is now G192.8–1.1; G308.7+0.0 is now incorporated into G308.8–0.1). G337.0–0.1 now refers to a small (1.5 arcmin) remnant, rather than larger supposed remnant at this position (see Sarma *et al.* 1997), and G350.0–2.0 now incorporates the previously catalogued G350.0–1.8, based on the improved observations of Gaensler (1998).

2.2 New SNRs

The following remnants were added to Version II of the catalogue: G0.9+0.1, G1.9+0.3, G5.9+3.1, G6.4+4.0, G8.7–0.1, G16.8–1.1, G18.9–1.1, G20.0–0.2, G27.8+0.6, G30.7+1.0, G31.5–0.6, G36.6–0.7, G42.8+0.6, G45.7–0.4, G54.1+0.3, G73.9+0.9, G179.0+2.6, G312.4–0.4, G357.7+0.3 and G359.1–0.5.

The following remnants were added to Version III of the catalogue: G4.2–3.5, G5.2–2.6, G6.1+1.2, G8.7–5.0, G13.5+0.2, G15.1–1.6, G16.7+0.1, G17.4–2.3, G17.8–2.6, G30.7–2.0, G36.6+2.6, G43.9+1.6, G59.8+1.2, G65.1+0.6, G68.6–1.2, G69.7+1.0, G279.0+1.1, G284.3–1.8 (=MSH 10–53), G358.4–1.9 and G359.0–0.9 (although, as noted above, G358.4–1.9 was subsequently removed).

The following remnants were added to Version IV of the catalogue: G59.5+0.1, G67.7+1.8, G84.9+0.5, G156.2+5.7, G318.9+0.4, G322.5–0.1, G343.1–2.3 and G348.5–0.0 (although, as noted above, G84.9+0.5 was subsequently removed).

The following remnants were added to 1995 July version of the catalogue: G1.0–0.1, G1.4–0.1, G3.7–0.2, G3.8+0.3, G28.8+1.5, G76.9+1.0, G272.2–3.2, G341.2+0.9, G354.1+0.1, G355.6–0.0, G356.3–0.3, G356.3–1.5 and G359.1+0.9.

The following remnants were added to the 1996 August version of the catalogue: G13.3–1.3, G286.5–1.2, G289.7–0.3, G294.1–0.0, G299.2–2.9, G299.6–0.5, G301.4–1.0, G308.1–0.7, G310.6–0.3, G310.8–0.4, G315.9–0.0, G317.3–0.2, G318.2+0.1, G320.6–1.6, G321.9–1.1, G327.4+1.0, G329.7+0.4, G342.1+0.9, G343.1–0.7, G345.7–0.2, G349.2–0.1, G351.7+0.8, G351.9–0.9 and G354.8–0.8.

The following remnants were added to the 1998 September version of the catalogue: G0.3+0.0, G32.1–0.9, G55.0+0.3, G63.7+1.1 and G182.4+4.3.

The following remnants were added to the 2000 August version of the catalogue: G7.0–0.1, G16.2–2.7, G29.6+0.1, G266.2–1.2 and G347.3–0.5.

The following remnants were added to the 2001 December version of the catalogue: G4.8+6.2, G28.6–0.1, G85.4+0.7, G85.9–0.6, G106.3+2.7, G292.2–0.5, G343.0–6.0, G353.9–2.0, G356.2+4.5 and G358.0+3.8.

G312.5–3.0 was added to the 2004 January version of the catalogue.

The following remnants were added to the 2006 April version of the catalogue: G5.5+0.3, G6.1+0.5, G6.5–0.4, G7.2+0.2, G8.3–0.0, G8.9+0.4, G9.7–0.0, G9.9–0.8, G10.5–0.0, G11.0–0.0, G11.1–0.7, G11.1–1.0, G11.1+0.1, G11.8–0.2, G12.2+0.3, G12.5+0.2, G12.7–0.0, G12.8–0.0, G14.1–0.1, G14.3+0.1, G15.4+0.1, G16.0–0.5, G16.4–0.5, G17.0–0.0, G17.4–0.1, G18.1–0.1, G18.6–0.2, G19.1+0.2, G20.4+0.1, G21.0–0.4, G21.5–0.1, G32.4+0.1, G96.0+2.0, G113.0+0.2 and G337.2+0.1.

The following remnants have been added to this version of the catalogue.

- G83.0–0.3, which had been suggested as a SNR by Taylor, Wallace & Goss (1992), and is now included in the catalogue following improved observations by Kothes *et al.* (2006) which confirm its nature.
- G108.2–0.6 identified by Tian, Leahy & Foster (2007).
- G315.1+2.7 and G332.5–5.6 – which had been suggested as SNR candidates by Duncan *et al.* (1995, 1997) – have been confirmed as SNRs by further observations reported by Stupar, Parker & Filipović (2007) and Reynoso & Green (2007) respectively.

- G327.2–0.1 a shell remnant found around the magnetar 1E 1547.0–5408, see Gelfand & Gaensler (2007).
- G350.1–0.3 was listed in early versions of the catalogue, but was removed (in Version III), as observations by Salter *et al.* (1986) did not allow a clear identification of the nature of this source. Recently Gaensler *et al.* (2008) have presented new observations of this source, including HI absorption observations which indicate it is Galactic, which – along with other observations, including its X-ray emission – support an SNR identification. However, its structure at radio wavelengths is rather different from other known remnants.
- G353.6–0.7 a shell remnant associated with HESS J1731–347 identified by Tian *et al.* (2008).
- Three sources – G355.4+0.7, G358.1+0.1, G358.5–0.9 – which had been identified as possible SNRs by Gray (1994b), have now been added to the catalogue, following further observations by Roy & Bhatnagar (2006) which confirm their nature.

2.3 Possible and probable SNRs not listed in the catalogue

The following are possible or probable SNRs for which further observations are required to confirm their nature or parameters, or for which observations are not yet in the published literature.

2.3.1 Radio

- G35.6–0.4 was listed in some early catalogues of Galactic SNRs (e.g. Milne 1970), but was identified as a likely thermal source instead by Caswell & Clark (1975). However, from VGPS and other data, this may in fact be a SNR (Green 2009).
- A possible SNR near the Galactic centre reported by Ho *et al.* (1985) from radio observations (see also Coil & Ho 2000; Lu, Wang & Lang 2003; Senda, Murakami & Koyama 2003, and references therein).
- Gosachinskiĭ (1985) reported evidence for non-thermal radio emission, presumably from SNRs, associated with several bright, thermal Galactic sources. Some of these sources have been included in the catalogue, following improved observations (but also see Odegard 1986, who questions the reliability of some of Gosachinskiĭ’s results).
- G300.1+9.4, a possible SNR nearly 2° in diameter reported by Dubner, Colomb & Giacani (1986).
- Routledge & Vaneldik (1988) report a possible faint radio shell SNR nearly 2° in diameter, near the young pulsar PSR 1930+22 – see also Gómez-González & del Romero (1983), who report a smaller (about 40 arcmin) possible SNR (G57.1+1.7) associated with this pulsar, and see Caswell, Landecker & Feldman (1985) and Kovalenko (1989).
- Gorham (1990) lists many SNR candidates from the Clark Lake 30.9 MHz survey of the first quadrant, following Kassim (1988), although several have been shown not to be SNRs by Gorham, Kulkarni & Prince (1993). Gorham *et al.* do report a poorly defined possible remnant G41.4+1.2. See also Aharonian *et al.* (2008) for observations of γ - and X-ray emission possibly associated with one of the candidates (G44.6+0.1) listed by Gorham.
- Four possible remnants (G45.9–0.1, G71.6–0.5, G72.2–0.3 and G85.2–1.2) of the eleven reported by Taylor, Wallace & Goss (1992) from a radio survey of part of the Galactic plane (see also Kothes *et al.* 2006). (Five of the other possible SNRs reported by Taylor *et al.*, are included in the catalogue as G55.0+0.3, G59.5+0.1, G63.7+1.1, G76.9+1.0 and G83.0–0.2, following improved observations which have confirmed their nature.)
- G356.6+0.1, G357.1–0.2, G358.7+0.7, G359.2–1.1, G3.1–0.6 and G4.2+0.0, which are among the possible SNRs listed by Gray (1994b) from radio observations near the Galactic centre. See also Roy & Pramesh Rao (2002) who present additional observations of G356.3–0.3, G356.6+0.1, G357.1–0.2 and G3.1–0.6 which they consider as possible SNRs, and Bhatnagar (2002) for additional observations of G4.2+0.0 which appears to be a thermal source.
- Duncan *et al.* (1995) and Duncan *et al.* (1997) list several large-scale (1.5 to 10 degree), and smaller, low radio surface-brightness candidate SNRs from the Parkes 2.4-GHz survey of $270^\circ < l < 360^\circ$. Several of these candidates have been confirmed as SNRs by subsequent, improved observations, and are included in the catalogue. (See also Camilo *et al.* 2004a, who detected a young pulsar near one of these candidate SNRs, G309.8–2.6, and Russeil *et al.* 2005, who detected optical filaments from another).

- Whiteoak & Green (1996), from their radio survey of much of the southern Galactic plane, list several possible SNRs (G308.4–1.4, G317.5+0.9, G319.9–0.7, G320.6–0.9, G322.7+0.1, G322.9–0.0, G323.2–1.0, G324.1+0.1, G325.0–0.3, G331.8–0.0, G337.2+0.1, G339.6–0.6, G345.1+0.2, G345.1–0.2, and G348.8+1.1). See also Schaudel *et al.* (2002) and Hui & Becker (2007) for X-ray observations of G308.3–1.4 and G319.9–0.7 respectively.
- Several candidate SNRs reported by Combi & Romero (1998), Combi, Romero & Arnal (1998), Combi, Romero & Benaglia (1998), Punsly *et al.* (2000) and Combi *et al.* (2001).
- A possible SNR, near $l = 313^\circ$, which is close to an unidentified Galactic plane γ -ray source (see Roberts *et al.* 1999), and to a pulsar (Roberts, Romani & Johnston 2001). See also Aharonian *et al.* (2006a).
- G359.07–0.02, a possible SNR noted by LaRosa *et al.* (2000).
- A possible SNR near G6.4–0.1 (=W28) noted by Yusef-Zadeh *et al.* (2000).
- Gaensler *et al.* (2000), in a search for pulsar wind nebulae, found a small shell of radio emission near PSR B1356–60 – which they designate G311.28+1.09 – which may be a supernova remnant.
- A possible SNR, G328.6–0.0, noted by McClure-Griffiths *et al.* (2001) in the test region of the *Southern Galactic Plane Survey*.
- G346.5–0.1, an arc of radio emission observed by Gaensler *et al.* (2001), which is potentially part of a SNR, but requires further observations to confirm its nature.
- Giacani *et al.* (2001) presented observations of a pulsar wind nebula around PSR J1709–4428, which may be part of the catalogued remnant G343.1–2.3, or may represent another object.
- Several possible SNRs reported by Trushkin (2001), which were identified from Galactic radio surveys (one of which, G6.1+0.5, is included in the catalogue, due to improved subsequent observations).
- Two possible SNRs (G336.1–0.2 and G352.2–0.1) discussed briefly by Manchester *et al.* (2002).
- G282.8–1.2, a possible young SNR noted by Misanovic, Cram & Green (2002).
- Three possible remnants – G41.5+0.4, G42.0–0.0 and G43.5+0.6 – identified by Kaplan *et al.* (2002).
- Two faint SNR candidates shown in Reich (2002).
- A possible faint remnant, G213.0–0.6, noted by Reich, Zhang & Fürst (2003), which is not well defined by current observations (this incorporates one of the faint remnants which was proposed by Bonsignori-Facondi & Tomasi 1979, see above).
- G107.5–1.5, a probable remnant identified at by Kothes (2003), but the full extent of which is not well defined at present (see also Kothes *et al.* 2006).
- Zhang (2003) identified four candidate SNRs from radio surveys. One of these – called G41.9+0.04 by Zhang – is close to one of the possible remnant by Kaplan *et al.* (2002), see above. A second – G74.8+0.63 – which Zhang identified as a possible remnant partly on the basis of its non-thermal radio spectrum, actually has a flat, thermal radio spectrum, and has long been identified as an HII region (e.g. Weiler & Shaver 1978; Pineault & Chastenay 1990). Another of the sources – G47.8+2.03 – also may have a thermal radio spectrum, given its published 2.7-GHz flux density (Fürst *et al.* 1990b).
- Brogan *et al.* (2006) identify 35 new SNRs in the region $4^\circ 5' < l < 22^\circ$, $|b| < 1^\circ 25'$, of which the 31 which are classed as ‘I’ or ‘II’ (i.e. those thought to be very or fairly confidently identified as SNRs) were included in the 2006 version of the catalogue. Four other possible SNRs – labelled G5.71–0.08, G6.31+0.54, G15.51–0.15 and G19.13+0.90 – which comprise Brogan *et al.*’s class ‘III’, are not included in the catalogue, as further observations are required to confirm their nature and better define their parameters.
- Helfand *et al.* (2006) list many SNR candidates in the region $5^\circ < l < 32^\circ$, $|b| < 0^\circ 8'$ from MAGPIS. Many of these correspond to sources in Brogan *et al.*, and several are included in the catalogue, with the others requiring further observations.
- A likely shell SNR G64.5+0.9, noted by Tian & Leahy (2006), see also Hurley-Walker *et al.* (2009).
- Martí *et al.* (2007), report extended radio emission near the X-ray source KS 1741–293 near the Galactic centre which may be a SNR (see also Cherepashchuk 1994).
- A poorly defined possible SNR, near $l = 151^\circ$, $b = 3^\circ$ has been reported by Kerton, Murphy & Patterson (2007).
- Roberts & Brogan (2008) propose a new SNR, G8.7–1.7, from non-thermal radio emission near an pulsar wind nebula, although currently the extent of the remnant is not well defined.

2.3.2 UV/Optical/Infra-red

- A possible SNR overlapping G296.1–0.5, identified from optical (and X-ray) observations by Hutchings, Crampton & Cowley (1981).
- Winkler *et al.* (1989) report a possible small (4 arcmin) SNR within the Puppis A remnant, from optical observations. This has not been detected at radio wavelengths (see Dubner *et al.* 1991).
- A possible SNR (G32.1+0.1) reported from optical spectroscopy by Thompson, Djorgovski & de Carvalho (1991), following up radio and infrared observations of Jones, Garwood & Dickey (1988), although this appears to have a thermal radio spectrum.
- G75.5+2.4, a possible large (about 2°) old SNR in Cygnus suggested by Nichols-Bohlin & Fesen (1993) from infra-red and optical observations (see also Dewdney & Lozinskaya 1994; Marston 1996; Esipov *et al.* 1996; Kothes *et al.* 2006).
- A possible optical SNR (G247.8+4.9) noted by Weinberger (1995), which may be Balmer dominated (see also Weinberger *et al.* 1998 and Zanin & Kerber 2000).
- An optical shell around the Coalsack Nebula (near $l = 300^\circ$, $b = 0^\circ$) identified by Walker & Zealey (1998). This coincides with one of the large possible SNRs suggested by Duncan *et al.* (1995), from radio observations.
- Two possible SNRs, G340.5+0.7 and G342.1+0.1, identified by Walker, Zealey & Parker (2001) from filaments seen in H α survey observations.
- A probable SNR which was identified by Bally & Reipurth (2001) – which they label as G110.3+11.3 – from optical filaments (and which is also associated with a large HI and CO cavity, and soft X-ray enhancement).
- A possible remnant, near $l = 70^\circ$, $b = 2^\circ$ noted by Mavromatakis & Strom (2002), for which Kothes *et al.* (2006) do not find any radio counterpart.
- Optical filaments in Pegausus (Boumis *et al.* 2002) which suggest one or more possible SNRs.
- A possible remnant identified from optical filaments to the NE of the known SNR G116.5+1.1, as observed by Mavromatakis *et al.* (2005).
- A suggested small, young remnant observed by Spitzer (Morris *et al.* 2006).
- Russell *et al.* (2007) report a small (about 7 arcmin in extent) optical ring, which is very faint at radio wavelengths, which is just to the NW of Cyg X-1, which may be a SNR if it is not associated with Cyg X-1 (see also Gallo *et al.* 2005).
- Stupar, Parker & Filipović (2008) report several SNRs identified from H α observations, several of which correspond to SNR candidates first suggested by Duncan *et al.* (1995, 1997) from radio observations. The full extent of most of these are not well defined, but two are currently included in the main catalogue (G315.1+2.7, and G332.5–5.6).

2.3.3 X-ray/ γ -ray

- H1538–32 a large X-ray source in Lupus, near $l = 307^\circ$, $b = +20^\circ$ (Riegler, Agrawal & Gull 1980; see also Colomb, Dubner & Giacani 1984; Gahm *et al.* 1990) which is a possible old SNR;
- G189.6+3.3, a faint, possible SNR overlapping G189.1+3.0 (=IC443) identified by Asaoka & Aschenbach (1994) from ROSAT X-ray observations.
- G117.7+0.6, a faint shell of soft X-ray emission near G116.9+0.2 (=CTB 1), which contains a pulsar (Hailey & Craig 1995; see also Craig, Hailey & Pisarski 1997, Esposito *et al.* 2008 and Kothes *et al.* 2006).
- A possible SNR identified in X-rays around the pulsar B1828–13 (see Finley, Srinivasan & Park 1996).
- A possible, large SNR, G69.4+1.2, identified as an X-ray shell by Yoshita, Miyata & Tsunemi (1999, 2000). See also Mavromatakis, Boumis & Paleologou (2002) and Kothes *et al.* (2006).
- Possible SNRs identified in the ROSAT All-Sky Survey are discussed briefly by Schaudel *et al.* (2002).
- G0.570–0.018 a small ring of X-ray emission near the Galactic Centre, which has been proposed as a very young remnant by Senda, Murakami & Koyama (2002). See also Senda, Murakami & Koyama (2003), who identify other possible SNRs near the Galactic Centre from their X-ray emission, Renaud *et al.* (2006) and Mori *et al.* (2008).

- Two probable SNRs (G25.5+0.0 and G26.6–0.1) identified by Bamba *et al.* (2003) from their hard X-ray emission.
- Ueno *et al.* (2004) identify several candidate SNRs in the first quadrant from the ASCA Galactic Plane Survey (see also Yamuguchi *et al.* 2004). Two of these are included in the catalogue (as G28.6–0.1 and G32.4+0.1), as additional observations confirm their nature.
- A possible SNR identified from X-ray and γ -ray observations (Malizia *et al.* 2005).
- Cui & Konopelko (2006) identify an extended X-ray source near $l = 8^{\circ}4$, $b = +0^{\circ}1$.
- An excess of Fe X-ray line emission in Sgr B, near $l = 0^{\circ}61$, $b = 0^{\circ}01$ may be from a SNR (Koyama *et al.* 2007).
- Nobukawa *et al.* (2008) report a region of X-ray emission, G0.42–0.04, near the Galactic centre, which may be part of a SNR.

2.3.4 Other

- G287.8–0.5, which is associated with η Carinae, was listed in Version I as a SNR, but was removed from the catalogue in Version II as its parameters are uncertain (see Jones 1973; Retallack 1984; Tateyama, Strauss & Kaufmann 1991; and the discussion in Version II).
- G359.2–0.8 (the ‘mouse’), near the Galactic centre, which has been suggested as being analogous to the central region of G69.0+2.7 (=CTB 80) by Predehl & Kulkarni (1995), i.e. a pulsar powered nebula (see also Camilo *et al.* 2002).

It should also be noted that: (a) some large radio continuum and HI loops in the Galactic plane (e.g. Berkhuijsen 1973) may be parts of very large, old SNRs, but they have not been included in the catalogue (see also Combi *et al.* 1995; Maciejewski *et al.* 1996; Kim & Koo 2000; Normandeau *et al.* 2000; Woermann, Gaylard & Otrupcek 2001; Stil & Irwin 2001; Uyaniker & Kothes 2002; Olano, Meschin & Niemela 2006), also see Koo, Kang & Salter (2006) and Kang & Koo (2007) who identify faint Galactic HI features at forbidden velocities as indicators of old, otherwise undetectable SNRs; (b) some large ($> 10^{\circ}$) regions of X-ray emission that are indicative of a SNR are not included in the catalogue (e.g. the Monogem ring, near $l = 203^{\circ}$, $b = +12^{\circ}$, see Nousek *et al.* 1981, Plucinsky *et al.* 1996, Thorsett *et al.* 2003, Amenomori *et al.* 2005, and references therein, plus Weinberger, Temporin & Stecklum 2006, for observations of optical filaments; in the Gum Nebula near $l = 250^{\circ}$, $b = 0^{\circ}$, see Leahy, Nousek & Garmire 1992, and also see Reynolds 1976, Dubner *et al.* 1992, Duncan *et al.* 1996, Reynoso & Dubner 1997, Heiles 1998; in Eridanus near $l = 200^{\circ}$, $b = -40^{\circ}$, see Naranan *et al.* 1976, Burrows *et al.* 1993, Snowden *et al.* 1995, Heiles 1998, Boumis *et al.* 2001, Ryu *et al.* 2006); a large approximately 24° diameter, X-ray and optical loop in Antlia, see McCullough, Fields & Pavlidou 2002, Shinn *et al.* 2007); (c) the distinction between filled-centre remnants and pulsar wind nebulae (PWNe) is not clear, and isolated, generally faint, pulsar wind nebulae are also not included in the catalogue. Kaspi, Roberts & Harding (2006) provide a catalogue of PWNe (see also <http://www.physics.mcgill.ca/~pulsar/pwnecat.html>, and Camilo *et al.* 2004b, Aharonian *et al.* 2005, Hessels *et al.* 2005, Aharonian *et al.* 2006b, Gonzalez *et al.* 2006, Wang, Lu & Gotthelf 2006, Aharonian *et al.* 2007, Hinton *et al.* 2007, Bhattacharyya 2008, Gotthelf & Halpern 2008, Munro *et al.* 2008).

2.4 Questionable SNRs listed in the catalogue

As noted in Versions II and IV of the catalogue, the following sources are listed as SNRs, although, as discussed in each case, the identifications are not certain: G5.4–1.2, G39.7–2.0 (=W50), G69.0+2.7 (=CTB 80), G318.9+0.4 and G357.7–0.1. The nature of G76.9+1.0 (an unusual radio source similar to G65.7+1.2), and of G354.1+0.1 (which may be similar to G357.7–0.1 (=MHS 17–39)) are also uncertain (see Landecker, Higgs & Wendker 1993 and Frail, Goss & Whiteoak 1994).

There are also some objects that have been identified as SNRs and are listed in the catalogue, although they have been barely resolved in the available observations, or are faint, and have not been well separated from confusing background or nearby thermal emission, and their identification as SNRs, or at least their parameters remain uncertain.

Acknowledgements

This research has made use of NASA’s Astrophysics Data System Bibliographic Services.

References

- Aharonian F. A. *et al.*, 2005, *A&A*, 442, L25.
Aharonian F. *et al.*, 2006a, *A&A*, 456, 245.
Aharonian F. *et al.*, 2006b, *A&A*, 460, 365.
Aharonian F. *et al.*, 2007, *A&A*, 472, 489.
Aharonian F. *et al.*, 2008, *A&A*, 484, 435.
Amenomori M. *et al.*, 2005, *ApJ*, 635.
Andersson B.-G., Wannier P. G., Moriarty-Schieven G. H. & Bakker E. J., 2000, *AJ*, 119, 1325.
Arendt R. G., 1989, *ApJS*, 70, 181.
Asaoka I. & Aschenbach B., 1994, *A&A*, 284, 573.
Bally J. & Reipurth B., 2001, *ApJ*, 552, L159.
Bally J. *et al.*, 1989, *ApJ*, 338, L65.
Bamba A., Ueno M., Koyama K. & Yamauchi S., 2003, *ApJ*, 589, 253.
Becker R. H. & Fesen R. A., 1988, *ApJ*, 334, L35.
Becklin E. E., Zuckerman B., McLean I. S. & Geballe T., 1994, *ApJ*, 430, 774.
Berkhuijsen E. M., 1973, *A&A*, 24, 143.
Bhatnagar S., 2002, *MNRAS*, 332, 1.
Bhattacharyya S., 2008, *MNRAS*, 391, L117.
Bonsignori-Facondi S. R. & Tomasi P., 1979, *A&A*, 77, 93.
Boumis P., Dickinson C., Meaburn J., Goudis C. D., Christopoulou P. E., López J. A., Bryce M. & Redman M. P., 2001, *MNRAS*, 320, 61.
Boumis P., Mavromatakis F., Paleologou E. V. & Becker W., 2002, *A&A*, 396, 225.
Brogan C. L., Gelfand J. D., Gaensler B. M., Kassim N. E. & Lazio T. J. W., 2006, *ApJ*, 639, L25.
Burrows D. N., Singh K. P., Nousek J. A., Garmire G. P. & Good J., 1993, *ApJ*, 406, 97.
Cameron P. B. & Kulkarni S. R., 2007, *ApJ*, 665, L135.
Camilo F., Manchester R. N., Gaensler B. M. & Lorimer D. R., 2002, *ApJ*, 579, L25.
Camilo F. *et al.*, 2004a, *ApJ*, 611, L25.
Camilo F., Gaensler B. M., Gotthelf E. V., Halpern J. P. & Manchester R. N., 2004b, *ApJ*, 616, 1118.
Caswell J. L., 1985, *AJ*, 90, 1224.
Caswell J. L., 1988, in *Supernovae and Supernova Remnants*, (IAU Colloquium 145), eds McCray R. & Wang Z., (Cambridge University Press), p269.
Caswell J. L. & Clark D. H., 1975, *AuJPA*, 37, 57.
Caswell J. L. & Stewart R. T., 1991, *PASA*, 9, 103.
Caswell J. L., Landecker T. L. & Feldman P. A., 1985, *AJ*, 90, 488.
Channan G. A., Helfand D. J., Spinrad H. & Ebnetter K., 1986, *Nature*, 320, 41.
Cherepashchuk A. M., Goranskij V. P., Karitskaya E. A., Nadjip A. E., Savage A., Shakura N. I., Sunyaev R. A. & Volchkov, A. A., 1994, *A&A*, 289, 419.
Clark S. J., Steele I. A. & Langer N., 2000, *ApJ*, 541, 67.
Coil A. L. & Ho P. T. P., 2000, *ApJ*, 533, 245.
Colomb F. R., Dubner G. M. & Giacani E. B., 1984, *A&A*, 130, 294.
Combi J. A. & Romero G. E., 1998, *A&AS*, 128, 423.
Combi J. A., Romero G. E. & Arnal E. M., 1998, *A&A*, 333, 298.
Combi J. A., Romero G. E. & Benaglia P., 1998, *A&A*, 333, L91.
Combi J. A., Testori J. C., Romero G. E. & Colomb F. R., 1995, *A&A*, 296, 514.
Combi J. A., Romero G. E., Benaglia P. & Jonas J. L., 2001, *A&A*, 366, 1047.
Corbel S. & Eikenberry S. S., 2004, *A&A*, 419, 191.
Cowan J. J., Ekers R. D., Goss W. M., Sramek R. A., Roberts D. A. & Branch D., 1989, *MNRAS*, 241, 613.
Craig W. W., Hailey C. J. & Pisarski R. L., 1997, *ApJ*, 488, 307.
Cui W. & Konopelko A., 2006, *ApJ*, 652, L109.
de Muizon M., Strom R. G., Oort M. J. A., Claas J. J. & Braun R., 1988, *A&A*, 193, 248.
Dewdney P. E. & Lozinskaya T. A., 1994, *AJ*, 108, 2212.
Dubner G. M., Colomb F. R. & Giacani E. B., 1986, *AJ*, 91, 343.
Dubner G. M., Braun R., Winkler P. F. & Goss 1991, *AJ*, 101, 1466.
Dubner G., Giacani E., Cappa de Nicolau C. & Reynoso E., 1992, *A&AS*, 96, 505.
Duncan A. R., Stewart R. T., Haynes R. F. & Jones K. L., 1995, *MNRAS*, 277, 36.
Duncan A. R., Stewart R. T., Haynes R. F. & Jones K. L., 1996, *MNRAS*, 280, 252.
Duncan A. R., Stewart R. T., Haynes R. F. & Jones K. L., 1997, *MNRAS*, 287, 722.
Esipov V. F., Lozinskaya T. A., Mel'nikov V. V., Pravdikova V. V., Sitnik T. G. & Nichol-Bohlin J., 1996, *ALet*, 22, 509.
Esposito P., de Luca A., Tiengo A., Paizis A., Mereghetti S. & Caraveo, P. A., 2008, *MNRAS*, 384, 225.
Fesen R. A. & Hurford A. P., 1996, *ApJS*, 106, 563.
Finley J. P., Srinivasan R. & Park S., 1996, *ApJ*, 466, 938.
Frail D. A., Goss W. M. & Whiteoak J. B. Z., 1994, *ApJ*, 437, 781.
Foster T., Kothes R., Sun X. H., Reich W. & Han J. L., 2006, *A&A*, 454, 517.
Foster T. J., Kothes R., Kerton C. R. & Arvidsson K., 2007, *ApJ*, 667, 248.
Fürst E., Reich W., Reich P. & Reif K., 1990a, *A&AS*, 85, 691.
Fürst E., Reich W., Reich P. & Reif K., 1990b, *A&AS*, 85, 805.

- Gaensler B. M., 1998, ApJ, 493, 781.
- Gaensler B. M., Stappers B. W., Frail D. A., Moffett D. A., Johnston S. & Chatterjee S., 2000, MNRAS, 318, 58.
- Gaensler B. M., Slane P. O., Gotthelf E. V. & Vasisht G., 2001, ApJ, 559, 963.
- Gaensler B. M., Tanna A., Slane P. O., Brogan C. L., Gelfand J. D., McClure-Griffiths N. M., Camilo F., Ng C.-Y. & Miller J. M., 2008, ApJ, 680, L37.
- Gahm G. F., Gebeyehu M., Lindgren M., Magnusson P., Mødigh P. & Nordh H. L., 1990, A&A, 228, 477.
- Gallo E., Fender R., Kaiser C., Russell D., Morganti R., Oosterloo T. & Heinz S., 2005, Nature, 436, 819.
- Gelfand J. D. & Gaensler B. M., 2007, ApJ, 667, 1111.
- Giacani E. B., Frail D. A., Goss W. M. & Vieytes M., 2001, AJ, 121, 3133.
- Gómez-González J. & del Romero A., 1983, A&A, 123, L5.
- Gonzalez M. E., Kaspi V. M., Pivovarov M. J. & Gaensler B. M., 2006, ApJ, 652, 569.
- Gorham P. W., 1990, ApJ, 364, 187.
- Gorham P. W., Kulkarni S. K. & Prince T. A., 1993, AJ, 105, 314.
- Gosachinskiĭ I. V., 1985, SvA, 29, 128.
- Goss W. M. & Lozinskaya T. A., 1995, ApJ, 439, 637.
- Gotthelf E. V. & Halpern J. P., 2008, ApJ, 681, 515.
- Gray A. D., 1994a, MNRAS, 270, 835.
- Gray A. D., 1994b, MNRAS, 270, 847.
- Green A. J., Cram L. E., Large M. I. & Ye T. S., 1999, ApJS, 122, 207.
- Green D. A., 1984, MNRAS, 209, 449. (Version I)
- Green D. A., 1986, MNRAS, 219, 39P.
- Green D. A., 1988, Ap&SS, 148, 3. (Version II)
- Green D. A., 1990, AJ, 100, 1241.
- Green D. A., 1991, PASP, 103, 209. (Version III)
- Green D. A., 1996, in *Supernovae and Supernova Remnants*, (IAU Colloquium 145), eds McCray R. & Wang Z., (Cambridge University Press), p.419. (Version IV)
- Green D. A., 2004, BASI, 32, 335. (Version V)
- Green D. A., 2009, MNRAS, in press.
- Green D. A. & Gull S. F., 1984, Nature, 312, 527.
- Green D. A. & Gull S. F., 1986, Nature, 320, 42.
- Green D. A. & Joncas G., 1994, A&AS, 104, 481.
- Hailey C. J. & Craig W. W., 1995, ApJ, 455, L151.
- Halpern J. P., Camilo F., Gotthelf E. V., Helfand D. J., Kramer M., Lyne A. G., Leighly K. M. & Eracleous M., 2001, ApJ, 552, L125.
- Heiles C., 1998, ApJ, 498, 689.
- Helfand D. J. & Channan G. A., 1989, AJ, 98, 1652.
- Helfand D. J., Becker R. H., White R. L., Fallon A. & Tuttle S., 2006, AJ, 131, 2525.
- Hessels J. W. T., Roberts M. S. E., Ransom S. M., Kaspi V. M., Romani R. W., Ng C.-Y., Freire P. C. C. & Gaensler B. M., 2004, ApJ, 612, 389.
- Hinton J. A., Funk S., Carrigan S., Gallant Y. A., de Jager O. C., Kosack K., Lemièrre A. & Pühlhofer G., 2007, A&A, 476, L25.
- Hui C. Y. & Becker W., 2007, A&A, 470, 965.
- Hutchings J. B., Crampton D. & Cowley P. A., 1981, AJ, 86, 871.
- Hurley-Walker N., *et al.*, 2009, MNRAS, in press.
- Ho P. T., Jackson J. M., Barrett A. H. & Armstrong J. T., 1985, ApJ, 288, 575.
- Jones B. B., 1973, AuJPh, 26, 545.
- Jones T. J., Garwood R. & Dickey J. M., 1988, ApJ, 328, 559.
- Kang J.-H. & Koo B.-C., 2007, ApJS, 173, 85.
- Kaplan D. L., Kulkarni S. R., Frail D. A. & van Kerkwijk M. H., 2002, ApJ, 566, 378.
- Kaspi V. M., Roberts M. S. E. & Harding A. K., 2006, in *Compact stellar X-ray sources*, (Cambridge Astrophysics Series, Volume 39), eds by Lewin W. & van der Klis M., (Cambridge University Press), p279.
- Kassim N. E., 1988, ApJ, 328, L55.
- Kassim N. E., 1989, ApJS, 71, 799.
- Kerton C. R., Murphy J. & Patterson J., 2007, MNRAS, 379, 289.
- Kim K.-T. & Koo B.-C., 2000, ApJ, 529, 229.
- Koo B.-C. & Heiles C., 1991, ApJ, 382, 204.
- Koo B.-C., Kang J. H. & Salter C. J., 2006, ApJ, 643, L49.
- Kothes R., 2003, A&A, 408, 187.
- Kothes R., Fedotov K., Foster T. J. & Uyaniker B., 2006, A&A, 457, 1081.
- Kovalenko A. V., 1989, SvAL, 15, 144.
- Kovalenko A. V., Pynzar' A. V. & Udal'tsov V. A., 1994, ARep, 38, 95.
- Koyama K. *et al.*, 2007, PASJ, 59, S221.
- Landecker T. L., Higgs L. A. & Wendker H. I., 1993, A&A, 276, 522.
- Langston G., Minter A., D'Addario L., Eberhart K., Koski K. & Zuber J., 2000, AJ, 119, 2801.
- LaRosa T. N., Kassim N. E., Lazio T. J. W. & Hyman S. D., 2000, AJ, 119, 207.
- Lazio T. J. W., Anantharamaiah K. R., Goss W. M., Kassim N. E. & Cordes J. M., 1999, ApJ, 515, 196.
- Leahy D. A., Nousek J. & Garmire G., 1992, ApJ, 385, 561.
- Leahy D. A., Tian W. & Wang Q. D., 2008, AJ, 136, 1477.
- Lu F. J., Wang Q. D. & Lang C. C., 2003, AJ, 126, 319.

-
- Maciejewski W., Murphy E. M., Lockman F. J. & Savage B. D., 1996, ApJ, 469, 238.
McClure-Griffiths N. M., Green A. J., Dickey J. M., Gaensler B. M., Haynes R. F. & Wieringa M. H., 2001, ApJ, 551, 394.
McCullough P. R., Fields B. D. & Pavlidou V., 2002, ApJ, 576, L41.
Malizia A. *et al.*, 2005, ApJ, 630, L157.
Manchester R. N. *et al.*, 2002, in *Neutron Stars in Supernova Remnants*, (ASP Conference Series, Volume 271), eds Slane P. O. & Gaensler B. M., (ASP, San Francisco), p.31.
Marston A. P., 1996, AJ, 112, 2828.
Martí J., Combi J. A., Pérez-Ramírez D., Garrido J. L., Luque-Escamilla P., Muñoz-Arjonilla A. J. & Sánchez-Sutil, J. R., 2007, A&A, 462, 1065.
Mavromatakis F. & Strom R. G., 2002, A&A, 382, 291.
Mavromatakis F., Boumis P. & Paleologou E. V., 2002, A&A, 387, 635.
Mavromatakis F., Boumis P., Xilouris E., Papamastorakis J. & Alikakos J., 2005, A&A, 435, 141.
Milne D. K., 1970, AuJPh, 23, 425.
Misanovic Z., Cram L. & Green A., 2002, MNRAS, 335, 114.
Mori H., Tsuru T. G., Hyodo Y., Koyama K. & Senda, A., 2008, PASJ, 60, S183.
Morris P. W., Stolovy S., Wachter S., Noriega-Crespo A., Pannuti T. G. & Hoard D. W., 2006, ApJ, 640, L179.
Muno M. P., Baganoff F. K., Brandt W. N., Morris M. R. & Starck J.-L., 2008, ApJ, 673, 251.
Naranan S., Shulman S., Friedman H. & Fritz G., 1976, ApJ, 208, 718.
Nichols-Bohlin J. & Fesen R. A., 1993, AJ, 105, 672.
Nobukawa M. *et al.*, 2008, PASJ, 60, S191.
Normandeau M., Taylor A. R., Dewdney P. E. & Basu S., 2000, AJ, 119, 2982.
Nousek J. A., Cowie L. L., Hu E., Lindblad C. J. & Garmire G. P., 1981, ApJ, 248, 152.
Odegard N., 1986, AJ, 92, 1372.
Olano C. A., Meschin P. I. & Niemela, V. S., 2006, MNRAS, 369, 867.
Onello J. S., DePree C. G., Phillips J. A. & Goss W. M., 1995, ApJ, 449, L127.
Pauls T. & Schwartz P. R., 1989, in *The Physics and Chemistry of Interstellar Molecular Clouds*, (Lecture Notes in Physics, Volume 331), eds Winnewisser G. & Armstrong T. J. (Springer), p.225.
Phillips J. P. & Ramos-Larios G., 2008, MNRAS, 390, 1170.
Phillips J. A., Onello J. S. & Kulkarni S. R., 1993, ApJ, 415, L143.
Pineault S. & Chastenay P., 1990, MNRAS, 246, 169.
Plucinsky P. P., Snowden S. L., Aschenbach B., Eggar R., Edgar R. J. & McCammon D., 1996, ApJ, 463, 224.
Polcaro V. F., Rossi C., Norci L. & Viotti R., 1995, A&A, 303, 211.
Predehl P. & Kulkarni S. R., 1995, A&A, 294, L29.
Punsly B., Romero G. E., Torres D. F. & Combi J. A., 2000, A&A, 364, 552.
Reach W. T. *et al.*, 2006, AJ, 131, 1479.
Reich W., 2002, in *Neutron Stars, Pulsars, and Supernova Remnants*, (MPE Report 278), eds Becker W., Lesch H. & Trümper J., (Max-Planck-Institut für extraterrestrische Physik, Garching bei München), p1.
Reich W., Zhang X. & Fürst E., 2003, A&A, 408, 961.
Reich W., Fürst E., Altenhoff W. J., Reich P. & Junkes N., 1985, A&A, 151, L10.
Reich W., Fürst E., Reich P. & Reif K., 1990, A&AS, 85, 633.
Reipurth B., Bally J. & Divine D., 1997, AJ, 114, 2708.
Retallack D. S., 1983, MNRAS, 204, 669.
Renaud M., Paron S., Terrier R., Lebrun F., Dubner G., Giacani E. & Bykov A. M., 2006, ApJ, 638, 220.
Reynolds R., 1976, ApJ, 206, 679.
Reynoso E. M. & Dubner G. M., 1997, A&AS, 123, 31.
Reynoso E. M. & Green, A. J., 2007, MNRAS, 375, 92.
Rho J. & Petre R., 1998, ApJ, 503, L167.
Riegler G. R., Agrawal P. C. & Gull S. F., 1980, ApJ, 235, L71.
Roberts M. S. E. & Brogan C. L., 2008, ApJ, 681, 320.
Roberts M. S. E., Romani R. W., Johnston S. & Green A. J., 1999, ApJ, 515, 712.
Roberts M. S. E., Romani R. W. & Johnston S., 2001, ApJ, 561, L187.
Routledge D. & Vaneldik J. F., 1988, ApJ, 326, 751.
Roy S. & Pramesh Rao A., 2002, MNRAS, 329, 775.
Roy S. & Bhatnagar S., 2006, JPhCS, 54, 152.
Russeil D., Adami C., Amram P., Le Coarer E., Georgelin Y. M., Marcellin M. & Parker Q., 2005, A&A, 429, 497.
Russell D. M., Fender R. P., Gallo E. & Kaiser C. R., 2007, MNRAS, 376, 1341.
Ryu K. *et al.*, 2006, ApJ, 644, L185.
Saken J. M., Fesen R. A. & Shull J. M., 1992, ApJS, 81, 715.
Sarma A. P., Goss W. M., Green A. J. & Frail D. A., 1997, ApJ, 483, 335.
Salter C. J., Patnaik A. R., Shaver P. A. & Hunt G. C., 1986, A&A, 162, 217.
Schaudel D., Beck W., Voges W., Aschenbach B., Reich W. & Weisskopf M., 2002, in *Neutron Stars in Supernova Remnants*, (ASP Conference Series, Volume 271), eds Slane P. O. & Gaensler B. M., (ASP, San Francisco), p.391.
Senda A., Murakami H. & Koyama K., 2002, ApJ, 565, 1017.
Senda A., Murakami H. & Koyama K., 2003, AN, 324 (Supplement 1), 151.

-
- Shinn J.-H. *et al.*, 2007, *ApJ*, 670, 1132.
- Snowden S. L., Burrows D. N., Sanders W. T., Aschenbach B. & Pfeffermann E., 1995, *ApJ*, 439, 399.
- Sramek R. A., Cowan J. J., Roberts D. A., Goss W. M. & Ekers R. D., 1992, *AJ*, 104, 704.
- Stephenson F. R. & Green D. A., 2002, *Historical Supernovae and their Remnants*, (Oxford University Press).
- Stil J. M. & Irwin J. A., 2001, *ApJ*, 563, 816.
- Stil J. M., Taylor A. R., Dickey J. M., Kavars D. W., Martin P. G., Rothwell T. A., Boothroyd A. I., Lockman F. J. & McClure-Griffiths N. M., 2006, *AJ*, 132, 1158.
- Stupar M., Parker Q. A. & Filipović M. D., 2007, *MNRAS*, 374, 1441.
- Stupar M., Parker Q. A. & Filipović M. D., 2008, *MNRAS*, 390, 1037.
- Subrahmanyan R., Ekers R. D., Wilson W. E., Goss W. M. & Allen D. A., 1993, *MNRAS*, 263, 868.
- Tateyama C. E., Strauss F. M. & Kaufmann P., 1991, *MNRAS*, 249, 716.
- Taylor A. R., Wallace B. J. & Goss W. M., 1992, *AJ*, 103, 931.
- Thompson D. J., Djorgovski S. & de Carvalho R. R., 1991, *PASP*, 103, 487.
- Thorsett S. E., Benjamin R. A., Briske W. F., Golden A. & Goss W. M., 2003, *ApJ*, 592, L71.
- Tian W. W. & Leahy D. A., 2006, *A&A*, 455, 1053.
- Tian W. W., Leahy D. A. & Foster T. J., 2007, *A&A*, 465, 907.
- Tian W. W., Leahy D. A., Haverkorn M. & Jiang, B., 2008, *ApJ*, 679, L85.
- Trushkin S. A., 1998, *BSAO*, 46, 62.
- Trushkin S. A., 2001, in *Exploring the gamma-ray universe*, eds Batrick N., Gimenez A., Reglero V. & Winkler C., (ESA, Noordwijk), p109.
- Ueno M., Yamaguchi H., Koyama K., Bamba A., Yamauchi S. & Ebisawa K., 2005, in *X-Ray and Radio Connections*, eds Sjouwerman L. O. & Dyer K. K., (4.18). Available from <http://www.aoc.nrao.edu/events/xraydio/>.
- Uyaniker B. & Kothes R., 2002, *ApJ*, 574, 805.
- Velusamy T., Goss W. M. & Arnal E. M., 1986, *JApA*, 7, 105.
- Walker A. & Zealey W. J., 1998, *PASA*, 15, 79.
- Walker A., Zealey W. J. & Parker Q. A., 2001, *PASA*, 18, 259.
- Wang Q. D., Lu F. J. & Gotthelf E. V., 2006, *MNRAS*, 367, 937.
- Ward-Thompson D. & Robson E. I., 1991, *MNRAS*, 248, 670.
- Weiler K. W. & Shaver P. A., 1978, *A&A*, 65, 305.
- Weinberger R., 1995, *PASP*, 107, 58.
- Weinberger R., Tempurin S. & Stecklum B., 2006, *A&A*, 448, 1095.
- Weinberger R., Tajitsu A., Tamura S. & Yadoumaru Y., 1998, *PASP*, 110, 722.
- Wendker H. I., Higgs L. A. & Landecker T. L., 1991, *A&A*, 241, 551.
- Wendker H. I., Higgs L. A., Landecker T. L. & Ward-Thompson D., 1993, *MNRAS*, 263, 543.
- White R. L. & Becker R. H., 1990, *MNRAS*, 244, 12P.
- White R. L., Becker R. H. & Helfand D. J., 2005, *AJ*, 130, 586.
- Whiteoak J. B. Z. & Green A. J., 1996, *A&AS*, 118, 329.
- Winkler P. F. & Reipurth B., 1992, *ApJ*, 389, L25.
- Winkler P. F., Kirshner R. P., Hughes J. P. & Heathcote S. R., 1989, *Nature*, 337, 48.
- Woermann B., Gaylard M. J. & Otrupcek R., 2001, *MNRAS*, 325, 1213.
- Yamaguchi H., Ueno M., Koyama K., Bamba A. & Yamauchi S., 2004, *PASJ*, 56, 1059.
- Yoshita K., Miyata E. & Tsunemi H., 1999, *AN*, 320, 344.
- Yoshita K., Miyata E. & Tsunemi H., 2000, *PASJ*, 52, 867.
- Yusef-Zadeh F., Cotton W. D. & Reynolds S. P., 1998, *ApJ*, 498, L55.
- Yusef-Zadeh F., Shure M., Wardle M. & Kassim N., 2000, *ApJ*, 540, 842.
- Zanin C. & Kerber F., 2000, *A&A*, 356, 274.
- Zhang X. Z., 2003, *AcASn*, 44 (Supplement), 183.
- Zijlstra A. A., 1991, *MNRAS*, 248, 11P.

l	b	RA (J2000.0) (h m s)	Dec ($^{\circ}$ $'$)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
0.0	+0.0	17 45 44	-29 00	3.5×2.5	S	100?	0.8?	Sgr A East
0.3	+0.0	17 46 15	-28 38	15×8	S	22	0.6	
0.9	+0.1	17 47 21	-28 09	8	C	18?	varies	
1.0	-0.1	17 48 30	-28 09	8	S	15	0.6?	
1.4	-0.1	17 49 39	-27 46	10	S	2?	?	
1.9	+0.3	17 48 45	-27 10	1.5	S	0.6	0.6	
3.7	-0.2	17 55 26	-25 50	14×11	S	2.3	0.65	
3.8	+0.3	17 52 55	-25 28	18	S?	3?	0.6	
4.2	-3.5	18 08 55	-27 03	28	S	3.2?	0.6?	
4.5	+6.8	17 30 42	-21 29	3	S	19	0.64	Kepler, SN1604, 3C358
4.8	+6.2	17 33 25	-21 34	18	S	3	0.6	
5.2	-2.6	18 07 30	-25 45	18	S	2.6?	0.6?	
5.4	-1.2	18 02 10	-24 54	35	C?	35?	0.2?	Milne 56
5.5	+0.3	17 57 04	-24 00	15×12	S	5.5	0.7	
5.9	+3.1	17 47 20	-22 16	20	S	3.3?	0.4?	
6.1	+0.5	17 57 29	-23 25	18×12	S	4.5	0.9	
6.1	+1.2	17 54 55	-23 05	30×26	F	4.0?	0.3?	
6.4	-0.1	18 00 30	-23 26	48	C	310	varies	W28
6.4	+4.0	17 45 10	-21 22	31	S	1.3?	0.4?	
6.5	-0.4	18 02 11	-23 34	18	S	27	0.6	
7.0	-0.1	18 01 50	-22 54	15	S	2.5?	0.5?	
7.2	+0.2	18 01 07	-22 38	12	S	2.8	0.6	
7.7	-3.7	18 17 25	-24 04	22	S	11	0.32	1814-24
8.3	-0.0	18 04 34	-21 49	5×4	S	1.2	0.6	
8.7	-5.0	18 24 10	-23 48	26	S	4.4	0.3	
8.7	-0.1	18 05 30	-21 26	45	S?	80	0.5	(W30)
8.9	+0.4	18 03 58	-21 03	24	S	9	0.6	
9.7	-0.0	18 07 22	-20 35	15×11	S	3.7	0.6	
9.8	+0.6	18 05 08	-20 14	12	S	3.9	0.5	
9.9	-0.8	18 10 41	-20 43	12	S	6.7	0.4	
10.5	-0.0	18 09 08	-19 47	6	S	0.9	0.6	
11.0	-0.0	18 10 04	-19 25	11×9	S	1.3	0.6	
11.1	-1.0	18 14 03	-19 46	18×12	S	5.8	0.6	
11.1	-0.7	18 12 46	-19 38	11×7	S	1.0	0.7	
11.1	+0.1	18 09 47	-19 12	12×10	S	2.3	0.4	
11.2	-0.3	18 11 27	-19 25	4	C	22	0.6	
11.4	-0.1	18 10 47	-19 05	8	S?	6	0.5	
11.8	-0.2	18 12 25	-18 44	4	S	0.7	0.3	
12.0	-0.1	18 12 11	-18 37	7?	?	3.5	0.7	
12.2	+0.3	18 11 17	-18 10	6×5	S	0.8	0.7	

l	b	RA (J2000.0) (h m s)	Dec ($^{\circ}$ $'$)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
12.5	+0.2	18 12 14	-17 55	6 \times 5	C?	0.6	0.4	
12.7	-0.0	18 13 19	-17 54	6	S	0.8	0.8	
12.8	-0.0	18 13 37	-17 49	3	C?	0.8	0.5	
13.3	-1.3	18 19 20	-18 00	70 \times 40	S?	?	?	
13.5	+0.2	18 14 14	-17 12	5 \times 4	S	3.5?	1.0?	
14.1	-0.1	18 15 52	-16 34	6 \times 5	S	0.5	0.6	
14.3	+0.1	18 15 58	-16 27	5 \times 4	S	0.6	0.4	
15.1	-1.6	18 24 00	-16 34	30 \times 24	S	5.5?	0.8?	
15.4	+0.1	18 18 02	-15 27	15 \times 14	S	5.6	0.6	
15.9	+0.2	18 18 52	-15 02	7 \times 5	S?	5	0.6?	
16.0	-0.5	18 21 56	-15 14	15 \times 10	S	2.7	0.6	
16.2	-2.7	18 29 40	-16 08	17	S	2	0.5	
16.4	-0.5	18 22 38	-14 55	13	S	4.6	0.7	
16.7	+0.1	18 20 56	-14 20	4	C	3.0	0.6	
16.8	-1.1	18 25 20	-14 46	30 \times 24?	?	2?	?	
17.0	-0.0	18 21 57	-14 08	5	S	0.5	0.5	
17.4	-2.3	18 30 55	-14 52	24?	S	4.8?	0.8?	
17.4	-0.1	18 23 08	-13 46	6	S	0.4	0.7	
17.8	-2.6	18 32 50	-14 39	24	S	4.0?	0.3?	
18.1	-0.1	18 24 34	-13 11	8	S	4.6	0.5	
18.6	-0.2	18 25 55	-12 50	6	S	1.4	0.4	
18.8	+0.3	18 23 58	-12 23	17 \times 11	S	33	0.4	Kes 67
18.9	-1.1	18 29 50	-12 58	33	C?	37	varies	
19.1	+0.2	18 24 56	-12 07	27	S	10	0.5	
20.0	-0.2	18 28 07	-11 35	10	F	10	0.0	
20.4	+0.1	18 27 51	-11 00	8	S	3.1	0.4	
21.0	-0.4	18 31 12	-10 47	9 \times 7	S	1.1	0.6	
21.5	-0.9	18 33 33	-10 35	4	C	6?	0.0	
21.5	-0.1	18 30 50	-10 09	5	S	0.4	0.5	
21.8	-0.6	18 32 45	-10 08	20	S	69	0.5	Kes 69
22.7	-0.2	18 33 15	-09 13	26	S?	33	0.6	
23.3	-0.3	18 34 45	-08 48	27	S	70	0.5	W41
23.6	+0.3	18 33 03	-08 13	10?	?	8?	0.3	
24.7	-0.6	18 38 43	-07 32	15?	S?	8	0.5	
24.7	+0.6	18 34 10	-07 05	30 \times 15	C?	20?	0.2?	
27.4	+0.0	18 41 19	-04 56	4	S	6	0.68	4C-04.71
27.8	+0.6	18 39 50	-04 24	50 \times 30	F	30	varies	
28.6	-0.1	18 43 55	-03 53	13 \times 9	S	3?	?	
28.8	+1.5	18 39 00	-02 55	100?	S?	?	0.4?	
29.6	+0.1	18 44 52	-02 57	5	S	1.5?	0.5?	

<i>l</i>	<i>b</i>	RA (J2000.0) (h m s)	Dec (° ′)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
29.7	-0.3	18 46 25	-02 59	3	C	10	0.7	Kes 75
30.7	-2.0	18 54 25	-02 54	16	?	0.5?	0.7?	
30.7	+1.0	18 44 00	-01 32	24 × 18	S?	6	0.4	
31.5	-0.6	18 51 10	-01 31	18?	S?	2?	?	
31.9	+0.0	18 49 25	-00 55	7 × 5	S	24	0.49	3C391
32.0	-4.9	19 06 00	-03 00	60?	S?	22?	0.5?	3C396.1
32.1	-0.9	18 53 10	-01 08	40?	C?	?	?	
32.4	+0.1	18 50 05	-00 25	6	S	0.25?	?	
32.8	-0.1	18 51 25	-00 08	17	S?	11?	0.2?	Kes 78
33.2	-0.6	18 53 50	-00 02	18	S	3.5	varies	
33.6	+0.1	18 52 48	+00 41	10	S	22	0.5	Kes 79, 4C00.70, HC13
34.7	-0.4	18 56 00	+01 22	35 × 27	C	230	0.37	W44, 3C392
36.6	-0.7	19 00 35	+02 56	25?	S?	?	?	
36.6	+2.6	18 48 49	+04 26	17 × 13?	S	0.7?	0.5?	
39.2	-0.3	19 04 08	+05 28	8 × 6	C	18	0.6	3C396, HC24, NRAO 593
39.7	-2.0	19 12 20	+04 55	120 × 60	?	85?	0.7?	W50, SS433
40.5	-0.5	19 07 10	+06 31	22	S	11	0.5	
41.1	-0.3	19 07 34	+07 08	4.5 × 2.5	S	22	0.48	3C397
42.8	+0.6	19 07 20	+09 05	24	S	3?	0.5?	
43.3	-0.2	19 11 08	+09 06	4 × 3	S	38	0.48	W49B
43.9	+1.6	19 05 50	+10 30	60?	S?	8.6?	0.2?	
45.7	-0.4	19 16 25	+11 09	22	S	4.2?	0.4?	
46.8	-0.3	19 18 10	+12 09	17 × 13	S	14	0.5	(HC30)
49.2	-0.7	19 23 50	+14 06	30	S?	160?	0.3?	(W51)
53.6	-2.2	19 38 50	+17 14	33 × 28	S	8	0.75	3C400.2, NRAO 611
54.1	+0.3	19 30 31	+18 52	1.5	F?	0.5	0.1	
54.4	-0.3	19 33 20	+18 56	40	S	28	0.5	(HC40)
55.0	+0.3	19 32 00	+19 50	20 × 15?	S	0.5?	0.5?	
55.7	+3.4	19 21 20	+21 44	23	S	1.4	0.6	
57.2	+0.8	19 34 59	+21 57	12?	S?	1.8?	?	(4C21.53)
59.5	+0.1	19 42 33	+23 35	15	S	3?	?	
59.8	+1.2	19 38 55	+24 19	20 × 16?	?	1.6	0.5	
63.7	+1.1	19 47 52	+27 45	8	F	1.8	0.3	
65.1	+0.6	19 54 40	+28 35	90 × 50	S	5.5	0.61	
65.3	+5.7	19 33 00	+31 10	310 × 240	S?	52?	0.6?	
65.7	+1.2	19 52 10	+29 26	22	F	5.1	varies	DA 495
67.7	+1.8	19 54 32	+31 29	15 × 12	S	1.0	0.5	
68.6	-1.2	20 08 40	+30 37	23	?	0.7?	0.0?	
69.0	+2.7	19 53 20	+32 55	80?	?	120?	varies	CTB 80
69.7	+1.0	20 02 40	+32 43	16 × 14	S	2.0	0.7	

<i>l</i>	<i>b</i>	RA (J2000.0) (h m s)	Dec (° ′)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
73.9	+0.9	20 14 15	+36 12	27	S?	9	0.23	
74.0	-8.5	20 51 00	+30 40	230 × 160	S	210	varies	Cygnus Loop
74.9	+1.2	20 16 02	+37 12	8 × 6	F	9	varies	CTB 87
76.9	+1.0	20 22 20	+38 43	9	?	1.2	0.60	
78.2	+2.1	20 20 50	+40 26	60	S	320	0.51	DR4, γ Cygni SNR
82.2	+5.3	20 19 00	+45 30	95 × 65	S	120?	0.5?	W63
83.0	-0.3	20 46 55	+42 52	9 × 7	S	1	0.4	
84.2	-0.8	20 53 20	+43 27	20 × 16	S	11	0.5	
85.4	+0.7	20 50 40	+45 22	24?	S	?	0.2	
85.9	-0.6	20 58 40	+44 53	24	S	?	0.2	
89.0	+4.7	20 45 00	+50 35	120 × 90	S	220	0.38	HB21
93.3	+6.9	20 52 25	+55 21	27 × 20	C?	9	0.45	DA 530, 4C(T)55.38.1
93.7	-0.2	21 29 20	+50 50	80	S	65	0.65	CTB 104A, DA 551
94.0	+1.0	21 24 50	+51 53	30 × 25	S	13	0.48	3C434.1
96.0	+2.0	21 30 30	+53 59	26	S	0.3	0.5	
106.3	+2.7	22 27 30	+60 50	60 × 24	C?	6	0.6	
108.2	-0.6	22 53 40	+58 50	70 × 54	S	8	0.5	
109.1	-1.0	23 01 35	+58 53	28	S	22	0.50	CTB 109
111.7	-2.1	23 23 26	+58 48	5	S	2720	0.77	Cassiopeia A, 3C461
113.0	+0.2	23 36 35	+61 22	40 × 17?	?	?	?	
114.3	+0.3	23 37 00	+61 55	90 × 55	S	5.5	0.5	
116.5	+1.1	23 53 40	+63 15	80 × 60	S	10	0.5	
116.9	+0.2	23 59 10	+62 26	34	S	8	0.61	CTB 1
119.5	+10.2	00 06 40	+72 45	90?	S	36	0.6	CTA 1
120.1	+1.4	00 25 18	+64 09	8	S	56	0.65	Tycho, 3C10, SN1572
126.2	+1.6	01 22 00	+64 15	70	S?	6	0.5	
127.1	+0.5	01 28 20	+63 10	45	S	12	0.45	R5
130.7	+3.1	02 05 41	+64 49	9 × 5	F	33	0.07	3C58, SN1181
132.7	+1.3	02 17 40	+62 45	80	S	45	0.6	HB3
156.2	+5.7	04 58 40	+51 50	110	S	5	0.5	
160.9	+2.6	05 01 00	+46 40	140 × 120	S	110	0.64	HB9
166.0	+4.3	05 26 30	+42 56	55 × 35	S	7	0.37	VRO 42.05.01
179.0	+2.6	05 53 40	+31 05	70	S?	7	0.4	
180.0	-1.7	05 39 00	+27 50	180	S	65	varies	S147
182.4	+4.3	06 08 10	+29 00	50	S	1.2	0.4	
184.6	-5.8	05 34 31	+22 01	7 × 5	F	1040	0.30	Crab Nebula, 3C144, SN1054
189.1	+3.0	06 17 00	+22 34	45	C	160	0.36	IC443, 3C157
192.8	-1.1	06 09 20	+17 20	78	S	20?	0.6?	PKS 0607+17
205.5	+0.5	06 39 00	+06 30	220	S	160	0.5	Monoceros Nebula
206.9	+2.3	06 48 40	+06 26	60 × 40	S?	6	0.5	PKS 0646+06

l	b	RA (J2000.0) (h m s)	Dec ($^{\circ}$ $'$)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
260.4	-3.4	08 22 10	-43 00	60 \times 50	S	130	0.5	Puppis A, MSH 08-44
261.9	+5.5	09 04 20	-38 42	40 \times 30	S	10?	0.4?	
263.9	-3.3	08 34 00	-45 50	255	C	1750	varies	Vela (XYZ)
266.2	-1.2	08 52 00	-46 20	120	S	50?	0.3?	RX J0852.0-4622
272.2	-3.2	09 06 50	-52 07	15?	S?	0.4	0.6	
279.0	+1.1	09 57 40	-53 15	95	S	30?	0.6?	
284.3	-1.8	10 18 15	-59 00	24?	S	11?	0.3?	MSH 10-53
286.5	-1.2	10 35 40	-59 42	26 \times 6	S?	1.4?	?	
289.7	-0.3	11 01 15	-60 18	18 \times 14	S	6.2	0.2?	
290.1	-0.8	11 03 05	-60 56	19 \times 14	S	42	0.4	MSH 11-61A
291.0	-0.1	11 11 54	-60 38	15 \times 13	C	16	0.29	(MSH 11-62)
292.0	+1.8	11 24 36	-59 16	12 \times 8	C	15	0.4	MSH 11-54
292.2	-0.5	11 19 20	-61 28	20 \times 15	S	7	0.5	
293.8	+0.6	11 35 00	-60 54	20	C	5?	0.6?	
294.1	-0.0	11 36 10	-61 38	40	S	>2?	?	
296.1	-0.5	11 51 10	-62 34	37 \times 25	S	8?	0.6?	
296.5	+10.0	12 09 40	-52 25	90 \times 65	S	48	0.5	PKS 1209-51/52
296.8	-0.3	11 58 30	-62 35	20 \times 14	S	9	0.6	1156-62
298.5	-0.3	12 12 40	-62 52	5?	?	5?	0.4?	
298.6	-0.0	12 13 41	-62 37	12 \times 9	S	5?	0.3	
299.2	-2.9	12 15 13	-65 30	18 \times 11	S	0.5?	?	
299.6	-0.5	12 21 45	-63 09	13	S	1.0?	?	
301.4	-1.0	12 37 55	-63 49	37 \times 23	S	2.1?	?	
302.3	+0.7	12 45 55	-62 08	17	S	5?	0.4?	
304.6	+0.1	13 05 59	-62 42	8	S	14	0.5	Kes 17
308.1	-0.7	13 37 37	-63 04	13	S	1.2?	?	
308.8	-0.1	13 42 30	-62 23	30 \times 20?	C?	15?	0.4?	
309.2	-0.6	13 46 31	-62 54	15 \times 12	S	7?	0.4?	
309.8	+0.0	13 50 30	-62 05	25 \times 19	S	17	0.5	
310.6	-0.3	13 58 00	-62 09	8	S	5?	?	Kes 20B
310.8	-0.4	14 00 00	-62 17	12	S	6?	?	Kes 20A
311.5	-0.3	14 05 38	-61 58	5	S	3?	0.5	
312.4	-0.4	14 13 00	-61 44	38	S	45	0.36	
312.5	-3.0	14 21 00	-64 12	20 \times 18	S	3.5?	?	
315.1	+2.7	14 24 30	-57 50	190 \times 150	S	?	?	
315.4	-2.3	14 43 00	-62 30	42	S	49	0.6	RCW 86, MSH 14-63
315.4	-0.3	14 35 55	-60 36	24 \times 13	?	8	0.4	
315.9	-0.0	14 38 25	-60 11	25 \times 14	S	0.8?	?	
316.3	-0.0	14 41 30	-60 00	29 \times 14	S	20?	0.4	(MSH 14-57)
317.3	-0.2	14 49 40	-59 46	11	S	4.7?	?	

l	b	RA (J2000.0) (h m s)	Dec ($^{\circ}$ $'$)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
318.2	+0.1	14 54 50	-59 04	40 × 35	S	>3.9?	?	
318.9	+0.4	14 58 30	-58 29	30 × 14	C	4?	0.2?	
320.4	-1.2	15 14 30	-59 08	35	C	60?	0.4	MSH 15-52, RCW 89
320.6	-1.6	15 17 50	-59 16	60 × 30	S	?	?	
321.9	-1.1	15 23 45	-58 13	28	S	>3.4?	?	
321.9	-0.3	15 20 40	-57 34	31 × 23	S	13	0.3	
322.5	-0.1	15 23 23	-57 06	15	C	1.5	0.4	
323.5	+0.1	15 28 42	-56 21	13	S	3?	0.4?	
326.3	-1.8	15 53 00	-56 10	38	C	145	varies	MSH 15-56
327.1	-1.1	15 54 25	-55 09	18	C	7?	?	
327.2	-0.1	15 50 55	-54 18	5	S	0.4	?	
327.4	+0.4	15 48 20	-53 49	21	S	30?	0.6	Kes 27
327.4	+1.0	15 46 48	-53 20	14	S	1.9?	?	
327.6	+14.6	15 02 50	-41 56	30	S	19	0.6	SN1006, PKS 1459-41
328.4	+0.2	15 55 30	-53 17	5	F	15	0.0	(MSH 15-57)
329.7	+0.4	16 01 20	-52 18	40 × 33	S	>34?	?	
330.0	+15.0	15 10 00	-40 00	180?	S	350?	0.5?	Lupus Loop
330.2	+1.0	16 01 06	-51 34	11	S?	5?	0.3	
332.0	+0.2	16 13 17	-50 53	12	S	8?	0.5	
332.4	-0.4	16 17 33	-51 02	10	S	28	0.5	RCW 103
332.4	+0.1	16 15 20	-50 42	15	S	26	0.5	MSH 16-51, Kes 32
332.5	-5.6	16 43 20	-54 30	35	S	2?	0.7?	
335.2	+0.1	16 27 45	-48 47	21	S	16	0.5	
336.7	+0.5	16 32 11	-47 19	14 × 10	S	6	0.5	
337.0	-0.1	16 35 57	-47 36	1.5	S	1.5	0.6?	(CTB 33)
337.2	-0.7	16 39 28	-47 51	6	S	1.5	0.4	
337.2	+0.1	16 35 55	-47 20	3 × 2	?	1.5?	?	
337.3	+1.0	16 32 39	-46 36	15 × 12	S	16	0.55	Kes 40
337.8	-0.1	16 39 01	-46 59	9 × 6	S	18	0.5	Kes 41
338.1	+0.4	16 37 59	-46 24	15?	S	4?	0.4	
338.3	-0.0	16 41 00	-46 34	8	C?	7?	?	
338.5	+0.1	16 41 09	-46 19	9	?	12?	?	
340.4	+0.4	16 46 31	-44 39	10 × 7	S	5	0.4	
340.6	+0.3	16 47 41	-44 34	6	S	5?	0.4?	
341.2	+0.9	16 47 35	-43 47	22 × 16	C	1.5?	0.6?	
341.9	-0.3	16 55 01	-44 01	7	S	2.5	0.5	
342.0	-0.2	16 54 50	-43 53	12 × 9	S	3.5?	0.4?	
342.1	+0.9	16 50 43	-43 04	10 × 9	S	0.5?	?	
343.0	-6.0	17 25 00	-46 30	250	S	?	?	RCW 114
343.1	-2.3	17 08 00	-44 16	32?	C?	8?	0.5?	

l	b	RA (J2000.0) (h m s)	Dec (° ′)	size /arcmin	type	Flux at 1 GHz/Jy	spectral index	other name(s)
343.1	-0.7	17 00 25	-43 14	27 × 21	S	7.8	0.55	
344.7	-0.1	17 03 51	-41 42	10	C?	2.5?	0.5	
345.7	-0.2	17 07 20	-40 53	6	S	0.6?	?	
346.6	-0.2	17 10 19	-40 11	8	S	8?	0.5?	
347.3	-0.5	17 13 50	-39 45	65 × 55	S?	?	?	
348.5	-0.0	17 15 26	-38 28	10?	S?	10?	0.4?	
348.5	+0.1	17 14 06	-38 32	15	S	72	0.3	CTB 37A
348.7	+0.3	17 13 55	-38 11	17?	S	26	0.3	CTB 37B
349.2	-0.1	17 17 15	-38 04	9 × 6	S	1.4?	?	
349.7	+0.2	17 17 59	-37 26	2.5 × 2	S	20	0.5	
350.0	-2.0	17 27 50	-38 32	45	S	26	0.4	
350.1	-0.3	17 17 40	-37 24	4?	?	6?	0.8?	
351.2	+0.1	17 22 27	-36 11	7	C?	5?	0.4	
351.7	+0.8	17 21 00	-35 27	18 × 14	S	10	0.5?	
351.9	-0.9	17 28 52	-36 16	12 × 9	S	1.8?	?	
352.7	-0.1	17 27 40	-35 07	8 × 6	S	4	0.6	
353.6	-0.7	17 32 00	-34 44	30	S	2.5?	?	
353.9	-2.0	17 38 55	-35 11	13	S	1?	0.5?	
354.1	+0.1	17 30 28	-33 46	15 × 3?	C?	?	varies	
354.8	-0.8	17 36 00	-33 42	19	S	2.8?	?	
355.4	+0.7	17 31 20	-32 26	25	S	5?	?	
355.6	-0.0	17 35 16	-32 38	8 × 6	S	3?	?	
355.9	-2.5	17 45 53	-33 43	13	S	8	0.5	
356.2	+4.5	17 19 00	-29 40	25	S	4	0.7	
356.3	-0.3	17 37 56	-32 16	11 × 7	S	3?	?	
356.3	-1.5	17 42 35	-32 52	20 × 15	S	3?	?	
357.7	-0.1	17 40 29	-30 58	8 × 3?	?	37	0.4	MSH 17-39
357.7	+0.3	17 38 35	-30 44	24	S	10	0.4?	
358.0	+3.8	17 26 00	-28 36	38	S	1.5?	?	
358.1	+0.1	17 37 00	-29 59	20	S	2?	?	
358.5	-0.9	17 46 10	-30 40	17	S	4?	?	
359.0	-0.9	17 46 50	-30 16	23	S	23	0.5	
359.1	-0.5	17 45 30	-29 57	24	S	14	0.4?	
359.1	+0.9	17 39 36	-29 11	12 × 11	S	2?	?	

Table II

Other names for SNRs

γ Cygni SNR G78.2+2.1	HB3 G132.7+1.3	NRAO 593 G39.2-0.3
	HB9 G160.9+2.6	NRAO 611 G53.6-2.2
1156-62 G296.8-0.3	HB21 G89.0+4.7	
1814-24 G7.7-3.7		PKS 0607+17 G192.8-1.1
	HC13 G33.6+0.1	PKS 0646+06 G206.9+2.3
3C10 G120.1+1.4	HC24 G39.2-0.3	PKS 1209-51/52 G296.5+10.0
3C58 G130.7+3.1	(HC30) G46.8-0.3	PKS 1459-41 G327.6+14.6
3C144 G184.6-5.8	(HC40) G54.4-0.3	
3C157 G189.1+3.0		Puppis A G260.4-3.4
3C358 G4.5+6.8	IC443 G189.1+3.0	
3C391 G31.9+0.0		R5 G127.1+0.5
3C392 G34.7-0.4	Kepler G4.5+6.8	
3C396 G39.2-0.3		RCW 86 G315.4-2.3
3C396.1 G32.0-4.9	Kes 17 G304.6+0.1	RCW 89 G320.4-1.2
3C397 G41.1-0.3	Kes 20A G310.6-0.3	RCW 103 G332.4-0.4
3C400.2 G53.6-2.2	Kes 20B G310.8-0.4	RCW 114 G343.0-6.0
3C434.1 G94.0+1.0	Kes 27 G327.4+0.4	
3C461 G111.7-2.1	Kes 32 G332.4+0.1	RX J0852.0-4622 G266.2-1.2
	Kes 40 G337.3+1.0	
4C-04.71 G27.4+0.0	Kes 41 G337.8-0.1	S147 G180.0-1.7
4C00.70 G33.6+0.1	Kes 67 G18.8+0.3	
(4C21.53) G57.2+0.8	Kes 69 G21.8-0.6	SN1006 G327.6+14.6
4C(T)55.38.1 G93.3+6.9	Kes 75 G29.7-0.3	SN1054 G184.6-5.8
	Kes 78 G32.8-0.1	SN1181 G130.7+3.1
	Kes 79 G33.6+0.1	SN1572 G120.1+1.4
CTA 1 G119.5+10.2		SN1604 G4.5+6.8
	Lupus Loop G330.0+15.0	
CTB 1 G116.9+0.2		SS433 G39.7-2.0
(CTB 33) G337.0-0.1		
CTB 37A G348.5+0.1	MSH 08-44 G260.4-3.4	
CTB 37B G348.7+0.3	MSH 10-53 G284.3-1.8	Sgr A East G0.0+0.0
CTB 80 G69.0+2.7	MSH 11-54 G292.0+1.8	
CTB 87 G74.9+1.2	MSH 11-61A G290.1-0.8	Tycho G120.1+1.4
CTB 104A G93.7-0.2	(MSH 11-62) G291.0-0.1	
CTB 109 G109.1-1.0	(MSH 14-57) G316.3-0.0	Vela (XYZ) G263.9-3.3
	MSH 14-63 G315.4-2.3	
Cassiopeia A G111.7-2.1	MSH 15-52 G320.4-1.2	VRO 42.05.01 G166.0+4.3
	MSH 15-56 G326.3-1.8	
Crab Nebula G184.6-5.8	(MSH 15-57) G328.4+0.2	W28 G6.4-0.1
	MSH 16-51 G332.4+0.1	(W30) G8.7-0.1
Cygnus Loop G74.0-8.5	MSH 17-39 G357.7-0.1	W41 G23.3-0.3
		W44 G34.7-0.4
DA 495 G65.7+1.2	Milne 56 G5.4-1.2	W49B G43.3-0.2
DA 530 G93.3+6.9		W50 G39.7-2.0
DA 551 G93.7-0.2	Monoceros Nebula G205.5+0.5	(W51) G49.2-0.7
		W63 G82.2+5.3
DR4 G78.2+2.1		

Journals

AdSpR	Advances in Space Research
A&A	Astronomy & Astrophysics
A&AS	Astronomy & Astrophysics Supplement
AJ	Astronomical Journal
AN	Astronomische Nachrichten
ApJ	Astrophysical Journal
ApJS	Astrophysical Journal Supplement
ApL	Astrophysical Letters
ApS&S	Astrophysics & Space Science
ARep	Astronomy Reports
AuJPA	Australian Journal of Physics Astrophysical Supplement
AuJPh	Australian Journal of Physics
BASI	Bulletin of the Astronomical Society of India
ChJAA	Chinese Journal of Astronomy & Astrophysics
JApA	Journal of Astrophysics & Astronomy
JPhCS	Journal of Physics Conference Series
MNRAS	Monthly Notices of the Royal Astronomical Society
NuPhS	Nuclear Physics B Proceedings Supplements
PASAU	Proceedings of the Astronomical Society of Australia
PASJ	Publications of the Astronomical Society of Japan
PASP	Publications of the Astronomical Society of the Pacific
RMxAA	Revista Mexicana de Astronomía y Astrofísica
SerAJ	Serbian Astronomical Journal
SvAL	Soviet Astronomy Letters

Proceedings

SNRISM is ‘*Supernova Remnants and the Interstellar Medium*’, (IAU Colloquium 101), eds Roger R. S. & Landecker T. L., (Cambridge University Press), 1988.

NSPS is ‘*Neutron Stars, Pulsars, and Supernova Remnants*’, (MPE Report 278), eds Becker W., Lesch H. & Trümper J., (Max-Planck-Institut für extraterrestrische Physik, Garching bei München), 2002.

XRRC is ‘*X-Ray and Radio Connections*’, eds Sjouwerman L. O. & Dyer K. K., (available at <http://www.aoc.nrao.edu/events/xraydio/>), 2005.

Radio Telescopes/Surveys

ATCA	Australia Telescope Compact Array
BIMA	Berkeley–Illinois–Maryland Array
CLFST	Cambridge Low Frequency Synthesis Telescope
DRAO	Dominion Radio Astrophysical Observatory
FIRST	Flours Synthesis Telescope
GBT	Green Bank Telescope
MOST	Molonglo Observatory Synthesis Telescope
NRAO	National Radio Astronomy Observatory
NRO	Nobeyama Radio Observatory
TPT	Clark Lake Teepee-Tee telescope
VLA	Very Large Array
VRO	Vermillion River Observatory
WSRT	Westerbork Synthesis Radio Telescope

(C/S/V)GPS (Canadian/Southern/VLA) Galactic Plane Survey

Satellites

HST	Hubble Space Telescope
ISO	Infrared Space Observatory
ASCA	Advanced Satellite for Cosmology and Astrophysics
EXOSAT	European X-ray Observatory Satellite
ROSAT	Röntgensatellit
XMM	X-ray Multi-Mirror(-Newton)