

G0.0+0.0

RA: 17^h45^m44^s
Dec: −29°00'

1-GHz flux/Jy: 100?
Spectral index: 0.8?

Sgr A East

Size/arcmin: 3.5×2.5
Type: S

Radio: Non-thermal shell, in complex region, interacting with molecular material to the west.

X-ray: Diffuse emission, centrally peaked.

Point sources: Compact X-ray/radio source.

References:

- Ekers *et al.* 1983, A&A, 122, 143. VLA at 1.4 GHz and 5 GHz (both 5''×8''), $S_{1.4 \text{ GHz}} = 77$ Jy, $S_{5.0 \text{ GHz}} = 31$ Jy.
 Pedlar *et al.* 1989, ApJ, 342, 769. VLA at 332 MHz (12''), 1.4 GHz (1''.3×2''.5) and 5 GHz (1''.3×2''.5).
 Mezger *et al.* 1989, A&A, 209, 337. Nearby molecular material.
 Ho *et al.* 1991, Nature, 350, 309. VLA of NH₃ emission from surroundings.
 Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (17''×33'').
 Serabyn *et al.* 1992, ApJ, 395, 166. Nearby molecular material.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×88'').
 Yusef-Zadeh & Mehringer 1995, ApJ, 452, L37. VLA of nearby H₂O masers.
 Yusef-Zadeh *et al.* 1999, ApJ, 512, 230. OH maser observations.
 Coil & Ho 2000, ApJ, 533, 245. NH₃ observations of surroundings.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24''×43'').
 see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Fatuzzo *et al.* 2001, ApJ, 549, 293. Electron–positron lines from the vicinity.
 Yusef-Zadeh *et al.* 2001, ApJ, 560, 749. Observations of nearby molecular hydrogen.
 Maeda *et al.* 2002, ApJ, 570, 671. Chandra observations.
 Sakano *et al.* 2003, AN, 324 (No S1), 197. XMM-Newton observations.
 Roy & Rao 2004, MNRAS, 349, L25. GMRT at 620 MHz (6''.6×11''.4).
 Sakano *et al.* 2004, MNRAS, 350, 129. XMM-Newton observa-

- tions.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (1''.2×2''.2).
 Park *et al.* 2005, ApJ, 631, 964. Chandra observations.
 Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.
 Koyama *et al.* 2007, PASJ, 59, S237. Suzaku observations.
 Lee *et al.* 2008, ApJ, 674, 247. Molecular H₂ observations of surroundings.
 Sjouwerman & Pihlström 2008, ApJ, 681, 1287. VLA at 1.7 GHz of OH masers.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Tsuboi *et al.* 2012, PASJ, 64, 111. CO observations of SW.
 Minh *et al.* 2013, ApJ, 773, 31. NH₃ observations of region.
 Zhao *et al.* 2013, ApJ, 777, 146. VLA at 4.8 GHz (0''.5×0''.7), 5.5 GHz (0''.6×1''.6), and 8.3 GHz (1''.0×2''.0).
 Nynka *et al.* 2013, ApJ, 778, L31. NuSTAR of compact X-ray source.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA observations of methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Tsuboi *et al.* 2015, PASJ, 67, 109. CO observations of region.
 Lau *et al.* 2015, Science, 348, 413. IR observations.
 Ajello *et al.* 2016, ApJ, 819, 44. γ -ray observations.
 Yusef-Zadeh *et al.* 2016, ApJ, 819, 60. VLA at 1.5 GHz (0''.5×1''.4) and other frequencies.
 McEwen *et al.* 2016, ApJ, 832, 129. CH₃OH maser observations in region.

G0.3+0.0

RA: 17^h46^m15^s
Dec: −28°38'

1-GHz flux/Jy: 22
Spectral index: 0.6

Size/arcmin: 15×8
Type: S

Has been called G0.33+0.04 and G0.4+0.1.

Radio: Bilateral shell, near Galactic Centre.

References:

- Kassim & Frail 1996, MNRAS, 283, L51. VLA at 333 MHz (23''×42''), plus review of flux densities and other observations.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (30'').
 see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.

- Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Ajello *et al.* 2016, ApJ, 819, 44. γ -ray observations.

G0.9+0.1

RA: 17^h47^m21^s
Dec: −28°09'

1-GHz flux/Jy: 18?
Spectral index: varies

Size/arcmin: 8
Type: C

Radio: Flat spectrum core within steep spectrum shell.

X-ray: Central core, with non-thermal spectrum.

Point sources: Central pulsar.

References:

- Helfand & Becker 1987, ApJ, 314, 203. VLA at 1.4 and 5 GHz, and Einstein observations.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×91'').
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Mereghetti *et al.* 1998, A&A, 331, L77. X-ray detection.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43''×24'').
 see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Sidoli *et al.* 2000, A&A, 361, 719. BeppoSAX observations.
 Gaensler *et al.* 2001, ApJ, 556, L107. Chandra observations.
 Porquet *et al.* 2003, A&A, 401, 197. XMM-Newton observations.

- Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.6×11''.3).
 Aharonian *et al.* 2005, A&A, 432, L25. γ-ray detection.
 Dubner *et al.* 2008, A&A, 487, 1033. ATCA and VLA at 1.4 GHz (1''.2×2''.5 : $S = 8.3 \pm 0.7$ Jy), 5 GHz (1''.6×2''.5) and 8.3 GHz (0''.8×1''.5).
 Camilo *et al.* 2009, ApJ, 700, L34. Pulsar detection.
 Holler *et al.* 2012, A&A, 539, A24. Chandra and XMM-Newton observations.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Archer *et al.* 2016, ApJ, 821, 129. γ-ray observations.

G1.0−0.1

RA: 17^h48^m30^s
Dec: −28°09'

1-GHz flux/Jy: 15
Spectral index: 0.6?

Size/arcmin: 8
Type: S

Has been called G1.05−0.1 and G1.05−0.15.

Radio: Incomplete shell, to the S of Sgr D.

X-ray: Possibly detected.

References:

- Downes *et al.* 1979, A&AS, 35, 1. Review of flux densities.
 Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (64''×100'': $S = 12.3$ Jy).
 Liszt 1992, ApJS, 82, 495. VLA at 1.6 GHz (13''×23'').
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×91'').
 Mehringer *et al.* 1998, ApJ, 493, 274. VLA at 1.6 GHz (15''×28'') and 5 GHz, including masers observations.
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43''×24'').
 see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Sidoli *et al.* 2001, A&A, 372, 651. BeppoSAX possible detec-

- tion.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Nobukawa *et al.* 2009, AdSpR, 43, 1045. Suzaku observations.
 Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G1.4−0.1

RA: 17^h49^m39^s
Dec: −27°46'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 10
Type: S

Radio: Shell, brightest in E.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×92'': $S = 2$ Jy).
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (2''.7×2''.4 : $S = 4.2 \pm 0.5$).
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.2×12''.2).
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for

- OH masers.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA observations of methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 McEwen *et al.* 2016, ApJ, 826, 189. NH₃ and CH₃OH observations.

G1.9+0.3

RA: $17^{\text{h}}48^{\text{m}}45^{\text{s}}$
Dec: $-27^{\circ}10'$

1-GHz flux/Jy: 0.6
Spectral index: 0.6

Size/arcmin: 1.5
Type: S

Radio: Shell, brighter to the N, brightening; shows secular increase.

X-ray: Shell, with bright limbs to E and W.

Distance: HI absorption gives < 10 kpc.

References:

- Green & Gull 1984, Nature, 312, 527. VLA at 5 GHz ($4''.4 \times 2''$).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 94''$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz ($8''.3 \times 12''$).
 Nord *et al.* 2004, AJ, 128, 1646. VLA at 330 MHz ($7'' \times 12''$).
 Green 2004, BASI, 32, 335. VLA at 1.5 GHz ($7''.2 \times 9''.4$).
 Reynolds *et al.* 2008, ApJ, 680, L41. Chandra observations.
 Green *et al.* 2008, MNRAS, 387, L54. VLA at 4.86 GHz ($4'' \times 10''$), for expansion studies.
 Murphy *et al.* 2008, MNRAS, 389, L23. MOST at 843 MHz for flux increase.
 Gómez & Rodríguez 2009, RMxAA, 45, 91. VLA at 1.5 GHz
 (5''.1 \times 10''.6).
 Reynolds *et al.* 2009, ApJ, 695, L149. Chandra spectroscopy.
 Borkowski *et al.* 2010, ApJ, 724, L161. Chandra observations.
 Carlton *et al.* 2011, ApJ, 737, L22. Chandra expansion studies.
 Borkowski *et al.* 2013, ApJ, 771, L9. Chandra observations.
 Abramowski *et al.* 2014, MNRAS, 441, 790. γ -ray observations.
 Borkowski *et al.* 2014, ApJ, 790, L18. Chandra expansion studies.
 Roy & Pal 2014, IAUS, 296, 197. GMRT HI observations.
 De Horta *et al.* 2014, SerAJ, 189, 41. ATCA at 1.4 (5''.4 \times 10''.4), 2.4 (2''.9 \times 6''.1) and 5 GHz (1''.2 \times 2''.8).
 Gök & Ergin 2015, AdSpR, 56, 1793. Suzaku and γ -ray observations.
 Zoglauer *et al.* 2015, ApJ, 798, 98. NuSTAR observations.

G3.7–0.2

RA: $17^{\text{h}}55^{\text{m}}26^{\text{s}}$
Dec: $-25^{\circ}50'$

1-GHz flux/Jy: 2.3
Spectral index: 0.65

Size/arcmin: 14×11
Type: S

Has been called G003.8–00.3.

Radio: Double arc.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43'' \times 99''$: $S=2.4$ Jy).
 Gaensler 1998, ApJ, 493, 781. VLA at 1.4 GHz ($9'' \times 15''$: $S=1.7 \pm 0.1$ Jy).
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz ($8''.4 \times 11''.4$).

G3.8+0.3

RA: $17^{\text{h}}52^{\text{m}}55^{\text{s}}$
Dec: $-25^{\circ}28'$

1-GHz flux/Jy: 3?
Spectral index: 0.6

Size/arcmin: 18
Type: S?

Radio: Incomplete shell.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43'' \times 100''$: $S=3.5$ Jy).
 Bhattacharjee 2002, MNRAS, 332, 1. GMRT at 327 MHz ($27'' \times 17''$: $S=6.0 \pm 0.4$).

G4.2–3.5

RA: $18^{\text{h}}08^{\text{m}}55^{\text{s}}$
Dec: $-27^{\circ}03'$

1-GHz flux/Jy: 3.2?
Spectral index: 0.6?

Size/arcmin: 28
Type: S

Radio: Elongated shell.

Optical: Detected.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4''.3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G4.5+6.8

RA: 17^h30^m42^s
Dec: -21°29'

1-GHz flux/Jy: 19
Spectral index: 0.64

Kepler, SN1604, 3C358

Size/arcmin: 3
Type: S

This is the remnant of Kepler's SN of AD1604.

Radio: Incomplete shell, brighter to the N.

Optical: Faint filaments.

X-ray: Shell, brighter to the N.

Distance: Optical expansion and proper motion indicates about 2.9 kpc, H_I observations suggest 3.4 to 6.4 kpc.

References:

- van den Bergh & Kamper 1977, ApJ, 218, 617. Optical proper motions.
Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
White & Long 1983, ApJ, 264, 196. Einstein observations.
Matsui *et al.* 1984, ApJ, 287, 295. VLA at 1.4 (2''.5×3''.2) and 5 GHz (3''.2×4''.8) and Einstein image (5'').
Dickel *et al.* 1988, ApJ, 330, 254. VLA at 1.4 (1''.2×2''.3) and 5 GHz (0''.6×1''.0) at two epochs.
Smith *et al.* 1989, ApJ, 347, 925. EXOSAT observations.
Hatsukade *et al.* 1990, PASJ, 42, 279. X-ray spectrum.
Blair *et al.* 1991, ApJ, 366, 484. Optical imaging and spectroscopy.
Bandiera & van den Bergh 1991, ApJ, 374, 186. Optical changes.
van den Bergh 1991, PASP, 103, 194. Optical imaging.
Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Hughes 1999, ApJ, 527, 298. ROSAT and Einstein image comparison for expansion studies.
Reynoso & Goss 1999, AJ, 118, 926. VLA at 1.4 GHz (23''×13'') for H_I studies.
Kinugasa & Tsunemi 1999, PASJ, 51, 239. ASCA observations.
Gerardy & Fesen 2001, AJ, 121, 2781. IR spectroscopy and imaging.
DeLaney *et al.* 2002, ApJ, 580, 914. VLA at 1.3 to 1.5 GHz and 5 GHz (7''.2) for spectral index studies.
Morgan *et al.* 2003, ApJ, 597, L33. Sub-mm dust observations.

- Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
Cassam-Chenai *et al.* 2004, A&A, 414, 545. XMM-Newton observations.
Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
Riesgo & López 2005, RMxAA, 41, 57. Optical observations of filament (previously classified as PN, H 2-12).
Blair *et al.* 2007, ApJ, 662, 998. Spitzer observations.
Reynolds *et al.* 2007, ApJ, 668, L135. Chandra observations.
Sankrit *et al.* 2008, AJ, 135, 538. HST observations.
Aharonian *et al.* 2008, A&A, 488, 219. γ -ray upper limit.
Enomoto *et al.* 2008, ApJ, 683, 383. γ -ray upper limit.
Katsuda *et al.* 2008, ApJ, 689, 225. Chandra proper motion studies.
Vink *et al.* 2008, ApJ, 689, 231. Chandra proper motion studies.
Gomez *et al.* 2012, MNRAS, 420, 3557. Herschel IR dust observations.
Williams *et al.* 2012, ApJ, 755, 3. Spitzer spectroscopy.
Burkey *et al.* 2013, ApJ, 764, 63. Chandra observations.
Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
Park *et al.* 2013, ApJ, 767, L10. Suzaku observations.
Katsuda *et al.* 2015, ApJ, 808, 49. XMM-Newton, Chandra and Suzaku observations.
Sankrit *et al.* 2016, ApJ, 817, 36. HST for proper motion studies.
Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G4.8+6.2

RA: 17^h33^m25^s
Dec: -21°34'

1-GHz flux/Jy: 3
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G4.5+6.2.

Radio: Faint shell.

References:

- Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10'.4).
Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (2''.2×1''.3:S=5.5±1.2 Jy), and NVSS at 1.4 GHz.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G5.2–2.6

RA: 18^h07^m30^s
Dec: -25°45'

1-GHz flux/Jy: 2.6?
Spectral index: 0.6?

Size/arcmin: 18
Type: S

Radio: Poorly resolved shell.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4''.3).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G5.4–1.2

RA: 18^h02^m10^s
Dec: −24°54'

1-GHz flux/Jy: 35?
Spectral index: 0.2?

Milne 56
Size/arcmin: 35
Type: C?

Part been called G5.3–1.0. Has been suggested that this is not a SNR.

Radio: Incomplete shell, including wide ‘v’ of emission to east with small flat-spectrum source at apex.

Optical: Detected.

X-ray: Pulsar detected, with faint extension.

Point sources: Pulsar nearby, in flat spectrum source.

Distance: H_I absorption suggests >4.3 kpc.

References:

- Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 38$ Jy).
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz ($3'$).
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8.4'$) and 5 GHz ($4.4'$).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6.8' : S = 21.9 \pm 2.4$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2.6'$).
 Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.
 Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 and 5 GHz.
 Helfand & Becker 1985, Nature, 313, 118. Suggesting it is not a SNR.
 Manchester *et al.* 1985, MNRAS, 212, 975. Pulsar detection.
 Caswell *et al.* 1987, MNRAS, 225, 329. MOST at 843 MHz ($42'' \times 110''$).
 Frail & Kulkarni 1991, Nature, 352, 785. Pulsar and remnant association.
 Manchester *et al.* 1991, MNRAS, 253, 7P. Pulsar and remnant association.
 Milne *et al.* 1992, MNRAS, 255, 707. Parkes 64-m at 4.75 ($4.5' : S = 30.8 \pm 2.1$ Jy) and 8.4 GHz ($3' : S = 24 \pm 3$ Jy), including polarisation.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3.0 \times 4.9' : S = 38$ Jy).
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz ($68'' \times 73''$), plus H_I absorption.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gaensler & Frail 2000, Nature, 406, 158. Pulsar observations, including proper motion.
 Kaspi *et al.* 2001, ApJ, 562, L163. X-ray detection of pulsar, and upper limit for remnant.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.
 Blazek *et al.* 2006, ApJ, 652, 1523. Proper motion study of pulsar.
 Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion study of pulsar.
 Liszt *et al.* 2009, A&A, 508, 1331. CO and IR observations of region.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.

G5.5+0.3

RA: 17^h57^m04^s
Dec: −24°00'

1-GHz flux/Jy: 5.5
Spectral index: 0.7

Size/arcmin: 15×12
Type: S

Has been called G5.55+0.32.

Radio: Shell.

Optical: Detected.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 14.3 \pm 0.3$ Jy), plus other observations.
 Liszt *et al.* 2009, A&A, 508, 1331. CO and IR observations of region.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H_α observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G5.9+3.1

RA: 17^h47^m20^s
Dec: −22°16'

1-GHz flux/Jy: 3.3?
Spectral index: 0.4?

Size/arcmin: 20
Type: S

Radio: Asymmetric shell.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Effelsberg 100-m at 2.7 GHz ($4.3'$).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.1+0.5

RA: $17^{\text{h}}57^{\text{m}}29^{\text{s}}$
Dec: $-23^{\circ}25'$

1-GHz flux/Jy: 4.5
Spectral index: 0.9

Size/arcmin: 18×12
Type: S

Has been called G6.10+0.53.

Radio: Partial shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 13.4 \pm 0.2$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H $_2$ IR observations.

G6.1+1.2

RA: $17^{\text{h}}54^{\text{m}}55^{\text{s}}$
Dec: $-23^{\circ}05'$

1-GHz flux/Jy: 4.0?
Spectral index: 0.3?

Size/arcmin: 30×26
Type: F

Has been called G6.1+1.15.

Radio: Faint, diffuse emission.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4/3), including polarisation.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4/3).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.4–0.1		W28
RA: 18 ^h 00 ^m 30 ^s	1-GHz flux/Jy: 310	Size/arcmin: 48
Dec: −23°26'	Spectral index: varies	Type: C
Has been called G6.6–0.2.		
Radio: Several non-thermal sources in a ring, with flat spectrum core.		
Optical: Diffuse emission.		
X-ray: Diffuse emission from most of the remnant.		
Point sources: Young pulsar near edge of remnant, but not thought to be related.		
Distance: H _I observations suggest 1.9 kpc.		
References:		
Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10 GHz (3').	Rho & Borkowski 2002, ApJ, 575, 201. ROSAT and ASCA observations.	
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical observations.	Claussen <i>et al.</i> 2002, ApJ, 580, 909. Observations of nearby source.	
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).	Yusef-Zadeh <i>et al.</i> 2003, ApJ, 583, 267. OH observations.	
Dopita <i>et al.</i> 1977, ApJ, 214, 179. Some optical line ratios.	Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz of associated OH masers.	
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).	Mavromatakis <i>et al.</i> 2004, A&A, 426, 567. Optical observations.	
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.	Reach <i>et al.</i> 2005, ApJ, 618, 297. Molecular lines and near IR observations.	
Andrews <i>et al.</i> 1983, ApJ, 266, 684. VLA at 1.4 GHz (8'') 4.9 GHz (3'') and 15 GHz (1'') of central region only, plus Einstein image of central region.	Hoffman <i>et al.</i> 2005, ApJ, 620, 257. OH maser observations.	
Bohigas <i>et al.</i> 1983, RMxAA, 8, 155. Optical spectra.	Kawasaki <i>et al.</i> 2005, ApJ, 631, 935. ASCA observations.	
Odenwald <i>et al.</i> 1984, ApJ, 279, 162. Nearby IR source.	Neufeld <i>et al.</i> 2007, ApJ, 664, 890. Spitzer observations.	
Andrews <i>et al.</i> 1985, AJ, 90, 310. VLA of central component.	Aharonian <i>et al.</i> 2008, A&A, 481, 401. γ -ray observations.	
Long <i>et al.</i> 1991, ApJ, 373, 567. Einstein and optical observations.	Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.	
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.1 \times 4'.4 : $S = 660$ Jy).	Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.	
Kaspi <i>et al.</i> 1993, ApJ, 409, L57. Pulsar association.	Guilani <i>et al.</i> 2010, A&A, 516, L11. γ -ray observations.	
Fratil <i>et al.</i> 1993, Nature, 365, 136. VLA at 327 MHz (smoothed to 65''), plus pulsar association.	Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.	
Fratil <i>et al.</i> 1994, ApJ, 424, L111. VLA of associated OH masers.	Abdo <i>et al.</i> 2010, ApJ, 718, 348. Fermi observations.	
Esposito <i>et al.</i> 1996, ApJ, 461, 820. Possible associated γ -ray emission.	Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.	
Fratil <i>et al.</i> 1996, AJ, 111, 1651. OH maser emission.	Sawada <i>et al.</i> 2012, PASJ, 64, 81. Suzaku observations.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	Nichols <i>et al.</i> 2012, MNRAS, 419, 251. CO observations of selected regions.	
Claussen <i>et al.</i> 1997, ApJ, 489, 143. VLA of OH masers.	Gusdorf <i>et al.</i> 2012, A&A, 542, L19. CO observations of regions in NE.	
Claussen <i>et al.</i> 1999, ApJ, 522, 349. High resolution observations of OH masers.	Vaupree <i>et al.</i> 2014, A&A, 568, A50. CO, HCO ⁺ and DCO ⁺ molecular line observations.	
Arikawa <i>et al.</i> 1999, PASJ, 51, L7. Observations of shocked CO.	Pihlstrom <i>et al.</i> 2014, AJ, 147, 73. VLA observations of methanol masers.	
Yusef-Zadeh <i>et al.</i> 2000, ApJ, 540, 842. VLA at 327 MHz (2'.6 \times 5'.5) and 1.48 GHz (40'' \times 65'').	Neufeld <i>et al.</i> 2014, ApJ, 781, 102. Herschel and Spitzer IR spectroscopy.	
Dubner <i>et al.</i> 2000, AJ, 120, 1933. VLA at 328 MHz (97'' \times 52'': $S = 425 \pm 40$ Jy) and 1415 MHz (88'' \times 48'': $S = 246 \pm 20$ Jy), and comparison with other observations.	Hanabata <i>et al.</i> 2014, ApJ, 786, 145. γ -ray observations of region.	
Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.	Zhou <i>et al.</i> 2014, ApJ, 791, 87. XMM-Newton observations.	
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.	Gusdorf <i>et al.</i> 2014, IAUS, 296, 178. CO observations.	
Rowell <i>et al.</i> 2000, A&A, 359, 337. Upper limit on high energy γ -rays.	Nakamura <i>et al.</i> 2014, PASJ, 66, 62. XMM-Newton observations of NE.	
Douvion <i>et al.</i> 2001, A&A, 373, 281. ISO observations.	Froebrich <i>et al.</i> 2015, MNRAS, 454, 2586. H ₂ IR observations.	
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.	Maxted <i>et al.</i> 2016, MNRAS, 462, 532. NH ₃ observations of region.	
Velázquez <i>et al.</i> 2002, AJ, 124, 2145. Parkes 64-m at 1.4 GHz (15') for H _I .	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.	

G6.4+4.0		Size/arcmin: 31
RA: 17 ^h 45 ^m 10 ^s	1-GHz flux/Jy: 1.3?	Type: S
Dec: −21°22'	Spectral index: 0.4?	

Radio: Faint asymmetric shell.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Effelsberg 100-m at 2.7 GHz (4'.3).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.5–0.4

RA: 18^h02^m11^s
Dec: −23°34'

1-GHz flux/Jy: 27
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G6.51–0.48, and part has been called G6.67–0.42.

Radio: Shell, overlapping G6.4–0.1.

Optical: Detected.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 330 MHz (2.6×5.5') and 1.4 GHz (0.7×1.1').
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S=60.8\pm0.4$ Jy), plus other observations.
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observa-

tions.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
Ajello *et al.* 2016, ApJ, 819, 44. γ -ray observations.

G7.0–0.1

RA: 18^h01^m50^s
Dec: −22°54'

1-GHz flux/Jy: 2.5?
Spectral index: 0.5?

Size/arcmin: 15
Type: S

Has been called G7.06–0.12.

Radio: Double rim, brightest in W, confused by bright H II region M20 in SE.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz (2.6×5.5') and 1.48 GHz (40''×65'').
Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz (97''×52'') and 1415 MHz (88''×48'').

G7.2+0.2

RA: 18^h01^m07^s
Dec: −22°38'

1-GHz flux/Jy: 2.8
Spectral index: 0.6

Size/arcmin: 12
Type: S

Has been called G7.20+0.20.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S=5.2\pm0.2$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

1814–24

G7.7–3.7

RA: 18^h17^m25^s
Dec: −24°04'

1-GHz flux/Jy: 11
Spectral index: 0.32

Size/arcmin: 22
Type: S

Radio: Shell, with high polarisation.

References:

Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8.4') and 5 GHz (4.4').
Milne *et al.* 1986, MNRAS, 223, 487. MOST at 843 MHz (44''×108'') and Parkes 64-m at 8.4 GHz (3' : $S=4.6\pm0.5$ Jy),

with polarisation, plus review of flux densities.
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (70''×35'': $S=9.9\pm0.1$ Jy), including polarisation.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G8.3–0.0

RA: $18^{\text{h}}04^{\text{m}}34^{\text{s}}$
Dec: $-21^{\circ}49'$

1-GHz flux/Jy: 1.2
Spectral index: 0.6

Size/arcmin: 5×4
Type: S

Has been called G8.31–0.09.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 2.3 \pm 0.1$ Jy), plus other observations.
Higashi *et al.* 2008, ApJ, 683, 957. γ -ray detection.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G8.7–5.0

RA: $18^{\text{h}}24^{\text{m}}10^{\text{s}}$
Dec: $-23^{\circ}48'$

1-GHz flux/Jy: 4.4
Spectral index: 0.3

Size/arcmin: 26
Type: S

Radio: Asymmetric shell.

References:

Reich *et al.* 1988, IAU Co, 101, 293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4.3'$).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

(W30)

G8.7–0.1

RA: $18^{\text{h}}05^{\text{m}}30^{\text{s}}$
Dec: $-21^{\circ}26'$

1-GHz flux/Jy: 80
Spectral index: 0.5

Size/arcmin: 45
Type: S?

Has been called G8.6–0.1.

Radio: Clumpy non-thermal shell, with low-frequency turnover.

X-ray: Northern edge detected.

Point sources: Pulsar inside western edge.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz ($7.2' \times 9.7' : S = 190 \pm 50$ Jy).
Kassim & Weiler 1990, Nature, 343, 146. VLA at 327 MHz ($3.0' \times 3.7'$).
Kassim & Weiler 1990, ApJ, 360, 184. VLA at 327 MHz ($2.8' \times 4.1' : S = 129 \pm 11$ Jy), and part at 1.4 GHz ($0.9' \times 1.8'$), plus review of flux densities.
Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz ($37'' \times 55''$).
Finley & Ögelman 1994, ApJ, 434, L25. ROSAT observations, including pulsar.
Aharonian *et al.* 2005, Science, 307, 1938. γ -ray detection.

Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations.
Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
Briskin *et al.* 2006, ApJ, 652, 554. Pulsar proper motion.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.
Castro & Slane 2010, ApJ, 717, 372. Fermi observations.
Ajello *et al.* 2012, ApJ, 744, 80. Fermi observations.
Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G8.9+0.4

RA: $18^{\text{h}}03^{\text{m}}58^{\text{s}}$
Dec: $-21^{\circ}03'$

1-GHz flux/Jy: 9
Spectral index: 0.6

Size/arcmin: 24
Type: S

Has been called G8.90+0.40.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 18.2 \pm 0.5$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G9.7–0.0

RA: 18^h07^m22^s
Dec: −20°35'

1-GHz flux/Jy: 3.7
Spectral index: 0.6

Size/arcmin: 15×11
Type: S

Has been called G9.7–0.1 and G9.70–0.06.

Radio: Shell.

References:

- Fratil *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 6.5 \pm 0.2$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.
- Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
Yeung *et al.* 2016, ApJ, 827, 41. γ-ray observations.

G9.8+0.6

RA: 18^h05^m08^s
Dec: −20°14'

1-GHz flux/Jy: 3.9
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Asymmetric shell.

References:

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
Caswell 1983, MNRAS, 204, 833. Molonglo at 408 MHz ($3' : S = 5.8 \pm 0.6$ Jy).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz
- ($41'' \times 63'' : S = 3.5 \pm 0.4$ Jy).
Fratil *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G9.9–0.8

RA: 18^h10^m41^s
Dec: −20°43'

1-GHz flux/Jy: 6.7
Spectral index: 0.4

Size/arcmin: 12
Type: S

Has been called G9.95–0.81.

Radio: Shell.

Optical: Detected.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 11.0 \pm 0.3$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Stupar & Parker 2011, MNRAS, 414, 2282. Hα observations.
- Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G10.5–0.0

RA: 18^h09^m08^s
Dec: −19°47'

1-GHz flux/Jy: 0.9
Spectral index: 0.6

Size/arcmin: 6
Type: S

Has been called G10.59–0.04.

Radio: Partial shell.

X-ray: Possibly detected.

References:

- Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 1.4 \pm 0.1$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G11.0–0.0

RA: $18^{\text{h}}10^{\text{m}}04^{\text{s}}$
Dec: $-19^{\circ}25'$

1-GHz flux/Jy: 1.3
Spectral index: 0.6

Size/arcmin: 11×9
Type: S

Has been called G11.0+0.0 and G11.03–0.05.

Radio: Partial shell.

X-ray: Diffuse emission.

References:

Bamba *et al.* 2003, ApJ, 589, 253. ASCA observations.
Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''),
1.5 GHz (25''), and 74 MHz.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S =$

3.1 ± 0.2 Jy), plus other observations.
Castelletti *et al.* 2016, A&A, 587, A71. VLA at 1.4 GHz
(4''.4 \times 8''.3) and CO observations of region.

G11.1–1.0

RA: $18^{\text{h}}14^{\text{m}}03^{\text{s}}$
Dec: $-19^{\circ}46'$

1-GHz flux/Jy: 5.8
Spectral index: 0.5

Size/arcmin: 18×12
Type: S

Has been called G11.2–1.1 and G11.17–1.04.

Radio: Shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S =$
 11.0 ± 0.3 Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5'':
 $S = 3.40 \pm 0.25$ Jy) including polarisation and review of flux
densities.

G11.1–0.7

RA: $18^{\text{h}}12^{\text{m}}46^{\text{s}}$
Dec: $-19^{\circ}38'$

1-GHz flux/Jy: 1.0
Spectral index: 0.7

Size/arcmin: 11×7
Type: S

Has been called G11.15–0.71.

Radio: Partial shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''),
1.5 GHz (25''), and 74 MHz.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S =$
 2.3 ± 0.1 Jy), plus other observations.

G11.1+0.1

RA: $18^{\text{h}}09^{\text{m}}47^{\text{s}}$
Dec: $-19^{\circ}12'$

1-GHz flux/Jy: 2.3
Spectral index: 0.4

Size/arcmin: 12×10
Type: S

Has been called G11.18+0.11.

Radio: Shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''),
1.5 GHz (25''), and 74 MHz.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S =$

3.5 ± 0.2 Jy), plus other observations.
Castelletti *et al.* 2016, A&A, 587, A71. VLA at 1.4 GHz
(4''.4 \times 8''.3) and CO observations of region.

G11.2–0.3

RA: 18^h11^m27^s
Dec: −19°25'

1-GHz flux/Jy: 22
Spectral index: 0.5

Size/arcmin: 4
Type: C

Probably associated with the SN of AD386.

Radio: Symmetrical clumpy shell, with flatter spectrum core.

X-ray: Shell, with hard spectrum centrally brightened region around pulsar.

Point sources: Central pulsar.

Distance: HI absorption indicates 4.4 kpc.

References:

- Shaver & Goss 1970, *AuJPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Radhakrishnan *et al.* 1972, *ApJS*, 24, 49. HI absorption.
 Becker & Kundu 1975, *AJ*, 80, 679. NRAO 140-ft at 10.6 GHz (3').
 Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Downes 1984, *MNRAS*, 210, 845. VLA at 1465 MHz (20''×25'') and Einstein observations, with review of flux densities.
 Becker *et al.* 1985, *ApJ*, 296, 461. VLA at 1.4 and 5 GHz, plus HI absorption, Einstein observations.
 Morsi & Reich 1987, *A&AS*, 71, 189. Effelsberg 100-m at 32 GHz (26.5': S=4.04±0.24 Jy).
 Green *et al.* 1988, *MNRAS*, 231, 735. VLA at 1.4 and 5 GHz.
 Kassim 1992, *AJ*, 103, 943. VLA at 327 MHz (3.2×4.1': S=39 Jy).
 Reynolds *et al.* 1994, *MNRAS*, 271, L1. ROSAT image and spectra.
 Vasisht *et al.* 1996, *ApJ*, 456, L59. ASCA observations.
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Torii *et al.* 1997, *ApJ*, 489, L145. ASCA detection of pulsar.
 Torii *et al.* 1999, *ApJ*, 523, L69. X-ray timing observations of pulsar.
 Kothes & Reich 2001, *A&A*, 372, 627. Effelsberg 100-m at 4.25, 10.45, 14.9 and 32 GHz (2.5', 1.1', 0.86' and 0.45': S=9.6±0.5, 6.3±0.4, 5.7±0.4 and 3.8±0.4).
 Kaspi *et al.* 2001, *ApJ*, 560, 371. Chandra observations.
 Tam *et al.* 2002, *ApJ*, 572, 202. VLA at 1.4/1.5 GHz (2.6'×1.8':

- S=16.6±0.9 Jy) and 5 GHz (2.1'×1.5': S=8.4±0.9 Jy) for spectral studies.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 14.7 GHz.
 Roberts *et al.* 2003, *ApJ*, 588, 992. Chandra observations.
 Tam & Roberts 2003, *ApJ*, 598, L27. Multi-epoch VLA observations at 1.4/1.5 GHz and 5 GHz, for expansion studies.
 Brogan *et al.* 2004, *AJ*, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.
 Bock & Gaensler 2005, *ApJ*, 626, 343. BIMA at 88.6 GHz (18'').
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Kaplan & Moon 2006, *ApJ*, 644, 1056. IR upper limit for pulsar.
 Koo *et al.* 2007, *ApJ*, 657, 308. IR observations.
 Dean *et al.* 2008, *MNRAS*, 384, L29. INTEGRAL observations of pulsar and nebula.
 Moon *et al.* 2009, *ApJ*, 703, L81. IR spectroscopy.
 Mizuno *et al.* 2010, *AJ*, 139, 1542. Spitzer observations.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9.5': S=9.0±0.5 Jy) including polarisation and review of flux densities.
 Anderson *et al.* 2011, *ApJ*, 742, 7. Spitzer observations.
 Lee *et al.* 2013, *ApJ*, 770, 143. IR observations.
 Froebrich *et al.* 2015, *MNRAS*, 454, 2586. H₂ IR observations.
 Kilpatrick *et al.* 2016, *ApJ*, 816, 1. CO observations, including broad lines.
 Borkowski *et al.* 2016, *ApJ*, 819, 160. Chandra observations.
 Koo *et al.* 2016, *ApJ*, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G11.4–0.1

RA: 18^h10^m47^s
Dec: −19°05'

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 8
Type: S?

Radio: Incomplete shell, possibly with central core.

References:

- Clark *et al.* 1975, *AuJPA*, 37, 1. Molonglo at 408 MHz (3': S=9.4 Jy) contaminated by sidelobes of a nearby source, and Parkes 64-m at 5 GHz (4': S=2.8 Jy).
 Angerhofer *et al.* 1977, *A&A*, 55, 11. NRAO 140-ft at 5 GHz (6.8': S=2.0±0.4 Jy).
 Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Kassim 1992, *AJ*, 103, 943. VLA at 327 MHz (3.2×4.1': S=18 Jy).
 Dubner *et al.* 1993, *AJ*, 105, 2251. VLA at 1.47 GHz

- (44''×63'': S=5.1±0.6 Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Brogan *et al.* 2004, *AJ*, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.
 Rangelov *et al.* 2014, *ApJ*, 796, 34. X-ray upper limit.
 Castelletti *et al.* 2016, *A&A*, 587, A71. VLA at 1.4 GHz (4.4×8'(3)) and CO observations of region.
 Acero *et al.* 2016, *ApJS*, 224, 8. γ-ray observations.

G11.8–0.2

RA: 18^h12^m25^s
Dec: −18°44'

1-GHz flux/Jy: 0.7
Spectral index: 0.3

Size/arcmin: 4
Type: S

Has been called G11.89–0.21.

Radio: Shell.

X-ray: Possibly detected.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 0.9 \pm 0.1$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.0–0.1

RA: 18^h12^m11^s
Dec: −18°37'

1-GHz flux/Jy: 3.5
Spectral index: 0.7

Size/arcmin: 7?
Type: ?

Radio: Incomplete shell, defined in E only.

X-ray: Detected, including possible PWN.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 6.6$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 1.1$ Jy).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'2 × 4'1).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41'' × 61'' : $S = 0.7$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Yamauchi *et al.* 2014, PASJ, 66, 20. Suzaku observations of possible PWN.

G12.2+0.3

RA: 18^h11^m17^s
Dec: −18°10'

1-GHz flux/Jy: 0.8
Spectral index: 0.7

Size/arcmin: 6 × 5
Type: S

Has been called G12.26+0.30.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 1.5 \pm 0.1$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G12.5+0.2

RA: 18^h12^m14^s
Dec: −17°55'

1-GHz flux/Jy: 0.6
Spectral index: 0.4

Size/arcmin: 6 × 5
Type: C?

Has been called G12.58+0.22.

Radio: Diffuse, central brightened.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 0.8 \pm 0.1$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.7–0.0

RA: 18^h13^m19^s
Dec: −17°54'

1-GHz flux/Jy: 0.8
Spectral index: 0.8

Size/arcmin: 6
Type: S

Has been called G12.72–0.00.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 2.0 \pm 0.1$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.8–0.0

RA: 18^h13^m37^s
Dec: −17°49'

1-GHz flux/Jy: 0.8
Spectral index: 0.5

Size/arcmin: 3
Type: C?

Has been called G12.82–0.02 and G12.83–0.02.

Radio: Shell.

X-ray: Diffuse.

Point sources: Central X-ray pulsar.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Brogan *et al.* 2005, ApJ, 629, L105. VLA at 330 MHz ($19'' \times 32''$), plus other observations.
Ubertini *et al.* 2005, ApJ, 629, L109. X-ray, γ -ray and other observations.
Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray detection.
Albert *et al.* 2006, ApJ, 637, L41. γ -ray observations.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S =$

1.2 ± 0.1 Jy), plus other observations.
Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
Funk *et al.* 2007, A&A, 470, 249. XMM-Newton observations, CO observations of surroundings.
Helfand *et al.* 2007, ApJ, 665, 1297. Chandra observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Gotthelf & Halpern 2009, ApJ, 700, L158. Pulsar detection.
Halpern *et al.* 2012, ApJ, 753, L14. Pulsar observations.

G13.3–1.3

RA: 18^h19^m20^s
Dec: −18°00'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 70×40
Type: S?

Radio: Amorphous emission.

Optical: Filaments in S.

X-ray: Elongated emission.

Distance: Absorption indicates 2–4 kpc.

References:

Seward *et al.* 1995, ApJ, 449, 681. ROSAT detection, optical studies and observations of CO.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G13.5+0.2

RA: 18^h14^m14^s
Dec: −17°12'

1-GHz flux/Jy: 3.5?
Spectral index: 1.0?

Size/arcmin: 5×4
Type: S

Has been called G13.46+0.16.

Radio: Elongated, incomplete shell.

References:

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ($S = 0.65 \pm 0.05$ Jy) and 1.4 GHz ($15'' : S = 2.67 \pm 0.5$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G14.1–0.1**RA:** 18^h16^m40^s
Dec: −16°41'**1-GHz flux/Jy:** 0.5
Spectral index: 0.6**Size/arcmin:** 6×5
Type: S

Has been called G14.18–0.12.

Radio: Shell.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.9 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G14.3+0.1**RA:** 18^h15^m58^s
Dec: −16°27'**1-GHz flux/Jy:** 0.6
Spectral index: 0.4**Size/arcmin:** 5×4
Type: S

Has been called G14.30+0.14. Has been suggested this is not an SNR.

Radio: Partial shell.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 1.2 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Pinheiro Gonçalves *et al.* 2011, AJ, 142, 47. IR observations.

G15.1–1.6**RA:** 18^h24^m00^s
Dec: −16°34'**1-GHz flux/Jy:** 5.5?
Spectral index: 0.0?**Size/arcmin:** 30×24
Type: S?**Radio:** Elongated, incomplete shell.**Optical:** Diffuse shell.**References:**

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4.3''$).
 Boumis *et al.* 2008, A&A, 481, 705. Optical detection.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5'' : S = 4.8 \pm 0.3$ Jy) including polarisation and review of flux densities.

G15.4+0.1**RA:** 18^h18^m02^s
Dec: −15°27'**1-GHz flux/Jy:** 5.6
Spectral index: 0.62**Size/arcmin:** 15×14
Type: C?

Has been called G15.42+0.18.

Radio: Shell.**X-ray:** Centrally brightened.**Distance:** H α observations suggest 4.8 kpc.**References:**

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 10.9 \pm 0.3$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5'' : S = 2.1 \pm 0.2$ Jy) including polarisation and review of flux densities.

Castelletti *et al.* 2013, A&A, 557, L15. GMRT at 1.4 GHz ($15''$) plus SGPS H α and CO observations of region.
 Abramowski *et al.* 2014, A&A, 562, A40. γ -ray and XMM-Newton observations.
 Supan *et al.* 2015, A&A, 576, A81. GMRT at 624 MHz ($10''$).

G15.9+0.2

RA: 18^h18^m52^s
Dec: −15°02'

1-GHz flux/Jy: 5.0
Spectral index: 0.63

Size/arcmin: 7×5
Type: S?

Radio: Incomplete shell, with bright concentration to the E.

X-ray: Shell, brighter to S and E.

Point sources: Central X-ray source.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 7.7$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 1.9$ Jy).
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6.8' : $S = 1.1 \pm 0.2$ Jy).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz (58'' × 44'').
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (77'' × 61'' : $S = 11.2 \pm 1.0$ Jy), 1.4 GHz (23'' × 14'' : $S =$

- 3.9 ± 0.1 Jy) and 4.9 GHz (13'' × 16'').
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Reynolds *et al.* 2006, ApJ, 652, L45. Chandra observations.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5' : $S = 2.0 \pm 0.3$ Jy) including polarisation and review of flux densities.
Klochov *et al.* 2016, A&A, 592, L12. Chandra observations.
Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G16.0−0.5

RA: 18^h21^m56^s
Dec: −15°14'

1-GHz flux/Jy: 2.7
Spectral index: 0.6

Size/arcmin: 15×10
Type: S

Has been called G16.05−0.57.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 4.9 \pm 0.2$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Beaumont *et al.* 2011, ApJ, 741, 14. CO of region.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G16.2−2.7

RA: 18^h29^m40^s
Dec: −16°08'

1-GHz flux/Jy: 2.5
Spectral index: 0.4

Size/arcmin: 17
Type: S

Radio: Double rim.

References:

- Trushkin 1999, A&A, 352, L103. Review of radio observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5' : $S = 1.28 \pm 0.10$ Jy) including polarisation and review of flux densities.

G16.4−0.5

RA: 18^h22^m38^s
Dec: −14°55'

1-GHz flux/Jy: 4.6
Spectral index: 0.3?

Size/arcmin: 13
Type: S

Has been called G16.41−0.55.

Radio: Partial shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 10.0 \pm 0.3$ Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5' : $S = 3.0 \pm 0.3$ Jy) including polarisation and review of flux densities.

G16.7+0.1

RA: 18^h20^m56^s
Dec: −14°20'

1-GHz flux/Jy: 3.0
Spectral index: 0.6

Size/arcmin: 4
Type: C

Has been called G16.73+0.08.

Radio: Asymmetric shell with flat-spectrum core.

X-ray: Non-thermal core.

References:

- Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ($S = 0.95$ Jy) and 1.4 GHz ($15'' : S = 2.43$ Jy) and Ooty at 327 MHz ($S = 5.13$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.
 Reynoso & Mangum 2000, ApJ, 545, 874. CO observations of surroundings.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Helfand *et al.* 2003, ApJ, 592, 941. XMM-Newton observations.
 Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz

- ($19'' \times 25''$).
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 6 GHz.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 1.23 \pm 0.11$ Jy) including polarisation and review of flux densities.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G17.0−0.0

RA: 18^h21^m57^s
Dec: −14°08'

1-GHz flux/Jy: 0.5
Spectral index: 0.5

Size/arcmin: 5
Type: S

Has been called G17.02−0.04.

Radio: Shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.7 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G17.4−2.3

RA: 18^h30^m55^s
Dec: −14°52'

1-GHz flux/Jy: 5
Spectral index: 0.5?

Size/arcmin: 24?
Type: S

Radio: Incomplete, poorly defined shell.

Optical: Filaments to SE, and diffuse emission.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4.3'$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2002, A&A, 385, 1042. Optical observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 2.3 \pm 0.2$ Jy) including polarisation and review of flux densities.

G17.4−0.1

RA: 18^h23^m08^s
Dec: −13°46'

1-GHz flux/Jy: 0.4
Spectral index: 0.7

Size/arcmin: 6
Type: S

Has been called G17.48−0.12.

Radio: Partial shell.

References:

- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.9 \pm 0.1$ Jy), plus other observations.
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G17.8–2.6

RA: 18^h32^m50^s
Dec: −14°39'

1-GHz flux/Jy: 5
Spectral index: 0.5

Size/arcmin: 24
Type: S

Radio: Well defined shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 2.23±0.13 Jy) including polarisation and review of flux densities.

G18.1–0.1

RA: 18^h24^m34^s
Dec: −13°11'

1-GHz flux/Jy: 4.6
Spectral index: 0.5

Size/arcmin: 8
Type: S

Has been called G18.1–0.2 and G18.16–0.16.

Radio: Shell.

X-ray: Possibly detected.

Distance: H_I absorption suggests 5.6 kpc.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (8').
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 7.6±0.1 Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Paron *et al.* 2013, MNRAS, 433, 1619. CO, optical and other observations of region.
 Leahy *et al.* 2014, MNRAS, 438, 1813. VGPS H_I observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Voison *et al.* 2016, MNRAS, 458, 2813. Molecular line observations of region.

G18.6–0.2

RA: 18^h25^m55^s
Dec: −12°50'

1-GHz flux/Jy: 1.4
Spectral index: 0.4

Size/arcmin: 6
Type: S

Has been called G18.62–0.28.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 1.9±0.1 Jy), plus other observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including

broad lines.
 Voison *et al.* 2016, MNRAS, 458, 2813. Molecular line observations of region.

G18.8+0.3		Kes 67
RA: 18 ^h 23 ^m 58 ^s	1-GHz flux/Jy: 33	Size/arcmin: 17×11
Dec: −12°23'	Spectral index: 0.46	Type: S
Has been called G18.9+0.3.		
Radio: Incomplete shell, in complex region near the HII region W39.		
Distance: Association with molecular cloud and HI absorption suggests 12 kpc.		
References:		
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 17 \pm 7$ Jy).		
Clark <i>et al.</i> 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 38$ Jy).		
Caswell <i>et al.</i> 1975, A&A, 45, 239. Parkes HI absorption.		
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).		
Milne <i>et al.</i> 1989, PASA, 8, 187. Parkes 64-m at 8.4 GHz ($3'0 : S = 12.9 \pm 1.0$ Jy), including polarisation.		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($2.9 \times 3.5' : S = 55$ Jy).		
Dubner <i>et al.</i> 1996, AJ, 111, 1304. VLA at 1.4 GHz ($75'' \times 55'' : S = 29.9 \pm 0.3$ Jy).		
Dubner <i>et al.</i> 1999, AJ, 118, 930. Parkes 64-m at 1.6 GHz (15'') for HI, VLA at 1.6 GHz (17''×12'') for OH, plus CO observations.		
Dubner <i>et al.</i> 2004, A&A, 426, 201. CO observations of environment.		
Tian <i>et al.</i> 2007, A&A, 474, 541. VGPS at 1.4 GHz (1') including HI, plus CO observations of region.		
Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 15.3 \pm 0.9$ Jy) including polarisation and review of flux densities.		
Vasquez <i>et al.</i> 2012, A&A, 545, A89. CO observations of region.		
Paron <i>et al.</i> 2012, A&A, 547, A60. CO and other molecular observations of region.		
Paron <i>et al.</i> 2015, A&A, 580, A51. CO observations in S.		
G18.9−1.1		
RA: 18 ^h 29 ^m 50 ^s	1-GHz flux/Jy: 37	Size/arcmin: 33
Dec: −12°58'	Spectral index: 0.39	Type: C?
Has been called G18.95−1.1 and G18.94−1.04.		
Radio: Non-thermal, diffuse partially limb-brightened, with central ridge.		
Optical: Detected.		
X-ray: Partial shell.		
Point sources: Compact X-ray source, with diffuse nebula.		
Distance: Various observations suggest 2 kpc.		
References:		
Fürst <i>et al.</i> 1985, Nature, 314, 720. Effelsberg 100-m at 4.75 GHz (2'4: $S = 23.8$ Jy), plus other flux densities.		
Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7.2×8': $S = 82 \pm 15$ Jy), plus review of flux densities.		
Barnes & Turtle 1988, IAUCo, 101, 347. Molonglo at 408 MHz (2.9×3.1': $S = 58 \pm 9$ Jy) and Parkes 64-m at 5 GHz (4'.4×4.1': $S = 23 \pm 6$ Jy).		
Patnaik <i>et al.</i> 1988, Nature, 332, 136. Ooty at 327 MHz (0'.6×1'.6).		
Fürst <i>et al.</i> 1989, A&A, 209, 361. Effelsberg 100-m at 4.75 GHz (2'.45: $S = 23.8$ Jy) and VLA at 1.5 and 4.9 GHz (19''×14''), and Effelsberg 100-m at 1.4 GHz (9') for HI.		
Aschenbach <i>et al.</i> 1991, A&A, 246, L32. ROSAT observations.		
Fürst <i>et al.</i> 1997, A&A, 319, 655. ROSAT observations, and		
Effelsberg 100-m at 10.6 GHz (1'.1).		
Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.		
Harris <i>et al.</i> 2004, ApJ, 603, 152. ROSAT and ASCA observations.		
Tüllmann <i>et al.</i> 2010, ApJ, 720, 848. Chandra detection of compact source.		
Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 19.6 \pm 1.0$ Jy) including polarisation and review of flux densities.		
Stupar & Parker 2011, MNRAS, 414, 2282. Hα observations.		
Froebrich <i>et al.</i> 2015, MNRAS, 454, 2586. H ₂ IR observations.		
Acero <i>et al.</i> 2016, ApJS, 224, 8. γ-ray observations.		
G19.1+0.2		
RA: 18 ^h 24 ^m 56 ^s	1-GHz flux/Jy: 10	Size/arcmin: 27
Dec: −12°07'	Spectral index: 0.5	Type: S
Has been called G19.15+0.27.		
Radio: Partial shell.		
References:		
Brogan <i>et al.</i> 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S = 17.4 \pm 0.4$ Jy), plus other observations.		

G20.0–0.2

RA: 18^h28^m07^s
Dec: −11°35'

1-GHz flux/Jy: 10
Spectral index: 0.1

Size/arcmin: 10
Type: F

Radio: Faint, filled-centre, polarised.

X-ray: Centrally brightened.

Point sources: OH source 20.1–0.1 is nearby.

References:

Becker & Helfand 1985, ApJ, 297, L25. VLA at 1.4 and 5 GHz (12'').
Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7'2×8' : $S = 8.5 \pm 2$ Jy), plus review of flux densities.
Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'.3), including polarisation.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S = 9.2 \pm 0.5$ Jy) including polarisation and review of flux densities.
Petriella *et al.* 2013, A&A, 554, A73. Chandra observations, plus CO and H_i observations of region.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G20.4+0.1

RA: 18^h27^m51^s
Dec: −11°00'

1-GHz flux/Jy: 9?
Spectral index: 0.1?

Size/arcmin: 8
Type: S?

Has been called G20.47+0.16.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 4.2 \pm 0.1$ Jy), plus other observations.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S = 7.5 \pm 0.5$ Jy) including polarisation and review of flux densities.

G21.0–0.4

RA: 18^h31^m12^s
Dec: −10°47'

1-GHz flux/Jy: 1.1
Spectral index: 0.6

Size/arcmin: 9×7
Type: S

Has been called G21.04–0.47.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S = 2.3 \pm 0.2$ Jy), plus other observations.

G21.5–0.9**RA:** 18^h33^m33^s
Dec: −10°35'**1-GHz flux/Jy:** 7
Spectral index: varies**Size/arcmin:** 5
Type: C

Early observations relate to the central core only.

Radio: Filled-centre, with high frequency turnover.

X-ray: Central core, with extended, faint halo.

Point sources: Central pulsar.

Distance: HI absorption indicates 4.6 kpc.

References:

- Wilson & Weiler 1976, A&A, 53, 89. WSRT at 5 GHz (6''×35'').
- Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20''×5'') and 8 GHz (7''×2''), plus review of flux densities.
- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2''.6).
- Becker & Szymkowiak 1981, ApJ, 248, L23. VLA at 5 GHz (8''), and Einstein observations.
- Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT X-ray spectrum, and VLA HI observations.
- Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (26''.5 : $S=5.64\pm0.29$ Jy).
- Fürst *et al.* 1988, PASJ, 40, 347. NRO array at 22.3 GHz (4''.4×7''.3).
- Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'.3), including polarisation.
- Salter *et al.* 1989, A&A, 225, 167. Observations at 90.7 (29''.5 : $S=3.8\pm0.4$ Jy) and 141.9 GHz ($S=2.5\pm1.2$ Jy).
- Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ($S=3.94\pm0.70$ Jy), plus review of flux densities.
- Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
- Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3''.3×3''.4 : $S=9$ Jy).
- Wallace *et al.* 1994, A&A, 286, 565. HI of surroundings.
- Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
- Slane *et al.* 2000, ApJ, 533, L29. Chandra observations identifying X-ray halo.
- Warwick *et al.* 2001, A&A, 365, L248. XMM-Newton observa-
- tions of X-ray halo.
- Bock *et al.* 2001, ApJ, 561, L203. BIMA at 94 GHz (8''.6×4''.6).
- Safi-Harb *et al.* 2001, ApJ, 561, 308. Chandra and other X-ray observations.
- Reich 2002, in NSPS, p1. Effelsberg 100-m at 22 GHz (8'') and 32 GHz, including polarisation.
- La Palombara & Mereghetti 2002, A&A, 383, 916. XMM-Newton upper limit on pulsations.
- Bocchino *et al.* 2005, A&A, 442, 539. XMM-Newton and Chandra observations.
- Gupta *et al.* 2005, CSci, 89, 853. Pulsar discovery.
- Camilo *et al.* 2006, ApJ, 637, 456. Pulsar discovery.
- Bietenholz & Bartel 2008, MNRAS, 386, 1411. VLA at 4.75 GHz (0''.53×0''.82) for expansion studies.
- Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1') including HI.
- Matheson & Safi-Harb 2010, ApJ, 724, 572. Chandra observations.
- Bietenholz *et al.* 2011, MNRAS, 412, 1221. VLA at 1.4 GHz (14''×18'').
- Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 6 GHz.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9''.5 : $S=6.5\pm0.4$ Jy) including polarisation and review of flux densities.
- Zajczyk *et al.* 2012, A&A, 542, A12. IR observations, including polarisation.
- Nynka *et al.* 2014, ApJ, 789, 72. NuSTAR observations.
- Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 70, 100 and 143 GHz.

G21.5–0.1**RA:** 18^h30^m50^s
Dec: −10°09'**1-GHz flux/Jy:** 0.4
Spectral index: 0.5**Size/arcmin:** 5
Type: S

Has been called G21.56–0.10.

Radio: Partial shell.

X-ray: Possibly detected.

References:

- Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
- Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : $S=0.5\pm0.1$ Jy), plus other observations.

G21.6–0.8

RA: 18^h33^m40^s
Dec: −10°25'

1-GHz flux/Jy: 1.4
Spectral index: 0.5?

Size/arcmin: 13
Type: S

Has been called G21.64–0.84.

Radio: Faint, irregular shell.

References:

- Bietenholz *et al.* 2011, MNRAS, 412, 1221. VLA at 327 MHz (85'': $S=2.8$ Jy) and 1.4 GHz (14'' × 18'').
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G21.8–0.6

RA: 18^h32^m45^s
Dec: −10°08'

1-GHz flux/Jy: 65
Spectral index: 0.56

Kes 69
Size/arcmin: 20
Type: S

Radio: Incomplete shell.

X-ray: Detected.

Distance: Association with CO indicates 5.2 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Wilson 1972, A&A, 19, 354. H₂CO absorption.
Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 5 GHz (6') and 10 GHz (3').
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft et 2.7 GHz (5'': $S=42.3\pm4.6$ Jy).
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8/4) and 5 GHz (4/4).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Seward 1990, ApJS, 73, 781. Einstein observations.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.3 × 3.5'': $S = 132$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.
Yusef-Zadeh *et al.* 2003, ApJ, 585, 319. X-ray observations.
- Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
Wood *et al.* 2008, AJ, 135, 2358. VLA at 4.8 GHz, including polarisation.
Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1') including HI.
Zhou *et al.* 2009, ApJ, 691, 516. CO and HCO⁺ of region.
Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5'': $S = 24.0\pm1.3$ Jy) including polarisation and review of flux densities.
Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G22.7–0.2

RA: 18^h33^m15^s
Dec: −09°13'

1-GHz flux/Jy: 33
Spectral index: 0.6

Size/arcmin: 26
Type: S?

Radio: Non-thermal ring in complex region, overlapping G23.3–0.3.

Point sources: Variable radio source near centre, and γ-ray source near edge.

Distance: association with CO indicates 4.4 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.3 × 3.4'': $S = 82$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
- Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Becker *et al.* 2010, AJ, 140, 157. Variable radio source detection.
Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
Su *et al.* 2014, ApJ, 796, 122. CO observations.
Su *et al.* 2015, ApJ, 811, 134. CO observations of region.
Abramowski *et al.* 2015, MNRAS, 446, 1163. γ-ray observations.

G23.3–0.3		W41
RA: 18 ^h 34 ^m 45 ^s	1-GHz flux/Jy: 70	Size/arcmin: 27
Dec: −08°48'	Spectral index: 0.5	Type: S
Radio: Distorted ring, in complex region, overlapping G22.7–0.2.		
X-ray: Possible extended emission, with compact sources.		
Point sources: Pulsar association suggested.		
Distance: H _I and CO observations indicate 4.2 kpc.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.3×3'.4 : $S = 138$ Jy).		
Gaensler & Johnston 1995, MNRAS, 275, L73. Possible pulsar association.		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.		
Aharonian <i>et al.</i> 2005, Science, 307, 1938. γ -ray detection.		
Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer possible detection.		
Aharonian <i>et al.</i> 2006, ApJ, 636, 777. γ -ray observations.		
Albert <i>et al.</i> 2006, ApJ, 643, L53. γ -ray observations.		
Landi <i>et al.</i> 2006, ApJ, 651, 190. X-ray observations.		
Tian <i>et al.</i> 2007, ApJ, 657, L25. VGPS at 1.4 GHz (1') including H _I , plus XMM-Newton observations.		
Leahy & Tian 2008, AJ, 135, 167. VGPS at 1.4 GHz (1') including H _I , plus CO observations.		
Mukherjee <i>et al.</i> 2009, ApJ, 691, 1707. XMM-Newton and γ -ray observations.		
Frail <i>et al.</i> 2013, ApJ, 773, L19. OH observations.		
Castro <i>et al.</i> 2013, ApJ, 774, 36. Fermi observations.		
Abramowski <i>et al.</i> 2015, A&A, 574, A27. γ -ray observations.		
Su <i>et al.</i> 2015, ApJ, 811, 134. CO observations of region.		
Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.		
G23.6+0.3		
RA: 18 ^h 33 ^m 03 ^s	1-GHz flux/Jy: 8?	Size/arcmin: 10?
Dec: −08°13'	Spectral index: 0.3	Type: ?
Has been suggested this is not an SNR.		
Radio: Not well resolved, in complex region.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observa-		
tions.		
Pinheiro Gonçalves <i>et al.</i> 2011, AJ, 142, 47. IR observations.		
Kilpatrick <i>et al.</i> 2016, ApJ, 816, 1. CO observations, including broad lines.		
G24.7–0.6		
RA: 18 ^h 38 ^m 43 ^s	1-GHz flux/Jy: 8	Size/arcmin: 15?
Dec: −07°32'	Spectral index: 0.5	Type: S?
Radio: Incomplete shell, defined in SW.		
References:		
Clark <i>et al.</i> 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 12.3$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 3.6$ Jy).		
Angerhofer <i>et al.</i> 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6.8' : $S = 2.2 \pm 0.5$ Jy).		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at		
4.9 GHz (2'.6).		
Dubner <i>et al.</i> 1993, AJ, 105, 2251. VLA at 1.47 GHz (44''×56'' : $S = 1.9$ Jy).		
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
G24.7+0.6		
RA: 18 ^h 34 ^m 10 ^s	1-GHz flux/Jy: 20?	Size/arcmin: 30×15
Dec: −07°05'	Spectral index: 0.2?	Type: C?
Radio: Filled-centre, with faint shell, and a compact H _{II} region to the S.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).		
Reich <i>et al.</i> 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4'.3' : $S = 19 \pm 3$ Jy) and 4.75 GHz (2'.4' : $S = 17 \pm 4$ Jy) and NRO		
45-m at 10.2 GHz (2'.7' : $S = 15 \pm 3$ Jy).		
Becker & Helfand 1987, ApJ, 316, 660. VLA at 1.4 GHz (12''), and X-ray upper limit.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Froebrich <i>et al.</i> 2015, MNRAS, 454, 2586. H ₂ IR observations.		
Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.		

G25.1–2.3

RA: 18^h45^m10^s
Dec: −08°00'

1-GHz flux/Jy: 8
Spectral index: 0.5?

Size/arcmin: 80×30?
Type: S

Radio: Incomplete shell, extent not well defined.

References:

Gao *et al.* 2011, A&A, 532, A144. Urumqi 25-m at 5 GHz (9'.5:S=3.7±0.4 Jy), plus other observations.

G27.4+0.0

RA: 18^h41^m19^s
Dec: −04°56'

1-GHz flux/Jy: 6
Spectral index: 0.68

4C–04.71
Size/arcmin: 4
Type: S

Early references refer to G27.3–0.1 (Kes 73), a supposed larger remnant.

Radio: Incomplete shell.

X-ray: Diffuse emission, with central low period pulsar.

Point sources: Central AXp.

Distance: H_I absorption suggests 7.5 to 9.8 kpc.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5').
Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3':S=4.4 Jy).
Dickey & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8:S=2.0±0.5 Jy).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz (60''×45'':S=3.5 Jy). Revise $S_{408\text{ MHz}}=10.4$ Jy, and $S_{5\text{ GHz}}=1.9\pm0.2$ Jy.
Kriss *et al.* 1985, ApJ, 288, 703. Einstein observations, plus VLA at 1.4 and 5 GHz (12'').
Sanbonmatsu & Helfand 1992, AJ, 104, 2189. VLA at 1.4 GHz for H_I absorption.
Helfand *et al.* 1994, ApJ, 434, 627. ROSAT observations, particularly of central source.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including nearby masers.
Vasisht & Gotthelf 1997, ApJ, 486, L129. ASCA detection of pulsar.
Gotthelf & Vasisht 1997, ApJ, 486, L133. ASCA observations.
Gotthelf *et al.* 1999, ApJ, 522, L49. X-ray timing observations of pulsar.
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Mereghetti *et al.* 2001, MNRAS, 321, 143. Search for optical/IR counterpart to pulsar.
Kuiper *et al.* 2004, ApJ, 613, 1173. X-ray observations of pulsar.
Tian & Leahy 2008, ApJ, 677, 292. VGPS at 1.4 GHz (1') including H_I.
Mizuno *et al.* 2010, AJ, 139, 1542. Spitzer observations.
An *et al.* 2013, ApJ, 779, 163. NuSTAR and γ-ray observations.
Kumar *et al.* 2014, ApJ, 781, 41. Chandra and XMM-Newton observations.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G27.8+0.6

RA: 18^h39^m50^s
Dec: −04°24'

1-GHz flux/Jy: 30
Spectral index: varies

Size/arcmin: 50×30
Type: F

Radio: Filled-centre, with spectral turnover.

X-ray: Possible pulsar wind nebula.

References:

- Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4'.3:S=23±2 Jy), and 4.75 GHz (2'.4:S=18±2 Jy) and NRO 45-m at 10.2 GHz (smoothed to 4'.3:S=8.5±2 Jy).
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.
Misanovic *et al.* 2010, ApJ, 725, 931. XMM-Newton pulsar/wind nebula search.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5:S=21.0±1.1 Jy) including polarisation and review of flux densities.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G28.6–0.1

RA: 18^h43^m55^s
Dec: −03°53'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 13×9
Type: S

Radio: Poorly defined regions of non-thermal emission.
X-ray: Diffuse shell, with thermal and non-thermal emission.

References:

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 1.4 GHz (15'') and 5 GHz (15'').
Bamba *et al.* 2001, PASJ, 53, L21. ASCA observations.
Ueno *et al.* 2003, ApJ, 588, 338. Chandra observations.
Ebisawa *et al.* 2005, ApJ, 635, 214. Chandra observations.

G28.8+1.5

RA: 18^h39^m00^s
Dec: −02°55'

1-GHz flux/Jy: ?
Spectral index: 0.4?

Size/arcmin: 100?
Type: S?

Radio: Part of rim detected.
X-ray: Diffuse, Centrally brightened.

References:

Schwentker 1994, A&A, 286, L47. ROSAT observations.
Song *et al.* 2000, PASJ, 52, 181. ASCA observations.
Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.
Misanovic *et al.* 2010, ApJ, 725, 931. XMM-Newton pulsar/wind nebula search.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G29.6+0.1

RA: 18^h44^m52^s
Dec: −02°57'

1-GHz flux/Jy: 1.5?
Spectral index: 0.5?

Size/arcmin: 5
Type: S

Radio: Diffuse shell.
Point sources: AXP associated.

References:

Gaensler *et al.* 1999, ApJ, 526, L37. VLA at 5 GHz (13'') and 8 GHz (8'').
Vasisht *et al.* 2000, ApJ, 542, L49. X-ray observations of AXP.
Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G29.7–0.3			Kes 75
RA: 18 ^h 46 ^m 25 ^s	1-GHz flux/Jy: 10	Size/arcmin: 3	
Dec: −02°59'	Spectral index: 0.63	Type: C	
Has erroneously been called G29.6+0.1.			
Radio: Shell with flatter spectrum emission from centre.			
X-ray: Thermal shell and non-thermal core, and central pulsar.			
Point sources: X-ray pulsar.			
Distance: Association with CO implies 11 kpc.			
References:			
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').			tions.
Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (8.4': $S = 5 \pm 20$ Jy).			Mereghetti <i>et al.</i> 2002, ApJ, 574, 873. BeppoSAX observations of pulsar.
Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').			Helfand <i>et al.</i> 2003, ApJ, 582, 783. Chandra observations.
Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20''×7'') and 8 GHz (25''×8''), plus review of flux densities.			Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (10''×13'').
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').			Morton <i>et al.</i> 2007, ApJ, 667, 219. Spitzer and Chandra observations.
Becker <i>et al.</i> 1983, ApJ, 268, L93. VLA at 1.4 GHz (3''), plus Einstein observations.			McBride <i>et al.</i> 2008, A&A, 477, 249. γ -ray and Chandra observations.
Becker & Helfand 1984, ApJ, 283, 154. VLA at 5 GHz (2.6''), plus HI.			Leahy & Tian 2008, A&A, 480, L25. VGPS at 1.4 GHz (1') including HI.
Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26.5': $S = 1.02 \pm 0.07$ Jy).			Kumar & Safi-Harb 2008, ApJ, 678, L43. Chandra observations.
Salter <i>et al.</i> 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.			Ng <i>et al.</i> 2008, ApJ, 686, 508. Chandra observations.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.4×3.7': $S = 27.4$ Jy).			Gavriil <i>et al.</i> 2008, Science, 319, 1802. X-ray observations of pulsar.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.			Su <i>et al.</i> 2009, ApJ, 694, 376. CO observations of region, plus Chandra observations.
Blanton & Helfand 1996, ApJ, 470, 961. ASCA observations.			Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 3.6 \pm 0.6$ Jy) including polarisation and review of flux densities.
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.			Kilpatrick <i>et al.</i> 2016, ApJ, 816, 1. CO observations, including broad lines.
Gotthelf <i>et al.</i> 2000, ApJ, 542, L37. X-ray pulsar detection.			Koo <i>et al.</i> 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
Sugizaki <i>et al.</i> 2001, ApJS, 134, 77. ASCA survey observa-			

G30.7–2.0			
RA: 18 ^h 54 ^m 25 ^s	1-GHz flux/Jy: 0.5?	Size/arcmin: 16	
Dec: −02°54'	Spectral index: 0.7?	Type: ?	
Radio: Poorly defined.			
References:			
Reich <i>et al.</i> 1988, IAUCo, 101, 293. Summary of parameters.			
Reich <i>et al.</i> 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4.3').			
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.			

G30.7+1.0			
RA: 18 ^h 44 ^m 00 ^s	1-GHz flux/Jy: 6	Size/arcmin: 24×18	
Dec: −01°32'	Spectral index: 0.4	Type: S?	
Radio: Non-thermal, highly polarised part shell?			
Point sources: Compact source near centre.			
References:			
Reich <i>et al.</i> 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2.4': $S = 3.4 \pm 0.4$ Jy), plus other flux densities.			Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.6×3.9': $S = 8.6$ Jy).			Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 2.93 \pm 0.19$ Jy) including polarisation and review of flux densities.

G31.5–0.6

RA: 18^h51^m10^s
Dec: −01°31'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 18?
Type: S?

Has been called G31.55–0.65.

Radio: Distorted shell? near HII region.

Optical: Diffuse, incomplete shell.

References:

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4), plus other flux densities.
 Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

G31.9+0.0

RA: 18^h49^m25^s
Dec: −00°55'

1-GHz flux/Jy: 25
Spectral index: varies

3C391

Size/arcmin: 7×5
Type: S

Radio: Shell, brightest in NW, with low frequency turnover.

X-ray: Diffuse with central core.

Distance: HI absorption suggests 7.2 kpc.

References:

- Radhakrishnan *et al.* 1972, ApJS, 24, 49. HI absorption.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
 Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15 GHz (58'' : $S > 1.6$).
 Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (20''×23'') and 8 GHz (24''×9''), plus review of flux densities.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Goss *et al.* 1979, A&A, 78, 75. FIRST at 1.4 GHz (54''×66'': $S = 20 \pm 2$ Jy) and Effelsberg 100-m at 10.7 GHz (77'' : $S = 7.5 \pm 0.8$ Jy).
 Wang & Seward 1984, ApJ, 279, 705. Einstein observations.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.4''×3.7'': $S = 44.8$ Jy).
 Reynolds & Moffett 1993, AJ, 105, 2226. VLA at 1.4 GHz (6''), including possible associated CO.
 Moffett & Reynolds 1994, ApJ, 425, 668. VLA at 330 MHz (smoothed to 30'': $S = 38.5 \pm 0.5$ Jy) 1.46 GHz (6.7') and 4.85 GHz (6.2''×6.4'), including spectral index and polarisation studies.
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.
 Rho & Petre 1996, ApJ, 467, 698. ROSAT observations.
 Wilner *et al.* 1998, AJ, 115, 247. CO observations of surroundings.
 Reach & Rho 1998, ApJ, 507, L93. ISO observations.
 Reach & Rho 1999, ApJ, 511, 836. CO, HCO⁺ and CS observations of surroundings.
 Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
 see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
- Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Chen & Slane 2001, ApJ, 563, 202. ASCA observations.
 Reach *et al.* 2002, ApJ, 564, 302. Observations of shocked molecular species.
 Chen *et al.* 2004, ApJ, 616, 885. Chandra observations.
 Brogan *et al.* 2005, AJ, 130, 148. VLA at 74 MHz, 330 MHz and 1.5 GHz (70'').
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Castro & Slane 2010, ApJ, 717, 372. Fermi observations.
 Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5' : $S = 8.9 \pm 0.6$ Jy) including polarisation and review of flux densities.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Neufeld *et al.* 2014, ApJ, 781, 102. Spitzer and Herschel IR spectroscopy.
 Ergin *et al.* 2014, ApJ, 790, 65. γ -ray and Suzaku observations.
 Gusdorf *et al.* 2014, IAUS, 296, 178. CO observations.
 Su *et al.* 2014, IAUS, 296, 372. VGPS for HI absorption.
 Lee *et al.* 2014, MNRAS, 443, 2650. [FeII] IR survey observations.
 Sato *et al.* 2014, PASJ, 66, 124. Suzaku observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G32.0–4.9	3C396.1
RA: 19 ^h 06 ^m 00 ^s	1-GHz flux/Jy: 22?
Dec: −03°00'	Spectral index: 0.5?
Radio: Possible large shell?	
References:	
Milne & Hill 1969, <i>AuJPh</i> , 22, 211. Parkes 64-m at 635 MHz (31' : $S = 25 \pm 30\%$ Jy), 1410 MHz (15' : $S = 19 \pm 15\%$ Jy) and 2650 MHz (8.4' : $S = 8.6 \pm 30\%$ Jy). Fluxes if size is 60', plus review of flux densities.	Caswell 1970, <i>AuJPh</i> , 23, 105. Revision of low frequency flux densities. Dickel & DeNoyer 1975, <i>AJ</i> , 80, 437. Arecibo at 111 MHz ($S = 105 \pm 30$ Jy).
G32.1–0.9	
RA: 18 ^h 53 ^m 10 ^s	1-GHz flux/Jy: ?
Dec: −01°08'	Spectral index: ?
Radio: Possible faint shell, not well defined.	Size/arcmin: 40?
X-ray: Diffuse, with clumps.	Type: C?
References:	
Folgheraiter <i>et al.</i> 1997, <i>MNRAS</i> , 292, 365. ROSAT and ASCA observations.	
Froebrich <i>et al.</i> 2015, <i>MNRAS</i> , 454, 2586. H ₂ IR observations.	
G32.4+0.1	
RA: 18 ^h 50 ^m 05 ^s	1-GHz flux/Jy: 0.25?
Dec: −00°25'	Spectral index: ?
Has been called G32.45+0.1.	Size/arcmin: 6 Type: S
Radio: Shell.	
X-ray: Shell.	
Distance: X-ray absorption suggests 17 kpc.	
References:	
Yamaguchi <i>et al.</i> 2004, <i>PASJ</i> , 56, 1059. XMM-Newton and other observations.	Acero <i>et al.</i> 2016, <i>ApJS</i> , 224, 8. γ-ray observations.
Ueno <i>et al.</i> 2005, in <i>XRRC</i> , E4.18. XMM-Newton observations.	Kilpatrick <i>et al.</i> 2016, <i>ApJ</i> , 816, 1. CO observations, including broad lines.
G32.8–0.1	Kes 78
RA: 18 ^h 51 ^m 25 ^s	1-GHz flux/Jy: 11?
Dec: −00°08'	Spectral index: 0.2?
Part has been called G33.1–0.1.	Size/arcmin: 17 Type: S?
Radio: Elongated shell?	
Optical: Detected.	
X-ray: Detected.	
Distance: Association with CO indicates 4.8 kpc.	
References:	
Velusamy & Kundu 1974, <i>A&A</i> , 32, 375. NRAO 300-ft et 2.7 GHz (5' : $S = 7.2 \pm 0.5$ Jy).	Green <i>et al.</i> 1997, <i>AJ</i> , 114, 2058. Parkes 64-m OH observations.
Dickel & DeNoyer 1975, <i>AJ</i> , 80, 437. Arecibo at 430 MHz ($S = 19.0 \pm 15.5$ Jy).	Koralesky <i>et al.</i> 1998, <i>AJ</i> , 116, 1323. VLA detection of compact OH emission.
Becker & Kundu 1975, <i>AJ</i> , 80, 679. NRAO 140-ft at 10.6 GHz (3').	Stupar & Parker 2011, <i>MNRAS</i> , 414, 2282. Hα observations.
Caswell <i>et al.</i> 1975, <i>AuJPA</i> , 37, 39. Molonglo at 408 MHz (3' : $S = 12.8$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 7.7$ Jy).	Zhou & Chen 2011, <i>ApJ</i> , 743, 4. XMM-Newton observations, plus CO of region.
Altenhoff <i>et al.</i> 1979, <i>A&AS</i> , 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').	Pihlström <i>et al.</i> 2014, <i>AJ</i> , 147, 73. VLA search for methanol masers.
Kassim 1992, <i>AJ</i> , 103, 943. VLA at 327 MHz (3.6' × 3.8' : $S = 31.3$ Jy).	Froebrich <i>et al.</i> 2015, <i>MNRAS</i> , 454, 2586. H ₂ IR observations.
	Bamba <i>et al.</i> 2016, <i>ApJ</i> , 818, 63. Suzaku observations.

G33.2–0.6

RA: 18^h53^m50^s
Dec: −00°02'

1-GHz flux/Jy: 3.5
Spectral index: varies

Size/arcmin: 18
Type: S

Radio: Incomplete shell.

Optical: Filaments and diffuse emission.

References:

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Reich 1982, A&A, 106, 314. Effelsberg 100-m at 2.7 GHz (4'.4 : $S=2.6\pm0.3$ Jy) and 4.75 GHz (2'.5 : $S=1.75\pm0.2$ Jy).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (68'' \times 52'': $S=2.7\pm0.3$ Jy).
- Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2009, A&A, 499, 789. Optical observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G33.6+0.1

RA: 18^h52^m48^s
Dec: +00°41'

1-GHz flux/Jy: 20
Spectral index: 0.51

Kes 79, 4C00.70, HC13

Size/arcmin: 10
Type: S

Has been called G33.7+0.0.

Radio: Shell, with bright central region, in complex region.

X-ray: Multiple shells and filaments.

Point sources: Central X-ray pulsar.

Distance: H_i absorption gives about 7.8 kpc.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S=35.5$ Jy) and Parkes 64-m at 5 GHz (4' : $S=7.8$ Jy).
 Caswell *et al.* 1975, A&A, 45, 239. H_i absorption.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430} MHz = 69 \pm 33 Jy.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3' : $S=6.8\pm1.5$ Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 : $S=11.4\pm1.1$ Jy).
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1'), plus observations of the nearby point source.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H_i absorption to nearby point source, possibly extragalactic.
 Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source, plus Einstein observations.
 Green 1989, MNRAS, 238, 737. OH absorption.
 Frail & Clifton 1989, ApJ, 336, 854. VLA at 1.4 GHz (1'×2.9'), including H_i absorption.
 Velusamy *et al.* 1991, AJ, 102, 676. VLA at 327 MHz (1'), 1.5 (7'×14'') and 5 GHz (7''), including spectral comparison.
 Green & Dewdney 1992, MNRAS, 254, 686. Observations of adjacent molecular material.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.6×3'.8 : $S=34.8$ Jy).
 Seward & Velusamy 1995, ApJ, 439, 715. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-tions.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emis-sion.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observa-tions.
 Tsunemi & Enoguchi 2002, PASJ, 54, 735. ASCA observations.
 Seward *et al.* 2003, ApJ, 584, 414. Chandra observations.
 Stanimirović 2003, ApJ, 592, 953. Arecibo OH absorption.
 Sun *et al.* 2004, ApJ, 605, 742. Chandra observations.
 Gotthelf *et al.* 2005, ApJ, 627, 390. XMM-Newton pulsar detec-tion.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Halpern *et al.* 2007, ApJ, 665, 1304. X-ray pulsar timing obser-vations.
 Giacani *et al.* 2009, A&A, 507, 841. VLA at 74 MHz (36''×39'' : $S=76\pm10$ Jy), 324 MHz (13'' : $S=39\pm8$ Jy) and 1.5 GHz (17''×19'' : $S=11.5\pm1.5$ Jy), plus review of flux densities and XMM-Newton observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S=9.4\pm0.5$ Jy) including polarisation and review of flux densities.
 Auchettl *et al.* 2014, ApJ, 783, 32. γ -ray detection.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Zhou *et al.* 2016, ApJ, 831, 192. CO and other observations.
 Sato *et al.* 2016, PASJ, 68, S8. Suzaku observations.

G34.7–0.4

RA: 18^h56^m00^s
Dec: +01°22'

1-GHz flux/Jy: 240
Spectral index: 0.37

W44, 3C392

Size/arcmin: 35×27
Type: C

Has been called G34.6–0.5.

Radio: Distorted shell, brighter to the E, with pulsar and associated nebula.

Optical: Diffuse emission.

X-ray: Centrally concentrated, thermal spectrum, plus pulsar wind nebula.

Point sources: Pulsar within the boundary of the remnant.

Distance: H_I absorption indicates ∼ 3 kpc.

References:

- Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10.7 GHz ($3' : S = 105 \pm 7$ Jy).
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 177 \pm 10$ Jy).
Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.
Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 299$ Jy).
Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430} MHz = 540±187 Jy.
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8.4') and 5 GHz (4.4').
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Smith *et al.* 1985, MNRAS, 217, 99. Einstein observations.
Wolszczan *et al.* 1991, ApJ, 372, L99. Pulsar detection.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz ($3.6' \times 3.8' : S = 469$ Jy).
Jones *et al.* 1993, MNRAS, 265, 631. VLA at 1.4 GHz (15'), plus X-ray spectra.
Rho *et al.* 1994, ApJ, 430, 757. Optical and ROSAT observations.
Koo & Heiles 1995, ApJ, 442, 679. H_I of surrounding shell.
Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ-ray emission.
Harrus *et al.* 1996, ApJ, 464, L161. ASCA observations.
Frail *et al.* 1996, ApJ, 464, L165. VLA at 1.5 and 8.4 GHz ($8.9' \times 7.8'$) of pulsar nebula.
Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.
Harrus *et al.* 1997, ApJ, 488, 781. X-ray observations.
Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.
Giacani *et al.* 1997, AJ, 113, 1379. VLA at 1.4 GHz (15'), plus optical images.
Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.
Cox *et al.* 1999, ApJ, 524, 179. Revision of distance.
Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
Petre *et al.* 2002, ApJ, 579, 404. Chandra observations of pulsar and wind nebula.
Mavromatakis *et al.* 2003, A&A, 405, 591. Optical observa-
- tions.
Shelton *et al.* 2004, ApJ, 611, 906. Chandra observations of part.
Reach *et al.* 2005, ApJ, 618, 297. Molecular line and near-IR observations.
Hoffman *et al.* 2005, ApJ, 627, 803. Observations of OH masers.
Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
Castelletti *et al.* 2007, A&A, 471, 537. VLA at 74 MHz ($36'' \times 39'' : S = 634 \pm 70$ Jy) and 324 MHz ($13'' : S = 411 \pm 50$ Jy).
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
Abdo *et al.* 2010, Science, 327, 1103. Fermi observations.
Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 118 \pm 6$ Jy) including polarisation and review of flux densities.
Giuliani *et al.* 2011, ApJ, 742, L30. γ-ray observations.
Uchiyama *et al.* 2012, ApJ, 749, L35. Fermi observations.
Uchida *et al.* 2012, PASJ, 64, 141. Suzaku observations.
Ackermann *et al.* 2013, Science, 339, 807. Fermi observations.
Yoshiike *et al.* 2013, ApJ, 768, 179. CO and H_I observations of region.
Sashida *et al.* 2013, ApJ, 774, 10. HCO⁺ and CO observations of region.
Park *et al.* 2013, ApJ, 777, 14. Arecibo H_I observations of region.
Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
Anderl *et al.* 2014, A&A, 569, A81. CO observations of regions in NE.
Su *et al.* 2014, IAUS, 296, 372. VGPS for H_I absorption.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30, 44 and 70 GHz.
Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
McEwen *et al.* 2016, ApJ, 826, 189. NH₃ and CH₃OH observations.
Acero *et al.* 2016, ApJS, 224, 8. γ-ray observations.

G35.6–0.4

RA: 18^h57^m55^s
Dec: +02°13'

1-GHz flux/Jy: 9
Spectral index: 0.5

Size/arcmin: 15×11
Type: S?

Re-identified as SNR in 2009.

Radio: Diffuse, with some limb brightening.

References:

- Green 2009, MNRAS, 399, 177. Identification in the radio as a SNR.
Paron & Giacani 2010, A&A, 509, L4. CO and IR observations of region.
- Zhu *et al.* 2013, ApJ, 775, 95. H_I and other observations.
Paredes *et al.* 2014, A&A, 561, A56. GMRT at 610 MHz ($12.2'' \times 4.8''$).

G36.6–0.7

RA: 19^h00^m35^s
Dec: +02°56'

1-GHz flux/Jy: 1.0
Spectral index: 0.7?

Size/arcmin: 25?
Type: S?

Radio: polarised arc, possibly part of a larger shell?

References:

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4), plus other flux densities.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.4×3'.7 : $S = 6.7$ Jy).
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S = 0.39 \pm 0.04$ Jy) including polarisation and review of flux densities.

G36.6+2.6

RA: 18^h48^m49^s
Dec: +04°26'

1-GHz flux/Jy: 0.7?
Spectral index: 0.5?

Size/arcmin: 17×13?
Type: S

Radio: Poorly resolved shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

G38.7–1.3

RA: 19^h06^m40^s
Dec: +04°28'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 32×19?
Type: S

G38.7–1.4 refers to the E portion.

Radio: Incomplete shell.

Optical: Arc of filaments, brighter to E.

X-ray: Detected in E.

References:

Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT observations of E.
Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

Huang *et al.* 2014, ApJ, 785, 118. XMM-Newton and Chandra observations.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H $_2$ IR observations.

G39.2–0.3

RA: 19^h04^m08^s
Dec: +05°28'

1-GHz flux/Jy: 18
Spectral index: 0.34

3C396, HC24, NRAO 593

Size/arcmin: 8×6
Type: C

Radio: Shell, brighter to W, with faint ‘tail’ to E.

X-ray: Diffuse, brighter to W, with central core.

Point sources: Central X-ray source.

Distance: H_I absorption suggests >7.7 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3').
 Dickey & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 54 \pm 38$ Jy.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3' : $S = 4.1 \pm 1.0$ Jy).
 Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (65'' × 48'').
 Becker & Helfand 1987, AJ, 94, 1629. VLA at 1.4 GHz (12'' : $S = 14$ Jy) and 5 GHz, plus Einstein observations.
 Patnaik *et al.* 1990, A&A, 232, 467. VLA at 1.5 GHz (25'') and 1.4 GHz (7.8'' × 7.5'') and 5 GHz (25'') including polarisation, plus Ooty at 327 MHz (100'' × 31''), including review of flux densities.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3.5'' × 3.6' : $S = 42.5$ Jy).
 Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz for spectral index studies.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

- Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Harris & Slane 1999, ApJ, 516, 811. ASCA observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Colbert *et al.* 2003, ApJ, 592, L45. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Scaife *et al.* 2007, MNRAS, 377, L69. 33 GHz observations.
 Lee *et al.* 2009, ApJ, 691, 1042. IR observations.
 Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5' : $S = 8.8 \pm 0.5$ Jy) including polarisation and review of flux densities.
 Su *et al.* 2011, ApJ, 727, 43. Chandra and CO observations of region.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Cruciani *et al.* 2016, MNRAS, 459, 4224. Parkes 64 m at 8.4, 13.5, 18.6 and 21.5 GHz, plus review of flux densities.

G39.7–2.0

RA: 19^h12^m20^s
Dec: +04°55'

1-GHz flux/Jy: 85?
Spectral index: 0.7?

W50, SS433

Size/arcmin: 120×60
Type: ?

Eastern part has been called G40.0–3.1. Is this a SNR?

Radio: Elongated shell, containing SS433, adjacent to the H_{II} region S74.

Optical: Faint filaments at the edge of the radio emission.

X-ray: Emission from SS433 and two lobes.

Point sources: SS433 is the compact source in the centre of the W50.

Distance: H_I absorption indicates 6.0 ± 0.5 kpc.

References:

- van den Bergh 1980, ApJ, 236, L23. Optical in H α and [SII].
 Zealey *et al.* 1980, MNRAS, 192, 731. Optical spectra.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H_I absorption to nearby point source (not SS433).
 Sequist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
 Watson *et al.* 1983, ApJ, 273, 688. X-ray observations.
 Downes *et al.* 1986, MNRAS, 218, 393. Effelsberg 100-m at 4.75 GHz (2.4' : $S = 34 \pm 4$ Jy), plus previous 1.7 and 2.7 GHz data.
 Romney *et al.* 1987, ApJ, 321, 822. VLBI of SS433, including distance.
 Elston & Baum 1987, AJ, 94, 1633. Mosaic with VLA at 1.4 GHz (30'') of fine structure only.
 Kawai *et al.* 1989, PASJ, 41, 491. X-ray observations of SS433.
 Band 1989, ApJ, 336, 937. Einstein and EXOSAT observations.
 Yamauchi *et al.* 1994, PASJ, 46, L109. X-ray spectral observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

- Brinkmann *et al.* 1996, A&A, 312, 306. ROSAT observations.
 Kotani *et al.* 1996, PASJ, 48, 619. X-ray line emission from SS433.
 Safi-Harb & Ögelman 1997, ApJ, 483, 868. ROSAT observations.
 Dubner *et al.* 1998, AJ, 116, 1842. VLA at 328 MHz (64'' × 60'': $S = 160 \pm 20$ Jy), and 1.4 GHz (56'' × 54''), plus NRAO 140-ft at 1.4 GHz (21') for H_I observations.
 Safi-Harb & Petre 1999, ApJ, 512, 784. X-ray observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Brinkmann *et al.* 2007, A&A, 463, 611. XMM-Newton observations of E lobe.
 Boumis *et al.* 2007, MNRAS, 381, 308. Optical observations.
 Lockman *et al.* 2007, MNRAS, 381, 881. H_I observations.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5' : $S = 37 \pm 4$ Jy), including polarisation and review of flux densities.

G40.5–0.5

RA: 19^h07^m10^s
Dec: +06°31'

1-GHz flux/Jy: 11
Spectral index: 0.4

Size/arcmin: 22
Type: S

Radio: Shell, brightest to the NE.

References:

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Downes *et al.* 1980, A&A, 92, 47. Effelsberg 100-m at 1.7 GHz (7'.6 : $S = 9.3 \pm 1.3$ Jy), and 2.7 GHz (4'.4 : $S = 7.2 \pm 0.5$ Jy), plus review of flux densities.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.

- Yang *et al.* 2006, ChJAA, 6, 210. CO observations of surroundings.
 Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S = 6.4 \pm 0.3$ Jy) including polarisation and review of flux densities.
 Aliu *et al.* 2014, ApJ, 787, 166. γ -ray observations of region.

G41.1–0.3

RA: 19^h07^m34^s
Dec: +07°08'

1-GHz flux/Jy: 25
Spectral index: 0.50

Size/arcmin: 4.5×2.5
Type: S

3C397

Radio: 3C397 is two sources: the E is the SNR, the W is a HII region.

X-ray: Brighter to the E and W, with central component.

Distance: H α absorption suggest 8 to 10 kpc.

References:

- Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 10 GHz (3' and 5 GHz (6').
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5' : $S = 21.3 \pm 1.2$ Jy for both components).
 Caswell *et al.* 1975, A&A, 45, 239. H α absorption.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 82 \pm 51$ Jy, also Algonquin 46-m at 10.6 GHz (3' : $S = 12 \pm 2$ Jy), and Haystack 36-m at 15.5 GHz (2'.3 : $S = 8.5 \pm 3.0$ Jy).
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S = 29.8$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 8.7$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (58'' × 52'').
 Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz (8''), plus Einstein observations.
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30' : $S = 4.10 \pm 0.19$ Jy).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'.5 × 3'.6 : $S = 46.3$ Jy).
 Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz, for spectral index studies.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Chen *et al.* 1999, ApJ, 520, 737. ASCA and ROSAT observa-

- tions.
 Dyer & Reynolds 1999, ApJ, 526, 365. VLA at 1.5 GHz (6''.9 × 6''.6) and 4.8 GHz (46''.4 × 5''.6), including polarisation and comparison with ROSAT image.
 Safi-Harb *et al.* 2000, ApJ, 545, 922. ROSAT, ASCA and other X-ray observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Safi-Harb *et al.* 2005, ApJ, 618, 321. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Jiang *et al.* 2010, ApJ, 712, 1147. CO observations of region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S = 18.5 \pm 1.1$ Jy) including polarisation and review of flux densities.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
 Yamaguchi *et al.* 2015, ApJ, 801, L31. Suzaku observations.
 Clark *et al.* 2015, ApJ, 809, L2. γ -ray observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Leahy & Ranasinghe 2016, ApJ, 817, 74. VGPS for H α absorption.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G41.5+0.4

RA: 19^h05^m50^s
Dec: +07°46'

1-GHz flux/Jy: 1?
Spectral index: ?

Size/arcmin: 10
Type: S?

Radio: Partial clumpy shell, brightest to NE.

References:

- Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 332 MHz (20'' : $S = 1.8 \pm 0.4$ Jy).
 Alves *et al.* 2012, MNRAS, 422, 2429. Radio observations.

G42.0–0.1

RA: 19^h08^m10^s
Dec: +08°00'

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 8
Type: S?

Radio: Irregular shell.

References:

Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 332 MHz ($20'' : S = 1.8 \pm 0.4$ Jy).
 Alves *et al.* 2012, MNRAS, 422, 2429. Radio observations.

G42.8+0.6

RA: 19^h07^m20^s
Dec: +09°05'

1-GHz flux/Jy: 3?
Spectral index: 0.5?

Size/arcmin: 24
Type: S

Has been called G42.8+0.65.

Radio: Faint shell.

Point sources: Near soft gamma repeater, and young pulsar.

References:

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz ($2.4' : S = 1.5 \pm 0.2$ Jy), plus other flux densities.
 Vasisht *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz ($3.2' \times 3.4'$).
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.

Lorimer & Xilouris 2000, ApJ, 545, 385. Pulsar detection.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz ($50''$), and other observations of the region.

G43.3–0.2

RA: 19^h11^m08^s
Dec: +09°06'

1-GHz flux/Jy: 38
Spectral index: 0.46

W49B

Size/arcmin: 4×3
Type: S

Radio: Shell, brightest to the SE and W, near the HII region W49A.

X-ray: Centrally brightened, elongated E–W.

Point sources: Compact X-ray source.

Distance: HI absorption indicates 10 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ($3'$) and Parkes 64-m at 2.7 GHz ($4'$).
 Downes & Wilson 1974, A&A, 34, 133. Effelsberg 100-m at 10.7 GHz ($1.3'$).
 Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz ($58'' : S = 9.0 \pm 0.7$ Jy).
 Lockhart & Goss 1978, A&A, 67, 355. HI absorption.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz ($2.6'$).
 Pye *et al.* 1984, MNRAS, 207, 649. Einstein observations, and VLA at 1.4 and 5 GHz (both $12''$).
 Smith *et al.* 1985, ApJ, 296, 469. EXOSAT spectrum.
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to $30'' : S = 6.90 \pm 0.38$ Jy).
 Moffett & Reynolds 1994, ApJ, 437, 705. VLA at 330 MHz ($6.7' \times 7.7' : S = 64.4$ Jy), 1.48 GHz ($4.8' \times 5.2' : S = 31.8$ Jy) and 4.85 GHz ($4.0' \times 4.1'$), including polarisation.
 Fujimoto *et al.* 1995, PASJ, 47, L31. ASCA observations.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Hwang *et al.* 2000, ApJ, 532, 970. ROSAT image and ASCA spectroscopy.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. Limit on high energy γ -rays.
 Brogan & Troland 2001, ApJ, 550, 799. VLA at 1.4 GHz

($27'' \times 24''$ and $5''$) for HI Zeeman splitting.
 Lacey *et al.* 2001, ApJ, 559, 954. VLA at 74 MHz ($26'' \times 23'' : S = 55.6$ Jy) and 326 MHz ($6.6' \times 6.2' : S = 56.0$ Jy).
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz ($50''$), and other observations of the region.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Miceli *et al.* 2006, A&A, 453, 567. XMM-Newton observations.
 Keohane *et al.* 2007, ApJ, 654, 938. IR and Chandra observations.
 Ozawa *et al.* 2009, ApJ, 706, L71. Suzaku observations.
 Abdo *et al.* 2010, ApJ, 722, 1303. Fermi observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 19.1 \pm 1.0$ Jy) including polarisation and review of flux densities.
 Rodes-Roca 2013, A&A, 555, A115. IR of compact X-ray source.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
 Lopez *et al.* 2013, ApJ, 777, 145. Chandra spectroscopy.
 Zhu *et al.* 2014, ApJ, 793, 95. Spitzer and other observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Aceri *et al.* 2016, ApJS, 224, 8. γ -ray observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G43.9+1.6

RA: 19^h05^m50^s
Dec: +10°30'

1-GHz flux/Jy: 9.0
Spectral index: 0.5

Size/arcmin: 60?
Type: S?

Radio: Large, poorly defined faint shell.

Point sources: Soft gamma repeater nearby.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
 Vasishth *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3'.2×3'.4).
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50''), and other observations of the region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 4.55±0.24 Jy) including polarisation and review of flux densities.

G45.7–0.4

RA: 19^h16^m25^s
Dec: +11°09'

1-GHz flux/Jy: 4.2?
Spectral index: 0.4?

Size/arcmin: 22
Type: S

Radio: Shell, brightest to the SE, poorly defined to NW.

References:

- Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'.4:S=2.6±0.3 Jy), plus other flux densities.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Acero *et al.* 2016, ApJS, 224, 8. γ-ray observations.

G46.8–0.3

RA: 19^h18^m10^s
Dec: +12°09'

1-GHz flux/Jy: 17
Spectral index: 0.54

(HC30)

Size/arcmin: 17×13
Type: S

Has been called G46.6–0.2.

Radio: Shell, two bright arcs to NNW and SSE.

Distance: H_I absorption suggests 6.8–8.8 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5' : S = 9.8±0.9 Jy), and 37-m at 1.7 GHz (S=14.5±5.5 Jy).
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : S=20.3 Jy) and Parkes 64-m at 5 GHz (4' : S=7.1 Jy).
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S₄₃₀ MHz = 46±21 Jy.
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'.8 : S=5.9±0.6 Jy).
 Sato 1979, ApL, 20, 43. H_I observations.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (58''×53'': S=13.3±0.1 Jy).
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : S = 7.02±0.18 Jy) including polarisation and review of flux densities.

G49.2–0.7		(W51)	
RA: 19 ^h 23 ^m 50 ^s	1-GHz flux/Jy: 160?	Size/arcmin: 30	Type: S?
Dec: +14°06'	Spectral index: 0.3?		
Has erroneously been called G49.1–0.1.			
Radio: In complex region, parameters uncertain.			
Optical: Some diffuse emission possibly associated.			
X-ray: Elongated east–west.			
Distance: Association with CO gives 6 kpc. H _I suggest 4.3 kpc.			
References:			
Shaver & Goss 1970, AuJPA, 14, 133. Parkes 64-m at 5 GHz (4').	Koo <i>et al.</i> 2002, AJ, 123, 1629. ASCA observations.		
Sato 1973, PASJ, 25, 135. H _I absorption.	Koo <i>et al.</i> 2005, ApJ, 633, 946. Chandra observations.		
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5'), $S = 51.5 \pm 3.2$ Jy, for the non-thermal component, but probably confused.	Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H _I . Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.		
Altenhoff <i>et al.</i> 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').	Abdo <i>et al.</i> 2009, ApJ, 706, L1. Fermi observations.		
Seward 1990, ApJS, 73, 781. Einstein observations.	Koo <i>et al.</i> 2010, AJ, 140, 262. H _I Zeeman splitting observations.		
Copetti & Schmidt 1991, MNRAS, 250, 127. 151 MHz observations.	Ceccarelli <i>et al.</i> 2011, ApJ, 740, L4. Molecular line observations of region.		
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'1 × 3'5).	Aleksić <i>et al.</i> 2012, A&A, 541, A13. γ -ray observations.		
Subrahmanyam & Goss 1995, MNRAS, 275, 755. VLA at 330 MHz (1'1).	Hanabata <i>et al.</i> 2013, PASJ, 65, 42. Suzaku observations.		
Koo <i>et al.</i> 1995, ApJ, 447, 211. ROSAT observations.	Tian & Leahy 2013, ApJ, 769, L17. H _I observations of region.		
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant.	Brogan <i>et al.</i> 2013, ApJ, 771, 91. VLA at 74 MHz (84'' × 92'') and 320 MHz (33'' × 35''), plus OH, molecular line and other observations.		
Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.	Park <i>et al.</i> 2013, ApJ, 777, 14. Arecibo of H _I in region.		
Green <i>et al.</i> 1997, AJ, 114, 2058. OH masers.	Sasaki <i>et al.</i> 2014, A&A, 563, A9. XMM-Newton observations.		
Koo & Moon 1997, ApJ, 475, 194. Arecibo (3') and VLA (40'' × 42'') at 1.4 GHz for H _I .	Pihlström <i>et al.</i> 2014, AJ, 147, 73. VLA search for methanol masers.		
Koo & Moon 1997, ApJ, 485, 263. NRAO 12-m CO and HCO ⁺ observations.	Dumes <i>et al.</i> 2014, ApJ, 786, L24. Molecular line observations.		
Brogan <i>et al.</i> 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.	Jogler & Funk 2016, ApJ, 816, 100. γ -ray observations.		
Mavromatakis <i>et al.</i> 2001, A&A, 370, 265. Optical observations.	McEwen <i>et al.</i> 2016, ApJ, 826, 189. NH ₃ and CH ₃ OH observations.		
	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.		
G53.6–2.2		3C400.2, NRAO 611	
RA: 19 ^h 38 ^m 50 ^s	1-GHz flux/Jy: 8	Size/arcmin: 33 × 28	Type: S
Dec: +17°14'	Spectral index: 0.50		
Has been called G53.7–2.2.			
Radio: Ring of emission, with extension to NW.			
Optical: Filaments and diffuse emission.			
X-ray: Centrally brightened, offset to NW.			
Distance: Association with H _I gives 2.8 kpc.			
References:			
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5': $S = 5.3 \pm 0.6$ Jy).	tions.		
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5': $S = 4.8 \pm 0.3$ Jy).	Winkler <i>et al.</i> 1993, ApJ, 405, 608. Optical imaging.		
Clark <i>et al.</i> 1975, AuJPA, 37, 75. Molonglo at 408 MHz (3': $S = 11.7$ Jy).	Dubner <i>et al.</i> 1994, AJ, 108, 207. VLA at 327 MHz (59'') and 1.49 GHz (52''), plus X-rays.		
Dickey & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 20 \pm 10$ Jy, $S_{318\text{ MHz}} = 20 \pm 3.6$ Jy.	Saken <i>et al.</i> 1995, ApJ, 443, 231. ROSAT observations.		
Goss <i>et al.</i> 1975, A&A, 43, 459. WSRT at 610 MHz (1' × 3'): $S = 13.2 \pm 1.6$ Jy).	Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.		
Sabbadin & d'Odorico 1976, A&A, 49, 119. Optical spectra.	Giacani <i>et al.</i> 1998, A&AS, 133, 61. DRAO at 1.4 GHz for H _I studies, including distance.		
van den Bergh 1978, ApJS, 38, 119. Optical observations.	Yoshita <i>et al.</i> 2001, PASJ, 53, 93. ASCA observations, and spectral comparison with ROSAT.		
Rosado 1983, RMxAA, 8, 59. Optical spectra.	Ambrocio-Cruz <i>et al.</i> 2006, RMxAA, 42, 241. Optical imaging and spectroscopy.		
Blair & Long 1988, PASP, 100, 461. Optical imaging and spectroscopy. see also: Blair & Long 1988, PASP, 100, 651. Erratum.	Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 4.00 \pm 0.22$ Jy) including polarisation and review of flux densities.		
Long <i>et al.</i> 1991, ApJ, 373, 567. Einstein and optical observa-	Broersen & Vink 2015, MNRAS, 446, 3885. Chandra observations.		

G54.1+0.3

RA: 19^h30^m31^s
Dec: +18°52'

1-GHz flux/Jy: 0.5
Spectral index: 0.1

Size/arcmin: 12?
Type: C?

Radio: Filled-centre core, with faint diffuse emission.

X-ray: Centrally concentrated, with more extended diffuse emission.

Point sources: Central pulsar.

Distance: H_I absorption suggests 4.5–9 kpc, association with CO suggest 8.2 kpc.

References:

- Green 1985, MNRAS, 216, 691. Radio at 2.7 GHz (7''×20'').
 Reich *et al.* 1985, A&A, 151, L10. Effelsberg 100-m at 4.75 GHz (2':S=0.37±0.04 Jy)
 Velusamy & Becker 1988, AJ, 95, 1162. VLA at 1.4 (14'':S=0.48±0.03 Jy), 1.6 (14'':S=0.42±0.03 Jy) and 5 GHz (5'':S=0.33±0.02 Jy), Ooty at 327 MHz (S=0.50±0.08 Jy), plus review of flux densities.
 Seward 1989, AJ, 97, 481. Einstein observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Lu *et al.* 2001, A&A, 370, 570. ROSAT and ASCA observations.
 Lu *et al.* 2002, ApJ, 568, L49. Chandra observations.
 Camilo *et al.* 2002, ApJ, 574, L71. Pulsar detection.

- Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.
 Leahy *et al.* 2008, AJ, 136, 1477. VGPS at 1.4 GHz (1') including H_I.
 Koo *et al.* 2008, ApJ, 673, L147. Akari observations of surroundings.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Bocchino *et al.* 2010, A&A, 520, A71. XMM-Newton and Suzaku observations.
 Lang *et al.* 2010, ApJ, 709, 1125. VLA at 1.4 GHz (6''6×6''.8), 4.7 GHz (3''.2×3''.3), and 8.2 GHz (3''.0×3''.2) and Spitzer observations.
 Acciari *et al.* 2010, ApJ, 719, L69. γ-ray observations.
 Lee *et al.* 2012, JKAS, 45, 117. CO observations of region.

G54.4–0.3

RA: 19^h33^m20^s
Dec: +18°56'

1-GHz flux/Jy: 28
Spectral index: 0.5

(HC40)
Size/arcmin: 40
Type: S

Has been called G54.5–0.3.

Radio: Shell, in complex region.

Optical: Faint filaments.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5':S=34.4±5.0 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
 Caswell 1985, AJ, 90, 1224. DRAO at 1.4 GHz (1'.3×2'.6:S=18±4 Jy).
 Velusamy *et al.* 1986, JApA, 7, 105. WSRT at 609 MHz (50''×191'' smoothed to 100''×200'').

- Junkes *et al.* 1992, A&AS, 96, 1. Surrounding CO.
 Junkes *et al.* 1992, A&A, 261, 289. Nearby IRAS sources.
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_I.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_I in region.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G55.0+0.3

RA: 19^h32^m00^s
Dec: +19°50'

1-GHz flux/Jy: 0.5?
Spectral index: 0.5?

Size/arcmin: 20×15?
Type: S

Has been called G55.2+0.5.

Radio: Faint, partial shell.

Point sources: Old pulsar nearby.

Distance: Association with H_I features implies 14 kpc.

References:

- Matthews *et al.* 1998, ApJ, 493, 312. WSRT at 327 MHz (1'.0×2'.9:S=0.98±0.15 Jy), DRAO at 1.4 GHz (1'.0×2'.9:S=0.25±0.12 Jy), plus H_I observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G55.7+3.4

RA: 19^h21^m20^s
Dec: +21°44'

1-GHz flux/Jy: 1?
Spectral index: 0.3?

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Point sources: Old pulsar within the boundary of the remnant.

References:

Goss *et al.* 1977, A&A, 61, 93. WSRT observations at 610 MHz (57''×156'': $S = 1.9 \pm 0.2$ Jy) and 1415 MHz (27''×72'': $S = 1.0 \pm 0.1$ Jy).
Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 1.3 to 1.9 GHz

(30'').
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 0.52 \pm 0.03$ Jy) including polarisation and review of flux densities.

G57.2+0.8

RA: 19^h34^m59^s
Dec: +21°57'

1-GHz flux/Jy: 1.8
Spectral index: 0.62

(4C21.53)
Size/arcmin: 12?
Type: S?

Radio: Extended non-thermal arc.

Point sources: Central magnetar.

References:

Sieber & Seiradakis 1984, A&A, 130, 257. Effelsberg 100-m at 1.4 GHz (8'.8: 1.34±0.1), 2.7 GHz (4'.3: 0.86±0.1), plus other surveys of the area.
Caswell *et al.* 1985, AJ, 90, 488. DRAO at 1.4 GHz (1'×3').
Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5':

$S = 0.74 \pm 0.04$ Jy) including polarisation and review of flux densities.
Surnis *et al.* 2016, ApJ, 826, 184. GMRT at 610 MHz and VLA at 1.4 GHz.
Isreal *et al.* 2016, MNRAS, 457, 3448. Chandra, XMM-Newton and Swift observations of magnetar.

G59.5+0.1

RA: 19^h42^m33^s
Dec: +23°35'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 15
Type: S

Has been called G59.6+0.1.

Radio: Incomplete shell.

Optical: Diffuse shell.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1'.0×2.5': $S = 5.1 \pm 0.2$ Jy), and northern sky survey at 4.9 GHz.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.
Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
Xu & Wang 2012, A&A, 543, A24. CO observations of SE.

G59.8+1.2

RA: 19^h38^m55^s
Dec: +24°19'

1-GHz flux/Jy: 1.5
Spectral index: 0.0

Size/arcmin: 20×16?
Type: ?

Has been called G59.7+1.2.

Radio: Poorly defined source.

Optical: Faint diffuse emission and filaments.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'.3), including polarisation.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).

Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 1.43 \pm 0.08$ Jy) including polarisation and review of flux densities.

G63.7+1.1

RA: 19^h47^m52^s
Dec: +27°45'

1-GHz flux/Jy: 1.8
Spectral index: 0.24

Size/arcmin: 8
Type: F

Radio: Centrally brightened, with core.

X-ray: Diffuse emission.

References:

Wallace *et al.* 1997, AJ, 114, 2068. WSRT at 1.4 GHz (14''×26'' : $S = 1.63$ Jy), DRAO at 1.4 GHz (smoothed to 2'), plus review of flux densities and other observations.
Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5' : $S = 1.12 \pm 0.06$ Jy) including polarisation and review of flux densities.
Matheson *et al.* 2016, ApJ, 825, 134. XMM-Newton and Chandra observations.

G64.5+0.9

RA: 19^h50^m25^s
Dec: +28°16'

1-GHz flux/Jy: 0.15?
Spectral index: 0.5

Size/arcmin: 8
Type: S?

Radio: Shell with central source.

References:

Hurley-Walker *et al.* 2009, MNRAS, 398, 249. Radio identification.

G65.1+0.6

RA: 19^h54^m40^s
Dec: +28°35'

1-GHz flux/Jy: 5.5
Spectral index: 0.61

Size/arcmin: 90×50
Type: S

Radio: Large, faint shell.

Point sources: Old pulsar nearby.

Distance: Possible association with H_I suggests 9 kpc.

References:

Landecker *et al.* 1990, A&A, 232, 207. DRAO at 408 MHz (3.5'×7.0' : $S = 9.5 \pm 0.1$ Jy), and 1.4 GHz (1.0'×2.0' : $S = 5.4 \pm 1.0$ Jy).
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Tian & Leahy 2006, A&A, 455, 1053. CGPS at 408 MHz (2.8'×5.9' : $S = 8.6 \pm 0.8$ Jy) and 1.4 GHz (0.8'×1.7' : $S = 4.9 \pm 0.5$ Jy) including H_I.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~

3' : $S = 9.1 \pm 1.0$ Jy) and 1420 MHz ($\sim 1' : S = 3.9 \pm 0.5$ Jy), including review of flux densities.
Aleksić *et al.* 2010, ApJ, 725, 1629. Fermi observations.
Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5' : $S = 3.2 \pm 0.3$ Jy), including polarisation and review of flux densities.
Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G65.3+5.7

RA: 19^h33^m00^s
Dec: +31°10'

1-GHz flux/Jy: 42
Spectral index: 0.6

Size/arcmin: 310×240
Type: S?

Has been called G65.2+5.7.

Radio: Large, faint ring, near S91 and S94.

Optical: Filamentary ring.

X-ray: Diffuse, centrally brightened.

Distance: Optical proper motions and velocities indicates 0.8 kpc.

References:

Gull *et al.* 1977, ApJ, 215, L69. Optical plates.
Reich *et al.* 1979, A&A, 72, 270. Effelsberg 100-m observations at 1.42 GHz (smoothed to 11' : $S = 42.4 \pm 1.6$ Jy), estimate $S_{408\text{ MHz}} = 91 \pm 5$ Jy from previous sky survey.
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
Rosado 1981, ApJ, 250, 222. Optical interferometry.
Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
Seward 1990, ApJS, 73, 781. Einstein observations.
Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
Mavromatakis *et al.* 2002, A&A, 388, 355. Optical observa-

tions.
Boumis *et al.* 2004, A&A, 424, 583. Optical observations.
Shelton *et al.* 2004, ApJ, 615, 275. ROSAT observations.
Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
Xiao *et al.* 2009, A&A, 503, 827. Effelsberg 100-m at 2.7 GHz (4.4' : $S = 22 \pm 3$ Jy), and Urumqi 21-m at 4.8 GHz (9.5' : $S = 16.8 \pm 1.8$ Jy) including polarisation and review of flux densities.
Gosachinskii 2010, AstL, 36, 260. H_I observations.
Kim *et al.* 2010, ApJ, 722, 388. Far UV observations.

G65.7+1.2

RA: 19^h52^m10^s
Dec: +29°26'

1-GHz flux/Jy: 5.1
Spectral index: varies

DA 495

Size/arcmin: 22
Type: F

Has mistakenly been called G55.7+1.2.

Radio: Centrally brightened with thick shell?

X-ray: Centrally brightened.

Point sources: Compact X-ray source near centre.

Distance: HI polarisation observations suggest 1.5 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 2.8 \pm 0.4$ Jy), and 37-m at 1.7 GHz ($S = 4.4 \pm 0.5$ Jy), plus review of flux densities.
see also: Willis 1973, A&A, 27, 483. Erratum.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}} = 8.7 \pm 4.9$ Jy, $S_{318\text{ MHz}} = 9.7 \pm 2.2$ Jy.
 Landecker & Caswell 1983, AJ, 88, 1810. DRAO at 1.4 GHz ($0.9' \times 1.5' : S = 4.4 \pm 0.2$ Jy).
 Velusamy *et al.* 1989, JApA, 10, 161. Ooty at 327 MHz ($36'' \times 64''$), WSRT at 610 MHz ($62''$) and VLA at 1.4 GHz ($36''$ and $12''$), including IRAS imaging.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2004, ApJ, 607, 855. HI polarisation absorption.
 Arzoumanian *et al.* 2004, ApJ, 610, L101. ROSAT and ASCA

- observations of compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 6.5 \pm 0.6$ Jy) and 1420 MHz ($\sim 1' : S = 4.0 \pm 0.2$ Jy), including polarisation and review of flux densities.
 Arzoumanian *et al.* 2008, ApJ, 687, 505. Chandra observations.
 Kothes *et al.* 2008, ApJ, 687, 516. CGPS at 408 MHz ($2.9' \times 6.0' : S = 6.5 \pm 0.5$ Jy) and 1.4 GHz ($0.82' \times 1.75' : S = 4.0 \pm 0.2$ Jy), Effelsberg 100-m at 4.85 GHz ($2.45' : S = 1.6 \pm 0.1$ Jy) and 10.55 GHz (smoothed to $2.45' : S = 1.1 \pm 0.1$ Jy), plus review of flux densities.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 1.95 \pm 0.10$ Jy) including polarisation and review of flux densities.
 Karpova *et al.* 2015, MNRAS, 453, 2241. Chandra and XMM-Newton observations.

G65.8–0.5

RA: 19^h59^m20^s
Dec: +28°38'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 10×6?
Type: S

Radio: Arc in W.

Optical: Diffuse shell, brighter in W.

References:

- Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G66.0–0.0

RA: 19^h57^m50^s
Dec: +29°03'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 31×25?
Type: S

Radio: Some emission in N.

Optical: Incomplete shell.

References:

- Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G67.6+0.9

RA: 19^h57^m45^s
Dec: +30°53'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 50×45?
Type: S

Radio: Arc in S.

Optical: Filamentary shell.

References:

- Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G67.7+1.8

RA: 19^h54^m32^s
Dec: +31°29'

1-GHz flux/Jy: 1.0
Spectral index: 0.61

Size/arcmin: 15×12
Type: S

Radio: Double arc shell.

Optical: Filaments in N.

X-ray: Detected.

Point sources: Compact X-ray source.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz ($1'9 \times 1'0$: $S = 1.9 \pm 0.1$ Jy), and northern sky survey at 4.9 GHz ($S = 0.42 \pm 0.05$ Jy).
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S = 1.1 \pm 0.1$ Jy) and 1420 MHz ($\sim 1'$: $S = 0.68 \pm 0.04$ Jy),

including polarisation and review of flux densities.
Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.
Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
Hui & Becker 2009, A&A, 494, 1005. Chandra observations.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5$: $S = 0.30 \pm 0.03$ Jy) including polarisation and review of flux densities.

G67.8+0.5

RA: 20^h00^m00^s
Dec: +30°51'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 7×5
Type: ?

Radio: Poorly resolved arc.

Optical: Diffuse shell, brighter to W.

References:

Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

G68.6–1.2

RA: 20^h08^m40^s
Dec: +30°37'

1-GHz flux/Jy: 1.1
Spectral index: 0.2

Size/arcmin: 23
Type: ?

Radio: Faint, poorly defined source.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz ($4'3$), including polarisation.
Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4'3$).
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz ($\sim 1'$: $S = 0.57 \pm 0.08$ Jy), including review of flux densities.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5$: $S = 0.80 \pm 0.04$ Jy), including polarisation and review of flux densities.

G69.0+2.7

RA: 19^h53^m20^s
Dec: +32°55'

1-GHz flux/Jy: 120?
Spectral index: varies

CTB 80

Size/arcmin: 80?
Type: ?

An association with a SN in AD1408 has been suggested. Has been called G68.8+2.8.

Radio: Compact core, flat spectrum plateau, and steeper spectrum extensions, with spectral break?

Optical: Expanding nebulosity near centre, with filaments to the SW and far NE.

X-ray: Diffuse emission with compact source.

Point sources: Pulsar at western edge of core.

Distance: H_I observations suggest 1.5 kpc.

References:

- Angerhofer *et al.* 1981, A&A, 94, 313. WSRT at 610 MHz ($56'' \times 103''$) 1.4 GHz ($24'' \times 44''$) and 5 GHz ($7'' \times 13''$), plus optical.
 Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.
 Sofue *et al.* 1983, PASJ, 35, 437. NRO 45-m at 10.2 GHz (2'.7).
 Velusamy & Kundu 1983, JApA, 4, 253. VLA of compact sources.
 Blair *et al.* 1984, ApJ, 282, 161. Optical images and spectra.
 Wang & Seward 1984, ApJ, 285, 607. Einstein observations.
 Strom *et al.* 1984, A&A, 139, 43. Radio observations of flat spectrum component, VLA 5 GHz (1''.7) and 1.4 GHz.
 Mantovani *et al.* 1985, A&A, 145, 50. Bologna at 408 MHz ($2''.6 \times 4''.9 : S = 67.5 \pm 10.5$ Jy), Effelsberg 100-m at 1.41 (9' : $S = 62 \pm 9$ Jy), 1.72 (7''.6 : $S = 66 \pm 5$ Jy), 2.7 (4''.5 : $S = 52 \pm 4$ Jy) and 4.75 GHz (2''.4 : $S = 44 \pm 3.3$ Jy), plus review of flux densities.
 Kulkarni *et al.* 1988, Nature, 331, 50. Pulsar detection.
 Angelini *et al.* 1988, ApJ, 330, L43. EXOSAT spectra.
 Fesen *et al.* 1988, Nature, 334, 229. IRAS of surrounding shell.
 Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4''.3), including polarisation.
 Whitehead *et al.* 1989, MNRAS, 237, 1109. Optical of core.
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
 Hester & Kulkarni 1989, ApJ, 340, 362. Optical imaging and spectroscopy.
 Koo *et al.* 1990, ApJ, 364, 178. Large, expanding H_I shell.
 Greidanus & Strom 1990, A&A, 240, 376. Optical kinematics of core.
 Koo *et al.* 1993, ApJ, 417, 196. VLA at 1.4 GHz of surrounding H_I (1'), plus IRAS.
- Safi-Harb *et al.* 1995, ApJ, 439, 722. ROSAT observations.
 Srinivasan 1997, ApJ, 489, 170. γ -ray observations of pulsar.
 Mavromatakis *et al.* 2001, A&A, 371, 300. Optical observations.
 Butler *et al.* 2002, A&A, 395, 845. HST detection of pulsar.
 Migliazzo *et al.* 2002, ApJ, 567, L141. Pulsar proper motion study.
 Castelletti *et al.* 2003, AJ, 126, 2114. GMRT at 240 and 618 MHz ($26'' \times 17''$ and $10'' \times 6''$) and VLA at 324 and 1380 MHz ($73'' \times 63''$ and $93'' \times 78''$).
 Moon *et al.* 2004, ApJ, 610, L33. Chandra and HST observations of core.
 Li *et al.* 2005, ApJ, 628, 931. Chandra observations of pulsar and surroundings.
 Golden *et al.* 2005, ApJ, 635, L153. High resolution radio observations of pulsar and surroundings.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 72 \pm 7$ Jy) and 1420 MHz ($\sim 1' : S = 56 \pm 5$ Jy), including polarisation and review of flux densities.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_I.
 Albert *et al.* 2007, ApJ, 669, 1143. γ -ray observations.
 Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion of pulsar.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9''.5 : $S = 36 \pm 4$ Jy), including polarisation and review of flux densities.
 Leahy & Ranasinghe 2012, MNRAS, 423, 718. CGPS at 1.4 GHz, including H_I, plus ROSAT observations.
 Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_I in region.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30 and 44 GHz.

G69.7+1.0

RA: 20^h02^m40^s
Dec: +32°43'

1-GHz flux/Jy: 2.0
Spectral index: 0.7

Size/arcmin: 16×14
Type: S

Radio: Poorly resolved source.

X-ray: Detected.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4''.3), including polarisation.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4''.3).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Yoshita *et al.* 2000, PASJ, 52, 867. ROSAT and ASCA observa-
- tions.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.2 \pm 0.4$ Jy) and 1420 MHz ($\sim 1' : S = 1.5 \pm 0.1$ Jy), including review of flux densities.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9''.5 : $S = 0.78 \pm 0.07$ Jy) including polarisation and review of flux densities.

G70.0–21.5**RA:** 21^h24^m00^s
Dec: +19°23'**1-GHz flux/Jy:** ?
Spectral index: ?**Size/arcmin:** 330×240
Type: S**Optical:** Large, faint shell of filaments.**X-ray:** Partially detected.**References:**

- Boumis *et al.* 2002, A&A, 396, 225. Optical and ROSAT observations.
 Fesen *et al.* 2015, ApJ, 812, 37. H α and other optical/UV line and ROSAT observations.

G73.9+0.9**RA:** 20^h14^m15^s
Dec: +36°12'**1-GHz flux/Jy:** 9
Spectral index: 0.23**Size/arcmin:** 27
Type: S?**Radio:** Diffuse, centrally brightened to SW.**Optical:** Faint shell.**References:**

- Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2.4 : $S = 6.7 \pm 0.5$ Jy), plus other flux densities.
 Chastenay & Pineault 1988, IAUCo, 101, 297. DRAO at 408 MHz (3.5×5.9) and 1.4 GHz (1.0×1.7).
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz (3.4×5.8 : $S = 12.7 \pm 1.2$ Jy) and 1.4 GHz (1.0×1.7 : $S = 7.4 \pm 1.0$ Jy).
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Pineault *et al.* 1996, AJ, 112, 201. DRAO at 1.4 GHz (smoothed to 2') for HI.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Mavromatakis 2003, A&A, 398, 153. Optical observations.

- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.0 \pm 1.7$ Jy) and 1420 MHz ($\sim 1' : S = 7.6 \pm 0.6$ Jy), including polarisation and review of flux densities.
 Sitnik 2010, ARep, 54, 317. H α and CO observations of region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5 : $S = 6.2 \pm 0.3$ Jy) including polarisation and review of flux densities.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.
 Zdziarski *et al.* 2016, MNRAS, 455, 1451. γ -ray observations.

G74.0–8.5		Cygnus Loop
RA: 20 ^h 51 ^m 00 ^s	1-GHz flux/Jy: 210	Size/arcmin: 230×160
Dec: +30°40'	Spectral index: varies	Type: S
Has been suggested that this is two overlapping remnants.		
Radio: Shell, brightest to the NE, with fainter breakout region to S, with spectral variations.		
Optical: Large filamentary loop, brightest to the NE, not well defined to the S or W.		
X-ray: Shell in soft X-rays.		
Point sources: Several compact radio sources within the boundary of the remnant, including CL4, plus X-ray sources in S.		
Distance: Optical proper motion and shock velocity gives 0.44 kpc.		
References:		
Green 1990, AJ, 100, 1927. DRAO at 408 MHz (3'3×6.7') for spectral index study, plus X-ray and optical.	2.7 GHz (4'3), with comparison with other data for spectral index studies.	
Graham <i>et al.</i> 1991, AJ, 101, 175. Shocked molecular H outside rim in NE.	Blair <i>et al.</i> 2005, AJ, 129, 2268. HST of outer filaments.	
Shull & Clarke 1991, PASP, 103, 811. Optical spectroscopy of nearside filaments.	Leahy 2005, AJ, 130, 165. DRAO at 1.4 GHz of SE.	
Fesen <i>et al.</i> 1992, AJ, 104, 719. H α imagery.	Levenson & Graham 2005, ApJ, 622, 366. Chandra observations of knot in SE.	
Long <i>et al.</i> 1992, ApJ, 400, 214. Optical of Balmer dominated filament.	Sun <i>et al.</i> 2006, A&A, 447, 937. Urumqi 25-m at 4.8 GHz (9'5': S=90±9 Jy), with comparisons with other data for spectral index studies.	
Arendt <i>et al.</i> 1992, ApJ, 400, 562. IRAS observations.	Kaplan <i>et al.</i> 2006, ApJS, 163, 344. X-ray upper limit on compact sources.	
Hester <i>et al.</i> 1994, ApJ, 420, 721. H α , [OIII] and other optical observations of Balmer dominated filaments in NE.	Seon <i>et al.</i> 2006, ApJ, 644, L175. Far UV observations.	
Gorham <i>et al.</i> 1996, ApJ, 458, 257. Pulsar search.	Sankrit <i>et al.</i> 2007, AJ, 133, 1383. UV observations of part.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	Tsunemi <i>et al.</i> 2007, ApJ, 671, 1717. XMM-Newton observations of NE to SW.	
Levenson <i>et al.</i> 1997, ApJ, 484, 304. ROSAT observations.	Miyata <i>et al.</i> 2007, PASJ, 59, S163. Suzaku observations of NE.	
Leahy <i>et al.</i> 1997, AJ, 114, 2081. DRAO at 1.4 GHz (1'×2'), including polarisation.	Nemes <i>et al.</i> 2008, ApJ, 675, 1293. XMM-Newton observations of NE.	
Leahy & Roger 1998, ApJ, 505, 784. DRAO at 1.4 GHz (1'0×1'9) and 408 MHz (3'4×6'9), for spectral index studies in comparison with other radio observations.	Katsuda <i>et al.</i> 2008, ApJ, 680, 1198. Chandra observations of NE.	
Levenson <i>et al.</i> 1998, ApJS, 118, 541. Optical images.	Uchida <i>et al.</i> 2008, ApJ, 688, 1102. XMM-Newton observations.	
Roger <i>et al.</i> 1999, A&AS, 137, 7. 22 MHz flux density (S=1400±400 Jy).	Uchida <i>et al.</i> 2009, PASJ, 61, 503. Suzaku observations of N.	
Bohigas <i>et al.</i> 1999, ApJ, 518, 324. Optical spectroscopy of surroundings.	Kimura <i>et al.</i> 2009, PASJ, 61, S137. Suzaku observations from NE to SW.	
Levenson <i>et al.</i> 1999, ApJ, 526, 874. ROSAT images.	Tsunemi <i>et al.</i> 2009, PASJ, 61, S147. Suzaku observations of SE.	
Blair <i>et al.</i> 1999, AJ, 118, 942. HST observations, for distance.	Kosugi <i>et al.</i> 2010, PASJ, 62, 1035. Suzaku observations of SE.	
Aschenbach & Leahy 1999, A&A, 341, 602. ROSAT image, and comparison with radio.	Sankrit <i>et al.</i> 2010, ApJ, 712, 1092. Spitzer observations.	
Danforth <i>et al.</i> 2000, AJ, 119, 2319. UV, optical and X-ray comparison of selected regions.	McEntaffer & Brantseg 2011, ApJ, 730, 99. Chandra observations of E.	
Ghavamian <i>et al.</i> 2001, ApJ, 547, 995. Optical spectroscopy.	Katagiri <i>et al.</i> 2011, ApJ, 741, 44. Fermi observations.	
Miyata <i>et al.</i> 2001, ApJ, 550, 1023. ASCA observations of compact X-ray sources.	Katsuda <i>et al.</i> 2012, ApJ, 754, L7. Suzaku and XMM-Newton observations.	
Miyata & Tsunemi 2001, ApJ, 552, 624. ASCA spectroscopy of regions in N and E.	Leahy & Hassan 2013, ApJ, 764, 55. Suzaku and XMM-Newton observations of SW.	
Danforth <i>et al.</i> 2001, AJ, 122, 938. Far-UV spectroscopy, H α and other optical observations of NE region.	Oakley <i>et al.</i> 2013, ApJ, 766, 51. X-ray spectroscopy.	
Levenson & Graham 2001, ApJ, 559, 948. HST of SE region.	Kim <i>et al.</i> 2014, ApJ, 784, 12. Far UV observations.	
Uyaniker <i>et al.</i> 2002, A&A, 389, L61. Effelsberg 100-m at 2.7 GHz (4'3) including polarisation, and comparison with ROSAT data.	Sankrit <i>et al.</i> 2014, ApJ, 787, 3. Spitzer spectroscopy in SE.	
Leahy 2002, AJ, 123, 2689. DRAO at 1.4 GHz (4'×2') for HI.	Medina <i>et al.</i> 2014, ApJ, 791, 30. Optical spectroscopy in NE.	
Blair <i>et al.</i> 2002, ApJS, 140, 367. UV spectroscopy.	Raymond <i>et al.</i> 2015, ApJ, 805, 152. HST observations in NE. see also: Raymond <i>et al.</i> 2015, ApJ, 814, 165. Erratum.	
Welsh <i>et al.</i> 2002, A&A, 391, 705. Optical absorption to background stars.	Roberts & Wang 2015, MNRAS, 449, 1340. Suzaku observations.	
Sankrit & Blair 2002, ApJ, 565, 297. UV observations of NE filament.	Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux density 30 GHz.	
Levenson <i>et al.</i> 2002, ApJ, 576, 798. Chandra observations of W edge.	Katsuda <i>et al.</i> 2016, ApJ, 819, L32. H α spectroscopy of NE.	
Reich <i>et al.</i> 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5': S=184±18 Jy).	Koo <i>et al.</i> 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).	
Leahy 2004, MNRAS, 351, 385. Chandra observations of SW.	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.	
Uyaniker <i>et al.</i> 2004, A&A, 426, 909. Effelsberg 100-m at		

G74.9+1.2		CTB 87
RA: 20 ^h 16 ^m 02 ^s	1-GHz flux/Jy: 9	Size/arcmin: 8×6
Dec: +37°12'	Spectral index: varies	Type: F

Radio: Filled-centre, with high polarisation and high frequency turnover.

X-ray: Centrally brightened.

Point sources: Compact X-ray source in SE.

Distance: H_I absorption indicates 12 kpc, optical extinction gives 6.1 kpc.

References:

- Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430} MHz = 12.2±9.7 Jy, S_{318} MHz = 17.7±5.0 Jy.
 Weiler & Shaver 1978, A&A, 70, 389. WSRT at 610 MHz (57''×94'': $S = 9.1 \pm 1.2$ Jy), 1.4 (24''×40'': $S = 8.7 \pm 1.2$ Jy) and 5 GHz (24''×40'': $S = 5.6 \pm 1.3$ Jy).
 Geldzahler *et al.* 1980, A&A, 84, 237. Effelsberg 100-m at 2.7 GHz (4.4': $S = 7.6 \pm 0.5$ Jy).
 Wilson 1980, ApJ, 241, L19. Einstein observations.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H_I absorption of nearby compact source.
 Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (smoothed to 40'': $S = 1.47 \pm 0.19$ Jy).
 Green & Gull 1989, MNRAS, 237, 555. VLA at 1.4 GHz (1.2'×1.4') including H_I.
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz, plus review of flux densities.
 Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz (3.4'×5.8': $S = 11.6 \pm 0.4$ Jy) and 1.4 GHz (1.0'×1.7': $S = 7.2 \pm 0.3$ Jy).
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Wendker *et al.* 1991, A&A, 241, 551. DRAO at 408 MHz (3.5'×5.2': $S = 13.3 \pm 0.8$ Jy) and Effelsberg 100-m at 4.8 GHz ($S = 7.5 \pm 0.7$ Jy).
 Wallace *et al.* 1994, A&A, 286, 565. H_I of surroundings.

- Cho *et al.* 1994, AJ, 108, 634. CO of adjacent molecular clouds.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Wallace *et al.* 1997, A&A, 317, 212. DRAO at 408 MHz (3.4'×5.5') and 1.4 GHz (1.0'×1.6') including H_I (smoothed to 2').
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2003, ApJ, 588, 852. CGPS at 1.4 GHz (1.6'×1') including H_I, plus CO observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3': $S = 11.9 \pm 0.9$ Jy) and 1420 MHz (~1': $S = 7.1 \pm 1.1$ Jy), including polarisation and review of flux densities.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Sitnik 2010, ARep, 54, 317. H_α and CO observations of region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 6.4 \pm 0.4$ Jy) including polarisation and review of flux densities.
 Matheson *et al.* 2013, ApJ, 774, 33. Chandra observations.
 Bassani *et al.* 2014, A&A, 561, A108. X-ray and γ-ray observations.
 Aliu *et al.* 2014, ApJ, 788, 78. γ-ray detection.
 Saha 2016, MNRAS, 460, 3563. γ-ray observations.
 Acero *et al.* 2016, ApJS, 224, 8. γ-ray observations.

G76.9+1.0		Size/arcmin: 9
RA: 20 ^h 22 ^m 20 ^s	1-GHz flux/Jy: 2?	Type: C
Dec: +38°43'	Spectral index: ?	

Radio: Bipolar shell.

Point sources: Central pulsar.

References:

- Landecker *et al.* 1993, A&A, 276, 522. VLA at 1.49 GHz (14''), 4.86 GHz (13''×16'') and 8.55 GHz (11''×12''), including polarisation and review of flux densities.
 Landecker *et al.* 1997, A&AS, 123, 199. Miyun at 232-MHz (3.8'×5.4').
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3': $S = 2.3 \pm 0.2$ Jy) and 1420 MHz (~1': $S = 1.35 \pm 0.07$ Jy), including polarisation and review of flux densities.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observa-

- tions at 14 to 18 GHz.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5': $S = 0.79 \pm 0.07$ Jy) including polarisation and review of flux densities.
 Marthi *et al.* 2011, MNRAS, 416, 2560. GMRT at 618 MHz (51''×54''), 1160 MHz (2''2×3''4), and Chandra observations of central source.
 Arzoumanian *et al.* 2011, ApJ, 739, 39. Pulsar detection.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G78.2+2.1	DR4, γ Cygni SNR
RA: 20 ^h 20 ^m 50 ^s	1-GHz flux/Jy: 320
Dec: +40°26'	Spectral index: 0.51
Has been called G78.1+1.8.	Size/arcmin: 60
Radio: In complex region (early catalogues refer to other proposed remnants in this region).	Type: S
Optical: Faint filaments, spectra indicate a SNR superposed on a HII region.	
X-ray: Weak emission from the SE of the remnant.	
Point sources: X-ray pulsar at edge of remnant, with nebula.	
Distance: Associations with other objects suggests 1.7 to 2.6 kpc.	
References:	
Higgs <i>et al.</i> 1977, AJ, 82, 718. DRAO at 1.4 GHz ($2' \times 3' : S = 270 \pm 40$ Jy) plus some 10 GHz ($4'$) survey data, reveals true extent of remnant.	Uchiyama <i>et al.</i> 2002, ApJ, 571, 866. ASCA observations.
d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.	Mavromatakis 2003, A&A, 408, 237. Optical observations.
van den Bergh 1978, ApJS, 38, 119. Optical observations.	Bykov 2004, A&A, 427, L21. Hard X-ray observations.
Landecker <i>et al.</i> 1980, A&AS, 39, 133. DRAO HI observations ($2' \times 3'1$).	Becker <i>et al.</i> 2004, ApJ, 615, 897. Chandra and other observations of compact sources.
Higgs <i>et al.</i> 1983, AJ, 88, 97. CO of surroundings.	Weisskopf <i>et al.</i> 2006, ApJ, 652, 387. Chandra and other observations of compact sources.
Bohigas <i>et al.</i> 1983, RMxAA, 8, 155. Optical spectra, find thermal only.	Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 500 \pm 35$ Jy) and 1420 MHz ($\sim 1' : S = 226 \pm 19$ Jy), including review of flux densities.
Braun & Strom 1986, A&AS, 63, 345. WSRT HI observations.	Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity HI.
Fukui & Tatematsu 1988, IAUCo, 101, 261. CO observations of the vicinity ($2'7$).	Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
Green 1989, MNRAS, 238, 737. OH observations.	Ladouceur & Pineault 2008, A&A, 490, 197. CGPS at 408 MHz ($2'9 \times 4'5$) and 1.4 GHz ($0'8 \times 1'5$).
Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz ($3'4 \times 5'8 : S = 480 \pm 60$ Jy) and 1.4 GHz ($1'0 \times 1'7 : S = 270 \pm 40$ Jy).	Gao <i>et al.</i> 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'5 : S = 170 \pm 18$ Jy), including polarisation and review of flux densities.
Wendker <i>et al.</i> 1991, A&A, 241, 551. DRAO at 408 MHz ($3'5 \times 5'2 : S = 540 \pm 40$ Jy) and Effelsberg 100-m at 4.8 GHz ($S = 150 \pm 15$ Jy).	Leahy <i>et al.</i> 2013, MNRAS, 436, 968. ROSAT and Chandra observations, and CGPS for HI.
Esposito <i>et al.</i> 1996, ApJ, 461, 820. Associated γ -ray emission.	Aliu <i>et al.</i> 2013, ApJ, 770, 93. γ -ray observations.
Brazier <i>et al.</i> 1996, MNRAS, 281, 1033. γ -ray and X-ray point source.	Lin <i>et al.</i> 2013, ApJ, 770, L9. Pulsar detection.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	Hui <i>et al.</i> 2015, ApJ, 799, 76. XMM-Newton and Chandra observations of pulsar, including proper motion.
Zhang <i>et al.</i> 1997, A&A, 324, 641. Multi-frequency radio comparison.	Fraija & Araya 2016, ApJ, 826, 31. γ -ray observations.
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.	

G82.2+5.3	W63
RA: 20 ^h 19 ^m 00 ^s	Size/arcmin: 95×65
Dec: +45°30'	Type: S
Has been called G82.5+5.3.	
Radio: Shell in the Cygnus X complex.	
Optical: In complex region, but spectra indicate SNR filaments.	
X-ray: Detected.	
References:	
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 59.0 \pm 3.5$ Jy).	Uyaniker <i>et al.</i> 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation, of part.
Sabbadin 1976, A&A, 51, 159. Optical spectra.	Mavromatakis <i>et al.</i> 2004, A&A, 415, 1051. ROSAT, ASCA and optical observations.
Angerhofer <i>et al.</i> 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'8$). Incomplete mapping.	Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 144 \pm 12$ Jy) and 1420 MHz ($\sim 1' : S = 93 \pm 5$ Jy), including review of flux densities.
Rosado & González 1981, RMxAA, 5, 93. Optical spectra.	Gao <i>et al.</i> 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'5 : S = 49 \pm 5$ Jy), including polarisation and review of flux densities.
Seward 1990, ApJS, 73, 781. Einstein observations.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.	
Reich <i>et al.</i> 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'5 : S = 82.5 \pm 5.5$ Jy).	

G83.0–0.3

RA: 20^h46^m55^s
Dec: +42°52'

1-GHz flux/Jy: 1
Spectral index: 0.4

Size/arcmin: 9×7
Type: S

Radio: Incomplete shell.

References:

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 1.2 \pm 0.3$ Jy) and 1420 MHz ($\sim 1' : S = 0.8 \pm 0.1$ Jy, including polarisation and review of flux densities).

G84.2–0.8

RA: 20^h53^m20^s
Dec: +43°27'

1-GHz flux/Jy: 11
Spectral index: 0.5

Size/arcmin: 20×16
Type: S

Radio: Elongated shell, with a filament aligned with the major axis.

X-ray: Detected.

Distance: H_I absorption suggests 6 kpc.

References:

Matthews *et al.* 1977, A&A, 55, 1. WSRT at 610 MHz ($56'' \times 81'' : S = 12.4 \pm 1.5$ Jy) and Effelsberg 100-m at 2.7 GHz ($4.4' : S = 6.8 \pm 1.3$ Jy).
 Matthews & Shaver 1980, A&A, 87, 255. WSRT at 1415 MHz ($23'' \times 32''$), and Effelsberg 100-m at 2.7 GHz ($4.4' : S = 5.6 \pm 0.5$ Jy).
 Feldt & Green 1993, A&A, 274, 421. DRAO at 1.4 GHz ($1' \times 1.5'$), including H_I, plus CO observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) in-

cluding polarisation.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 14.5 \pm 0.5$ Jy) and 1420 MHz ($\sim 1' : S = 7.2 \pm 0.8$ Jy), including review of flux densities.
 Leahy & Green 2012, ApJ, 760, 25. CGPS, including H_I, plus Chandra observations.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G85.4+0.7

RA: 20^h50^m40^s
Dec: +45°22'

1-GHz flux/Jy: ?
Spectral index: 0.2

Size/arcmin: 24?
Type: S

Radio: Faint, incomplete shell, within larger thermal shell.

X-ray: Centrally brightened.

Distance: H_I observations suggest 3.5 kpc.

References:

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($4.4' \times 2.8' : S < 0.45$ Jy) and 1.4 GHz ($1.1' \times 0.8'$), plus H_I, X-ray and optical data.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 2.9 \pm 0.5$ Jy) and 1420 MHz ($\sim 1' : S = 2.3 \pm 0.2$ Jy),

including review of flux densities.
 Jackson *et al.* 2008, ApJ, 674, 936. XMM-Newton and H_I observations.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G85.9–0.6

RA: 20^h58^m40^s
Dec: +44°53'

1-GHz flux/Jy: ?
Spectral index: 0.2

Size/arcmin: 24
Type: S

Radio: Faint, incomplete shell.

Optical: Diffuse shell.

X-ray: Centrally brightened.

Distance: H_I observations suggest 4.8 kpc.

References:

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($4.4' \times 2.8' : S < 0.9$ Jy) and 1.4 GHz ($1.1' \times 0.8'$), plus H_I, X-ray and optical data.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.0 \pm 1.3$ Jy) and 1420 MHz ($\sim 1' : S = 2.2 \pm 0.8$ Jy), including review of flux densities.

Jackson *et al.* 2008, ApJ, 674, 936. XMM-Newton and H_I observations.
 Gök *et al.* 2009, Ap&SS, 324, 17. Optical observations.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G89.0+4.7		HB21
RA: 20 ^h 45 ^m 00 ^s	1-GHz flux/Jy: 220	Size/arcmin: 120×90
Dec: +50°35'	Spectral index: 0.38	Type: S
Radio: Distorted shell (4C50.52, an extragalactic double, is within the boundary of the remnant).		
Optical: Filaments and patches.		
X-ray: Centrally brightened.		
Distance: Various associations imply 0.8 kpc.		
References:		
Hirabayashi & Takahashi 1972, PASJ, 24, 231. 30-m dish at 4.2 GHz ($11' : S = 160 \pm 40$ Jy).		
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 148 \pm 16$ Jy), plus optical filaments.		
Hill 1974, MNRAS, 169, 59. Half-Mile Telescope at 1.4 GHz ($3' \times 3.9'$).		
Haslam <i>et al.</i> 1975, A&A, 39, 453. Effelsberg 100-m at 2.7 GHz (4.4').		
Fukui & Tatematsu 1988, IAUCo, 101, 261. CO observations of the vicinity (2.7').		
Tatematsu <i>et al.</i> 1990, A&A, 237, 189. DRAO at 408 MHz ($3.5' \times 4.5'$) and 1.4 GHz ($1.0' \times 1.3'$), including H _i , plus CO observations of adjacent molecular cloud.		
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		
Leahy & Aschenbach 1996, A&A, 315, 260. ROSAT observations.		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
Koo <i>et al.</i> 2001, ApJ, 552, 175. NRAO 12-m and other CO observations ($27''$ and $45''$) of eastern part.		
Reich <i>et al.</i> 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14.5' : S = 228 \pm 5$ Jy).		
Uyaniker <i>et al.</i> 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1') including polarisation.		
Byun <i>et al.</i> 2006, ApJ, 637, 283. CO observations of surround-		
ings.		
Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.		
Leahy 2006, ApJ, 647, 1125. CGPS at 408 MHz ($2.8' \times 3.7'$) and 1.4 GHz ($0.8' \times 1.1'$).		
Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 259 \pm 19$ Jy) and 1420 MHz ($\sim 1' : S = 183 \pm 9$ Jy), including polarisation and review of flux densities.		
Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H _i .		
Mavromatakis <i>et al.</i> 2007, A&A, 461, 991. Optical observations.		
Shinn <i>et al.</i> 2009, ApJ, 693, 1883. IR observations.		
Pannuti <i>et al.</i> 2010, AJ, 140, 1787. ASCA and observations.		
Shinn <i>et al.</i> 2010, AdSpR, 45, 445. IR observations in S.		
Gao <i>et al.</i> 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9.5' : S = 107 \pm 11$ Jy), including polarisation and review of flux densities.		
Reichardt <i>et al.</i> 2012, A&A, 546, A21. γ -ray detection.		
Shinn <i>et al.</i> 2012, ApJ, 759, 34. Akari observations of H ₂ .		
Pivato <i>et al.</i> 2013, ApJ, 779, 179. Fermi observations.		
Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux densities at 30 and 44 GHz.		
Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.		
G93.3+6.9		DA 530, 4C(T)55.38.1
RA: 20 ^h 52 ^m 25 ^s	1-GHz flux/Jy: 9	Size/arcmin: 27×20
Dec: +55°21'	Spectral index: 0.45	Type: C?
Has been called G93.2+6.7.		
Radio: Shell, with two bright limbs, highly polarised.		
X-ray: Compact central source.		
Distance: H _i observations suggest 2.2 kpc.		
References:		
Roger & Costain 1976, A&A, 51, 151. DRAO at 1.42 GHz ($2' \times 2.4' : S = 6.9$ Jy).		
Haslam <i>et al.</i> 1980, A&A, 92, 57. Effelsberg 100-m at 1.72 GHz ($7.6' : S = 6.47 \pm 0.52$ Jy) and 2.7 GHz (4.4' : $S = 5.64 \pm 0.64$ Jy), plus review of flux densities.		
Lalitha <i>et al.</i> 1984, A&A, 131, 196. Effelsberg 100-m at 4.75 GHz (smoothed to 3' : $S = 4.01 \pm 0.57$ Jy).		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
Landcker <i>et al.</i> 1999, ApJ, 527, 866. DRAO at 408 MHz ($3.5' \times 4.3'$) and 1.4 GHz ($1.0' \times 1.2'$), including polarisation and H _i .		
Foster & Routledge 2003, ApJ, 598, 1005. H _i for distance.		
Kaplan <i>et al.</i> 2004, ApJS, 153, 269. Chandra limits for any compact source.		
Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.5 \pm 0.7$ Jy) and 1420 MHz, including review of flux densities.		
Jiang <i>et al.</i> 2007, ApJ, 670, 1142. Chandra observations.		
Bocchino <i>et al.</i> 2008, AdSpR, 41, 407. XMM-Newton observations.		
Jeong <i>et al.</i> 2012, Ap&SS, 342, 389. CO observations of region.		

G93.7–0.2

RA: 21^h29^m20^s
Dec: +50°50'

1-GHz flux/Jy: 65
Spectral index: 0.65

Has been called G93.6–0.2 and G93.7–0.3.

Radio: Distorted, faint shell.

Distance: Association with H_I features suggests 1.5 kpc.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 18.4 \pm 1.0$ Jy).
 Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ($7.6' : S = 53.5 \pm 5.0$ Jy), plus review of flux densities.
 Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 58 \pm 6$ Jy).
 Mantovani *et al.* 1991, A&A, 247, 545. Effelsberg 100-m at 4.75 GHz (smoothed to $3' : S = 33.5 \pm 4.0$ Jy), including polarisation, plus review of flux densities.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emis-

CTB 104A, DA 551

Size/arcmin: 80
Type: S

G94.0+1.0

RA: 21^h24^m50^s
Dec: +51°53'

1-GHz flux/Jy: 13
Spectral index: 0.45

3C434.1

Size/arcmin: 30×25
Type: S

Radio: Incomplete shell, containing H_I shell.

Distance: Association with stellar wind bubble implies 5.2 kpc.

References:

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 6.1 \pm 0.8$ Jy), and 37-m at 1.7 GHz ($S = 11 \pm 3$ Jy).
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 5.8 \pm 0.4$ Jy). Also NRAO 140-ft at 5 GHz ($6'$).
 Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz ($7.6' : S = 12.0 \pm 1.3$ Jy), plus review of flux densities.
 Goss *et al.* 1984, A&A, 138, 469. WSRT at 610 MHz (smoothed to $100'' : S = 16 \pm 1.7$ Jy) and Effelsberg 100-m at 4.75 GHz ($2.4' : S = 7.2 \pm 0.5$ Jy).
 Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to $2' : S = 16 \pm 3$ Jy).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation.

- Foster *et al.* 2004, A&A, 417, 79. DRAO at 1.4 GHz, including H_I.
 Foster 2005, A&A, 441, 1043. CGPS at 408 MHz ($2.8 \times 3.6'$) and 1.4 GHz ($0.8 \times 1.0'$) for spectral index studies, plus other observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 20 \pm 2$ Jy) and 1420 MHz ($\sim 1' : S = 11.3 \pm 1.0$ Jy), including review of flux densities.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 6.2 \pm 0.4$ Jy) including polarisation and review of flux densities.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
 Jeong *et al.* 2013, ApJ, 770, 105. CO observations of region.

G96.0+2.0

RA: 21^h30^m30^s
Dec: +53°59'

1-GHz flux/Jy: 0.35
Spectral index: 0.6

Size/arcmin: 26
Type: S

Radio: Faint, arc in S, poorly defined in N.

Distance: Association for H_I indicates 4 kpc.

References:

- Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz ($2.8 \times 3.5'$) and 1.4 GHz ($50'' \times 61''$) including H_I.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 0.42 \pm 0.06$ Jy) and 1420 MHz ($\sim 1' : S = 0.24 \pm 0.02$ Jy),

- including review of flux densities.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 0.14 \pm 0.02$ Jy) including polarisation and review of flux densities.

G106.3+2.7

RA: 22^h27^m30^s
Dec: +60°50'

1-GHz flux/Jy: 6
Spectral index: 0.6

Size/arcmin: 60×24
Type: C?

Incorporates the pulsar wind nebula G106.6+2.9 (the ‘Boomerang’).

Radio: Faint extended source, which brighter ‘head’ to NE.

X-ray: Pulsar and wind nebula.

Point sources: Pulsar.

References:

- Pineault & Joncas 2000, AJ, 120, 3218. DRAO at 408 MHz ($3'9 \times 3'5 : S = 10.5 \pm 0.3$ Jy) and 1.4 GHz ($1'2 \times 1'0 : S = 4.9 \pm 0.6$ Jy), plus HI.
 Halpern *et al.* 2001, ApJ, 547, 323. X-ray and radio observations of the ‘head’.
 Halpern *et al.* 2001, ApJ, 552, L125. Pulsar detection.
 Kothes *et al.* 2001, ApJ, 560, 236. CGPS at 1.4 GHz, including HI, plus CO and other observations.
 Ng & Romani 2004, ApJ, 601, 479. Chandra detection of pulsar wind nebula.
 Kothes *et al.* 2004, ApJ, 607, 855. HI polarisation absorption.
 Kothes *et al.* 2006, ApJ, 638, 225. Effelsberg 100-m at 4.85

- ($2'4$), 8.35 ($1'4$), 10.5 ($1'2$) and 32 GHz ($0'45$) of pulsar wind nebula, including polarisation.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 8.6 \pm 1.0$ Jy) and 1420 MHz ($\sim 1' : S = 4.8 \pm 0.5$ Jy), including polarisation and review of flux densities.
 Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Acciari *et al.* 2009, ApJ, 703, L6. γ -ray observations.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'5 : S = 2.0 \pm 0.3$ Jy), including polarisation and review of flux densities.

G108.2–0.6

RA: 22^h53^m40^s
Dec: +58°50'

1-GHz flux/Jy: 8
Spectral index: 0.5

Size/arcmin: 70×54
Type: S

Radio: Faint shell.

Distance: Possible associated HI structures suggest 3.2 kpc.

References:

- Tian *et al.* 2007, A&A, 465, 907. DRAO at 408 MHz ($2'8 \times 3'3 : S = 11.5 \pm 1.2$ Jy) and 1.4 GHz ($1'0 \times 1'2 : S = 6.6 \pm 0.7$ Jy) including HI.

G109.1–1.0		CTB 109
RA: 23 ^h 01 ^m 35 ^s	1-GHz flux/Jy: 22	Size/arcmin: 28
Dec: +58°53'	Spectral index: 0.45	Type: S
Radio: Semicircular shell, with the Molecular cloud S152 is to the immediate W.		
Optical: Faint optical filaments.		
X-ray: Semicircular shell, with pulsar at W edge.		
Point sources: Long period X-ray pulsar (magnetar).		
Distance: Various observations imply 3.2 kpc.		
References:		
Hughes <i>et al.</i> 1981, ApJ, 246, L127. WSRT at 610 MHz ($1' : S = 40 \pm 5$ Jy) shows bad CLEAN artefacts.	Parmar <i>et al.</i> 1998, A&A, 330, 175. X-ray observations.	
Blair & Kirshner 1981, Nature, 291, 132. Optical spectra.	Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.	
Downes 1983, MNRAS, 203, 695. Effelsberg 100-m at 2.7 GHz ($4.4' : S = 13.0 \pm 1.5$ Jy).	Koralesky <i>et al.</i> 1998, AJ, 116, 1323. VLA search for OH emission.	
Sofue <i>et al.</i> 1983, PASJ, 35, 447. NRO 45-m at 10.2 GHz ($2.7'$).	Patel <i>et al.</i> 2001, ApJ, 563, L45. Chandra observations of pulsar.	
Hughes <i>et al.</i> 1984, ApJ, 283, 147. WSRT at 610 MHz ($20'' : S = 26 \pm 3$ Jy) 1.4 GHz ($10''$), DRAO at 1.4 GHz ($1' \times 1.2' : S = 16.8 \pm 2$ Jy) and Algonquin 46-m at 6.5 GHz ($4.5' : S = 6.7 \pm 1$ Jy).	Kothes <i>et al.</i> 2002, ApJ, 576, 169. CGPS at 1.4 GHz ($68'' \times 59''$), including H α , plus CO observations.	
Braun & Strom 1986, A&AS, 63, 345. WSRT H α observations.	Sasaki <i>et al.</i> 2004, ApJ, 617, 322. XMM-Newton observations.	
Tatematsu <i>et al.</i> 1987, PASJ, 39, 755. NRO 45-m at 10 GHz ($2.7'$), plus polarisation.	Sasaki <i>et al.</i> 2006, ApJ, 642, L149. CO observations of surroundings, plus Chandra observations.	
Tatematsu <i>et al.</i> 1987, A&A, 184, 279. CO observations of the surroundings ($2.7'$).	Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 26 \pm 3$ Jy) and 1420 MHz ($\sim 1' : S = 17.4 \pm 1.2$ Jy), including polarisation and review of flux densities.	
Hanson <i>et al.</i> 1988, A&A, 195, 114. EXOSAT of pulsar. see also: Hanson <i>et al.</i> 1988, A&A, 207, 204. Erratum.	Tian <i>et al.</i> 2010, MNRAS, 404, L1. CGPS for H α absorption.	
Morini <i>et al.</i> 1988, ApJ, 333, 777. EXOSAT observations.	Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 9.8 \pm 0.5$ Jy) including polarisation and review of flux densities.	
Koyama <i>et al.</i> 1989, PASJ, 41, 461. X-ray observations of pulsar.	Kothes & Foster 2012, ApJ, 746, L4. H α and CO observations of region.	
Coe <i>et al.</i> 1989, MNRAS, 238, 649. IRAS observations of surroundings.	Castro <i>et al.</i> 2012, ApJ, 756, 88. Fermi observations.	
Green 1989, MNRAS, 238, 737. OH observations.	Sasaki <i>et al.</i> 2013, A&A, 552, A45. Chandra observations of NE.	
Tatematsu <i>et al.</i> 1990, ApJ, 351, 157. CO of surroundings, plus X-ray observations.	Tendulkar <i>et al.</i> 2013, ApJ, 772, 31. Pulsar proper motion study.	
Davies & Coe 1991, MNRAS, 249, 313. Optical and IR observations near pulsar.	Vogel <i>et al.</i> 2014, ApJ, 789, 75. NuSTAR observations of pulsar.	
Hurford & Fesen 1995, MNRAS, 277, 549. ROSAT imaging.	Nakano <i>et al.</i> 2015, PASJ, 67, 9. Suzaku observations.	
Rho & Petre 1997, ApJ, 484, 828. ROSAT observations.	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.	

G111.7–2.1

RA: 23^h23^m26^s
Dec: +58°48'

1-GHz flux/Jy: 2400
Spectral index: 0.77

Presumably the remnant of a late 17th century SN.

Radio: Bright shell with compact knots and extended plateau of emission; shows secular decline.

Optical: Fast knots and quasi-stationary flocculli, with many filaments at large radii, and NE ‘jet’.

X-ray: Incomplete shell, with hard spectral component.

Point sources: Central compact X-ray source.

Distance: Optical expansion gives 3.3 kpc.

References:

- Anderson & Rudnick 1995, ApJ, 441, 307. VLA at 1.4 GHz and 4.8 GHz, for proper motion studies.
 Kassim *et al.* 1995, ApJ, 455, L59. VLA at 74 (25'') and 332 MHz (6''), including spectral index comparisons.
 Anderson & Rudnick 1996, ApJ, 456, 234. VLA for spectral index comparisons.
 Schwarz *et al.* 1997, A&AS, 123, 43. WSRT at 1.4 GHz (30'') for H_I absorption.
 O’Sullivan & Green 1999, MNRAS, 303, 575. Flux density changes at 13.5, 15.5 and 16.5 GHz.
 Agüeros & Green 1999, MNRAS, 305, 957. 151 MHz observations for bulk expansion studies.
 Gotthelf *et al.* 2001, ApJ, 552, L39. Chandra observations, showing outer shock.
 Hwang *et al.* 2001, ApJ, 560, L175. Chandra observations of Doppler shifted lines.
 Fesen *et al.* 2001, AJ, 122, 2644. HST observations.
 Reynoso & Goss 2002, ApJ, 575, 871. VLA at 5 GHz (6.4'×6.0') for H₂CO absorption studies.
 Morse *et al.* 2004, ApJ, 614, 727. HST proper motion studies.
 Krause *et al.* 2004, Nature, 432, 596. IR and molecular line observations of surroundings.
 Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
 Wilson & Batrla 2005, A&A, 430, 561. Observational constraints on dust.
 Krause *et al.* 2005, Science, 308, 1604. Spitzer light echoes from surroundings.
 Fesen *et al.* 2006, ApJ, 636, 848. Optical and IR limits for central sources.
 Laming *et al.* 2006, ApJ, 644, 260. Chandra observations, particularly of NE ‘jet’.
 Fesen *et al.* 2006, ApJ, 645, 283. HST observations for expansion studies.
 Renaud *et al.* 2006, ApJ, 647, L41. γ -ray line observations.
 Lazendic *et al.* 2006, ApJ, 651, 250. Chandra spectroscopy of bright knots.
 Ennis *et al.* 2006, ApJ, 652, 376. Spitzer observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_I.
 Albert *et al.* 2007, A&A, 474, 937. γ -ray observations.
 Rho *et al.* 2008, ApJ, 673, 271. Spitzer observations.
 Uchiyama & Aharonian 2008, ApJ, 677, L105. X-ray variability of compact features.
 Rest *et al.* 2008, ApJ, 681, L81. SN light echo.
 Hammell & Fesen 2008, ApJS, 179, 195. HST of outer knots.
 Krause *et al.* 2008, Science, 320, 1195. SN light echo spectrum.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Helmboldt & Kassim 2009, AJ, 138, 838. Low radio frequency temporal variations.
 Smith *et al.* 2009, ApJ, 693, 713. Spitzer observations.
 Patnaude & Fesen 2009, ApJ, 697, 535. X-ray proper motions.
 Pavlov & Luna 2009, ApJ, 703, 910. Chandra observations of central source.
 Barlow *et al.* 2010, A&A, 518, L138. Herschel observations.
 Abdo *et al.* 2010, ApJ, 710, L92. Fermi observations.
- Acciari *et al.* 2010, ApJ, 714, 163. γ -ray observations.
 Sibthorpe *et al.* 2010, ApJ, 719, 1553. Akari and sub-mm observations of region.
 DeLaney *et al.* 2010, ApJ, 725, 2038. Spitzer and Chandra observations for 3-D structure.
 Patnaude *et al.* 2011, ApJ, 729, L28. Chandra observations of fading.
 Rest *et al.* 2011, ApJ, 732, 3. SN light echo observations.
 Fesen *et al.* 2011, ApJ, 736, 109. HST variability studies.
 Besel & Krause 2012, A&A, 541, L3. IR light echoes.
 Hwang & Laming 2012, ApJ, 746, 130. Chandra observations.
 Rho *et al.* 2012, ApJ, 747, L6. Akari CO observations.
 Vogt *et al.* 2012, ApJ, 750, 155. Spitzer light echoes.
 Isenmeier *et al.* 2012, ApJ, 757, 126. Spitzer spectroscopy.
 Asgekar *et al.* 2013, A&A, 551, L11. LOFAR of carbon recombination lines.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
 Rutherford *et al.* 2013, ApJ, 769, 64. Chandra spectroscopy.
 Milisavljevic & Fesen 2013, ApJ, 772, 134. Optical spectroscopy for 3-D structure.
 Koo *et al.* 2013, Science, 342, 1346. IR spectroscopy.
 Willström *et al.* 2013, A&A, 558, L2. CO observations.
 DeLaney *et al.* 2014, ApJ, 785, 7. VLA plus Pie Town at 74 MHz (9'') and comparison with higher frequencies.
 Arendt *et al.* 2014, ApJ, 786, 55. Spitzer and Herschel IR observations.
 Patnaude & Fesen 2014, ApJ, 789, 138. Multi-epoch optical and X-ray observations.
 Lee *et al.* 2014, ApJ, 789, 7. Spitzer and CO observations.
 Vinyaikin 2014, ARep, 58, 626. Time evolution of radio emission.
 Alarie *et al.* 2014, MNRAS, 441, 2996. Optical imaging/spectroscopy, including distance from expansion.
 Grefenstette *et al.* 2014, Nature, 506, 339. NuSTAR ⁴⁴Ti observations.
 Siegert *et al.* 2015, A&A, 579, A124. INTEGRAL observations.
 Grefenstette *et al.* 2015, ApJ, 802, 15. NuSTAR observations.
 Lee *et al.* 2015, ApJ, 808, 98. WISE, Spitzer and other IR observations.
 Gorbynov *et al.* 2015, OAP, 28, 224. Time evolution of radio emission.
 Milisavljevic & Fesen 2015, Science, 347, 526. Near-IR observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 7 frequencies between 30 and 353 GHz.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Fesen & Milisavljevic 2016, ApJ, 818, 17. HST [SIII] and [SII] observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Wang & Li 2016, ApJ, 825, 102. INTEGRAL observations.
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

Cassiopeia A, 3C461

Size/arcmin: 5
Type: S

G113.0+0.2

RA: 23^h26^m50^s
Dec: +61°26'

1-GHz flux/Jy: 4
Spectral index: 0.5

Size/arcmin: 40×17?
Type: ?

Radio: Elongated, extent not well defined.

Point sources: Contains old pulsar.

Distance: Association for H_I indicates 3.1 kpc.

References:

Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz (2.8×3.1') and 1.4 GHz (49''×55'') including H_I.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3') and 1420 MHz (~1').

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9.5 : $S = 1.9 \pm 0.5$ Jy) including polarisation and review of flux densities.

G114.3+0.3

RA: 23^h37^m00^s
Dec: +61°55'

1-GHz flux/Jy: 5.5
Spectral index: 0.5

Size/arcmin: 90×55
Type: S

Radio: Shell, with H_{II} region S165 within the boundary of the remnant.

Optical: Faint emission in centre and to S.

Point sources: Pulsar near centre of remnant.

Distance: Association with H_I and other features implies 0.7 kpc.

References:

Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4.4 : $S = 3.6$ Jy) and $S_{1.4 \text{ GHz}} = 4.4$ Jy from 1.4 GHz survey data, plus H_I from Maryland–Green Bank survey.
Kulkarni *et al.* 1993, Nature, 362, 135. Pulsar association.
Fürst *et al.* 1993, A&A, 276, 470. Pulsar association.
Becker *et al.* 1996, A&A, 306, 464. ROSAT of pulsar.
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
Mavromatakis *et al.* 2002, A&A, 383, 1011. Optical observations.

Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz (49''×55''), including H_I (1.0×1.1').
Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz (3.4×3.9 : $S = 12.0 \pm 6.0$ Jy) and 1.4 GHz (1.0×1.1 : $S = 9.8 \pm 0.8$ Jy).
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz (~1' : $S = 5.4 \pm 0.8$ Jy), including review of flux densities.
Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5 : $S = 6.9 \pm 0.7$ Jy), including polarisation and review of flux densities.

G116.5+1.1

RA: 23^h53^m40^s
Dec: +63°15'

1-GHz flux/Jy: 10
Spectral index: 0.5

Size/arcmin: 80×60
Type: S

Radio: Distinct shell, with high polarisation.

Optical: Detected.

Distance: Association with H_I features implies 1.6 kpc.

References:

Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4.4 : $S = 4.7 \pm 0.4$ Jy) and $S_{1.4 \text{ GHz}} = 8.0 \pm 0.8$ Jy from 1.4 GHz survey data, plus H_I from Maryland–Green Bank survey.
Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz (49''×55''), including H_I (1.0×1.1').
Mavromatakis *et al.* 2005, A&A, 435, 141. Optical observations.

Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz (3.4×3.8 : $S = 15.0 \pm 1.5$ Jy) and 1.4 GHz (1.0×1.1 : $S = 10.6 \pm 0$ Jy).
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : $S = 12.5 \pm 1.6$ Jy) and 1420 MHz (~1' : $S = 10.3 \pm 0.70$ Jy), including polarisation and review of flux densities.
Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5 : $S = 5.7 \pm 0.6$ Jy), including polarisation and review of flux densities.

G116.9+0.2		CTB 1
RA: 23 ^h 59 ^m 10 ^s	1-GHz flux/Jy: 8	Size/arcmin: 34
Dec: +62°26'	Spectral index: 0.57	Type: S
Has been called G117.3+0.1 and G116.9+0.1.		
Radio: Incomplete shell.		
Optical: Filaments on sky survey.		
X-ray: Centrally brightened, with NE ‘breakout’.		
Point sources: Pulsar to NE.		
Distance: Association with H _I features implies 1.6 kpc.		
References:		
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 3.9 \pm 1.0$ Jy), and 37-m at 1.7 GHz ($S = 5.5 \pm 2.0$ Jy), plus review of flux densities.		sar.
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical observations.		Fesen <i>et al.</i> 1997, AJ, 113, 767. Optical observations.
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 4.2 \pm 0.2$ Jy).		Craig <i>et al.</i> 1997, ApJ, 488, 307. ROSAT observations.
Angerhofer <i>et al.</i> 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6.8' : S = 3.0 \pm 0.3$ Jy).		Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.
Dickel & Willis 1980, A&A, 85, 55. WSRT at 610 MHz ($56'' \times 64''$) and 1.4 GHz ($22'' \times 25''$).		Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz ($69''$), including polarisation.
Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz ($4.4' : S = 4.8 \pm 0.4$ Jy) and $S_{1.4 \text{ GHz}} = 7.8 \pm 0.8$ Jy from 1.4 GHz survey data, plus H _I from Maryland–Green Bank survey.		Yar-Uyaniker <i>et al.</i> 2004, ApJ, 616, 247. CGPS at 1.4 GHz ($49'' \times 55''$), including H _I ($1.0' \times 1.1'$).
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.		Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
Landecker <i>et al.</i> 1982, AJ, 87, 1379. DRAO at 1.42 GHz and H _I ($2' \times 2.3' : S = 8.3 \pm 0.5$ Jy), plus review of flux densities.		Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz ($3.4' \times 3.8' : S = 15.0 \pm 1.5$ Jy) and 1.4 GHz ($1.0' \times 1.1' : S = 8.1 \pm 0.4$ Jy).
see also: Landecker <i>et al.</i> 1983, AJ, 88, 877. Erratum.		Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.5 \pm 0.8$ Jy) and 1420 MHz ($\sim 1' : S = 7.0 \pm 0.8$ Jy), including review of flux densities.
Fesen <i>et al.</i> 1985, ApJ, 292, 29. Optical spectra.		Pannuti <i>et al.</i> 2010, AJ, 140, 1787. ASCA and Chandra observations.
Hailey & Craig 1994, ApJ, 434, 635. Optical spectroscopy.		Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 3.6 \pm 0.4$ Jy) including polarisation and review of flux densities.
Hailey & Craig 1995, ApJ, 455, L151. ROSAT of nearby pul-		

G119.5+10.2		CTA 1
RA: 00 ^h 06 ^m 40 ^s	1-GHz flux/Jy: 36	Size/arcmin: 90?
Dec: +72°45'	Spectral index: 0.6	Type: S
Has been called G119.5+10.3.		
Radio: Incomplete shell, with ‘breakout’ to NW.		
Optical: Faint diffuse nebulosities.		
X-ray: Centrally brightened.		
Point sources: Central pulsar.		
Distance: Associated H _I shell indicates 1.4 kpc.		
References:		
Sieber <i>et al.</i> 1979, A&A, 74, 361. Effelsberg 100-m at 2.7 GHz (4'': $S=23.6\pm2.8$ Jy).		Reich 2002, in NSPS, p1. Effelsberg 100-m at 1.4 GHz (9'').
Sieber <i>et al.</i> 1981, A&A, 103, 393. Effelsberg 100-m at 2.7 GHz and 1.7 GHz (7'': $S=31.6\pm2.5$ Jy), and 151 MHz (4'': $S=62.6\pm6$ Jy).		Slane <i>et al.</i> 2004, ApJ, 601, 1045. XMM-Newton observations of central source.
Fesen <i>et al.</i> 1981, ApJ, 247, 148. Optical, including spectra.		Halpern <i>et al.</i> 2004, ApJ, 612, 398. Chandra observations of central nebula, plus optical and radio limits for compact source.
Fesen <i>et al.</i> 1983, ApJS, 51, 337. Deep [OIII] imagery.		Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
Seward 1990, ApJS, 73, 781. Einstein observations.		Abdo <i>et al.</i> 2008, Science, 322, 1218. Fermi detection of pulsar.
Simonetti 1992, ApJ, 386, 170. VLA observations of background sources for rotation measure studies.		Lin <i>et al.</i> 2010, ApJ, 725, L1. XMM-Newton observations of pulsar.
Pineault <i>et al.</i> 1993, AJ, 105, 1060. DRAO at 1.4 GHz (1'': $S=34\pm4$ Jy), plus H _I and IRAS.		Caraveo <i>et al.</i> 2010, ApJ, 725, L6. XMM-Newton observations of pulsar.
Seward <i>et al.</i> 1995, ApJ, 453, 284. ROSAT observations.		Sun <i>et al.</i> 2011, A&A, 535, A64. Urumqi 25-m at 4.8 GHz (9'': $S=11.6\pm1.2$ Jy) and Effelsberg 100-m at 2.6 GHz (4'': $S=20.3\pm2.0$ Jy) including polarisation.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.		Lin <i>et al.</i> 2012, MNRAS, 426, 2283. Suzaku observations.
Slane <i>et al.</i> 1997, ApJ, 485, 221. ASCA and ROSAT observations.		Mignani <i>et al.</i> 2013, MNRAS, 430, 1354. Optical limits for pulsar.
Pineault <i>et al.</i> 1997, A&A, 324, 1152. DRAO at 408 MHz (3'': and 1.4 GHz (1'')).		Aliu <i>et al.</i> 2013, ApJ, 764, 38. γ -ray observations.
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		Li <i>et al.</i> 2016, ApJ, 831, 19. γ -ray observations of pulsar.
Brazier <i>et al.</i> 1998, MNRAS, 295, 819. Studies of central, compact γ -ray source.		
Mavromatakis <i>et al.</i> 2000, A&A, 353, 371. Optical imagery.		
Roberts <i>et al.</i> 2001, ApJS, 133, 451. ASCA observations.		

G120.1+1.4		Tycho, 3C10, SN1572			
RA: 00 ^h 25 ^m 18 ^s	1-GHz flux/Jy: 56	Size/arcmin: 8	Type: S		
Dec: +64°09'	Spectral index: 0.58				
This is the remnant of the Tycho's SN of AD1572.					
Radio: Shell, brightest to the NE.					
Optical: Faint filaments/knots to the NNW, NE and E.					
X-ray: Shell, brighter to the NE.					
Point sources: Faint radio source near centre of the remnant, thought to be extragalactic.					
Distance: H _I observations suggest 2.3–3 kpc, optical proper motion and shock velocity gives 2.4 kpc.					
References:					
Duin & Strom 1975, A&A, 39, 33. WSRT at 610 MHz ($57'' \times 64''$) and 5 GHz ($7'' \times 8''$).					
Klein <i>et al.</i> 1979, A&A, 76, 120. Effelsberg 100-m at 10.7 GHz ($1''.2 : S = 13.1 \pm 0.8$ Jy), plus review of flux densities.					
Strom <i>et al.</i> 1982, MNRAS, 200, 473. WSRT at 1415 MHz ($27'' \times 31''$) from 1971 and 1979, for expansion.					
Dicke <i>et al.</i> 1982, ApJ, 257, 145. Comparison of radio, X-ray and optical observations.					
Reid <i>et al.</i> 1982, ApJ, 261, 485. Einstein observations.					
Seward <i>et al.</i> 1983, ApJ, 266, 287. Einstein observations.					
Tan & Gull 1985, MNRAS, 216, 949. Radio at 2.7 GHz ($4''$) from 1980 and 1983, and One-Mile Telescope at 1.4 GHz ($23''$) from 1965 and 1980, for expansion.					
Albinson <i>et al.</i> 1986, MNRAS, 219, 427. H _I observations.					
Green & Gull 1987, MNRAS, 224, 1055. VLA H _I absorption observations towards central radio source.					
Smith <i>et al.</i> 1988, ApJ, 325, 288. EXOSAT spectrum and image (smoothed to $13''$).					
Dicke <i>et al.</i> 1991, AJ, 101, 2151. VLA at 1.4 and 5 GHz ($1''.5$).					
Wood <i>et al.</i> 1992, AJ, 103, 1338. VLA at 5 GHz ($1''.5$) polarisation studies.					
Fink <i>et al.</i> 1994, A&A, 283, 635. X-ray spectra.					
Vancura <i>et al.</i> 1995, ApJ, 441, 680. X-ray spectra and ROSAT image.					
Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.					
Schwarz <i>et al.</i> 1995, A&A, 299, 193. WSRT and Effelsberg 100-m at 1.4 GHz ($50''$), and VLA at 1.4 GHz ($13''$), for neutral hydrogen studies.					
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.					
Hwang & Gotthelf 1997, ApJ, 475, 665. ASCA observations.					
Reynoso <i>et al.</i> 1997, ApJ, 491, 816. VLA at 1.4 GHz ($1''.4 \times 1''.5$) from 1984 and 1994 for expansion studies.					
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.					
Reynoso <i>et al.</i> 1999, AJ, 117, 1827. VLA at 1.4 GHz ($13''$), for H _I studies.					
Katz-Stone <i>et al.</i> 2000, ApJ, 529, 453. VLA at 333 MHz and 1.4 GHz ($9''.5 \times 7''.7$), for spectral index studies.					
Ghavamian <i>et al.</i> 2000, ApJ, 535, 266. Optical observations of shock precursors.					
Hughes 2000, ApJ, 545, L53. ROSAT X-ray expansion.					
Decourchelle <i>et al.</i> 2001, A&A, 365, L218. XMM-Newton observations.					
Ghavamian <i>et al.</i> 2001, ApJ, 547, 995. Optical spectroscopy.					
Douville <i>et al.</i> 2001, A&A, 373, 281. ISO observations.					
Aharonian <i>et al.</i> 2001, A&A, 373, 292. Upper limit on high energy γ -rays.					
Hwang <i>et al.</i> 2002, ApJ, 581, 1101. Chandra observations.					
Lee <i>et al.</i> 2004, ApJ, 605, L113. Observations of molecular clouds in vicinity.					
Bamba <i>et al.</i> 2005, ApJ, 621, 793. Chandra observations of rim.					
Warren <i>et al.</i> 2005, ApJ, 634, 376. Chandra observations.					
Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 86 \pm 5$ Jy) and 1420 MHz ($\sim 1' : S = 40.5 \pm 1.5$ Jy), including polarisation and review of flux densities.					
Lee <i>et al.</i> 2007, ApJ, 659, L133. H α observations.					
Cassam-Chenaï <i>et al.</i> 2007, ApJ, 665, 315. Chandra observations.					
Rest <i>et al.</i> 2008, ApJ, 681, L81. SN light echo.					
Krause <i>et al.</i> 2008, Nature, 456, 617. SN light echo spectrum.					
Hurley-Walker <i>et al.</i> 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.					
Katsuda <i>et al.</i> 2010, ApJ, 709, 1387. Chandra proper motions study.					
Raymond <i>et al.</i> 2010, ApJ, 712, 901. H α spectroscopy.					
Lee <i>et al.</i> 2010, ApJ, 715, L146. H α observations.					
Hayato <i>et al.</i> 2010, ApJ, 725, 894. Suzaku observations of expansion.					
Ishihara <i>et al.</i> 2010, A&A, 521, L61. Akari observations.					
Tian & Leahy 2011, ApJ, 729, L15. H _I and CO observations.					
Acciari <i>et al.</i> 2011, ApJ, 730, L20. γ -ray observations.					
Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9''.5 : S = 20.0 \pm 2.0$ Jy) including polarisation and review of flux densities.					
Giordano <i>et al.</i> 2012, ApJ, 744, L2. Fermi observations.					
Gomez <i>et al.</i> 2012, MNRAS, 420, 3557. Herschel IR dust observations.					
Yang <i>et al.</i> 2013, ApJ, 766, 44. Suzaku spectroscopy.					
Williams <i>et al.</i> 2013, ApJ, 770, 129. Spitzer observations.					
Yamaguchi <i>et al.</i> 2014, ApJ, 780, 136. Suzaku observations.					
Wang & Li 2014, ApJ, 789, 123. INTEGRAL observations.					
Troja <i>et al.</i> 2014, ApJ, 797, L6. Swift observations.					
Miceli <i>et al.</i> 2015, ApJ, 805, 120. XMM-Newton observations.					
Lu <i>et al.</i> 2015, ApJ, 805, 142. Chandra observations.					
Katsuda <i>et al.</i> 2015, ApJ, 808, 49. Suzaku observations.					
Tran <i>et al.</i> 2015, ApJ, 812, 101. Chandra observations.					
Lopez <i>et al.</i> 2015, ApJ, 814, 132. NuSTAR observations.					
Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux densities at 5 frequencies between 30 and 143 GHz.					
Koo <i>et al.</i> 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).					
Williams <i>et al.</i> 2016, ApJ, 823, L32. Multi-epoch X-ray and radio observations for expansion.					
Zhou <i>et al.</i> 2016, ApJ, 826, 34. CO observations of region.					

G126.2+1.6

RA: 01^h22^m00^s
Dec: +64°15'

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 70
Type: S?

Radio: Poorly defined shell.

Optical: Filaments, mostly in W.

References:

- Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz (9' : $S = 6.8 \pm 0.7$ Jy) and 2.7 GHz (4'4 : $S = 3.9 \pm 0.4$ Jy).
 Blair *et al.* 1980, ApJ, 242, 592. Optical detection and spectra.
 Rosado 1982, RMxAA, 5, 127. Optical spectra.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz (4'4) and 4.8 GHz (2'6).
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz (3'5 × 3'9 : $S = 12 \pm 2.5$ Jy) and part at 1.4 GHz (1'1 × 1'0), plus review of flux densities.
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5 : $S = 6.1 \pm 1.6$ Jy).
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
 Tian & Leahy 2006, A&A, 447, 205. CGPS at 408 MHz (3'4 × 3'8 : $S = 9.7 \pm 3.9$ Jy) and 1.4 GHz (1'0 × 1'1 : $S = 6.7 \pm 2.1$ Jy), plus other observations for spectral index studies.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : $S = 5.7 \pm 0.7$ Jy) and 1420 MHz (~1' : $S = 6.4 \pm 1.1$ Jy), including review of flux densities.
 Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz (9'5 : $S = 2.6 \pm 0.6$ Jy), including polarisation.
 see also: Sun *et al.* 2007, A&A, 469, 1003. Erratum.

G127.1+0.5

RA: 01^h28^m20^s
Dec: +63°10'

1-GHz flux/Jy: 12
Spectral index: 0.45

Size/arcmin: 45
Type: S

R5

Has been called G127.3+0.7.

Radio: Distinct shell, with bright central source.

Optical: Detected.

Point sources: Flat radio spectrum (extragalactic) source at centre of remnant.

Distance: 1.2–1.3 kpc if associated with NGC 559.

References:

- Caswell 1977, MNRAS, 181, 789. Half-Mile Telescope at 1.42 GHz (3'5 × 3'9), plus other observations of central source.
 Pauls 1977, A&A, 59, L13. Effelsberg 100-m at 1.4 GHz (9' : $S = 8 \pm 1$ Jy).
 Salter *et al.* 1978, A&A, 66, 77. Effelsberg 100-m at 2.7 GHz (4'4), plus 5 and 8.7 GHz of the central source.
 Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz (9' : $S = 10.8 \pm 1.3$ Jy).
 Pauls *et al.* 1982, A&A, 112, 120. WSRT at 610 MHz (56'' × 62'') and HI absorption to the point source with the VLA.
 Geldzahler & Shaffer 1982, ApJ, 260, L69. Observations of central source.
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz (4'4) and 4.8 GHz (2'6).
 Goss & van Gorkom 1984, JApA, 5, 425. WSRT HI absorption of central source.
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz (3'5 × 3'9 : $S = 17.9 \pm 2.0$ Jy) and 1.4 GHz (1'13 × 1'0 : $S = 10.1 \pm 0.8$ Jy), plus review of flux densities.
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5 : $S = 14.6 \pm 0.8$ Jy).
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact sources.
 Leahy & Tian 2006, A&A, 451, 251. CGPS at 408 MHz (3'4 × 3'8 : $S = 17.1 \pm 1.7$ Jy) and 1.4 GHz (1'0 × 1'2 : $S = 10.0 \pm 0.8$ Jy).
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : $S = 15.9 \pm 1.0$ Jy) and 1420 MHz (~1' : $S = 9.7 \pm 0.6$ Jy), including polarisation and review of flux densities.
 Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz (9'5 : $S = 6.3 \pm 0.7$ Jy), including polarisation.
 see also: Sun *et al.* 2007, A&A, 469, 1003. Erratum.
 Zhou *et al.* 2014, ApJ, 791, 109. CO observations.

G130.7+3.1

RA: 02^h05^m41^s
Dec: +64°49'

1-GHz flux/Jy: 33
Spectral index: 0.07

This is the remnant of the SN of AD1181.

Radio: Filled-centre, highly polarised, with high frequency turnover.

Optical: Faint filaments.

X-ray: Centrally brightened, with faint jet.

Point sources: Central pulsar.

Distance: H_I absorption indicates 2 kpc.

References:

- Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz ($58'' : S = 26.7 \pm 0.5$ Jy).
 Wilson & Weiler 1976, A&A, 49, 357. WSRT at 610 MHz ($58'' \times 64''$), 1.4 GHz ($24'' \times 27''$) and 5 GHz ($7'' \times 8''$).
 van den Bergh 1978, ApJ, 220, L9. Optical observations.
 Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.
 Green & Gull 1982, Nature, 299, 606. H_I absorption distance.
 Fesen 1983, ApJ, 270, L53. Optical spectra.
 Reynolds & Aller 1985, AJ, 90, 2312. VLA at 1.4 GHz (2'), for limits of shell.
 Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT spectrum.
 Green 1986, MNRAS, 218, 533. 151 MHz observations ($1.2 \times 1.3' : S = 36 \pm 4$ Jy), plus 2.7 GHz (4''), plus Einstein observations for limit on shell.
 Green 1987, MNRAS, 225, 11P. Flux density increase at 408 MHz.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz ($26'' : S = 24.2 \pm 1.4$ Jy).
 Reynolds & Aller 1988, ApJ, 327, 845. VLA at 1.4 (2.4') and 4.9 GHz (2.5').
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ($90'' : S = 15.0 \pm 2.0$ Jy), plus review of flux densities.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Green & Scheuer 1992, MNRAS, 258, 833. IRAS upper limits.
 Roberts *et al.* 1993, A&A, 274, 427. H_I absorption.
 Wallace *et al.* 1994, A&A, 286, 565. H_I of surroundings.
 Helfand *et al.* 1995, ApJ, 453, 741. ROSAT observations.
 Torii *et al.* 2000, PASJ, 52, 875. ASCA observations.
 Bietenholz *et al.* 2001, ApJ, 560, 772. VLA at 74 MHz ($26'' : S = 33.6$ Jy) and 327 MHz ($8''.2 : S = 33.9$ Jy), for spectral index and expansion studies.
 Bocchino *et al.* 2001, A&A, 369, 1078. XMM-Newton observations.
 Murray *et al.* 2002, ApJ, 568, 226. Chandra pulsar detection.
 Camilo *et al.* 2002, ApJ, 571, L41. Pulsar detection in radio.
- Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz ($26''$) for polarised intensity.
 Slane *et al.* 2002, ApJ, 571, L45. Chandra observations.
 Slane *et al.* 2004, ApJ, 616, 403. Deep Chandra imaging.
 Bietenholz 2006, ApJ, 645, 1180. VLA at 1.4 GHz ($1''.36$) for expansion studies.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 32.2 \pm 2.0$ Jy) and 1420 MHz ($\sim 1' : S = 31.9 \pm 1.0$ Jy), including polarisation and review of flux densities.
 Gotthelf *et al.* 2007, ApJ, 654, 267. XMM-Newton observations.
 Slane *et al.* 2008, ApJ, 676, L33. Spitzer and other IR observations.
 Fesen *et al.* 2008, ApJS, 174, 379. Optical observations for proper motion studies.
 Shearer & Neustroev 2008, MNRAS, 390, 235. Optical observations of pulsar nebula.
 Shibanov *et al.* 2008, A&A, 486, 273. Optical observations of pulsar nebula.
 Abdo *et al.* 2009, ApJ, 699, L102. Fermi observations of pulsar.
 Livingstone *et al.* 2009, ApJ, 706, 1163. Pulsar observations.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9''.5 : S = 31.7 \pm 3.0$ Jy) including polarisation and review of flux densities.
 Bietenholz *et al.* 2013, MNRAS, 431, 2590. Proper motion study of pulsar.
 Kothes *et al.* 2013, A&A, 560, A18. CGPS H_I observations for distance.
 Aleksić *et al.* 2014, A&A, 567, L8. γ-ray detection.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 7 frequencies between 30 and 353 GHz.

3C58, SN1181

Size/arcmin: 9×5
Type: F

G132.7+1.3		HB3
RA: 02 ^h 17 ^m 40 ^s	1-GHz flux/Jy: 45	Size/arcmin: 80
Dec: +62°45'	Spectral index: 0.6	Type: S
Has been called G132.4+2.2.		
Radio: Faint shell, adjacent to W3/4/5 complex.		
Optical: Complete, filamentary shell, shock excited spectra.		
X-ray: Partial shell.		
Point sources: Pulsar nearby.		
Distance: Interaction with surroundings gives 2 kpc.		
References:		
van den Berg <i>et al.</i> 1973, ApJS, 26, 19. Optical observations.		
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5':S=33.8\pm7.0$ Jy).		
d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.		
Read 1981, MNRAS, 194, 863. Radio at 151 MHz ($4.4'$) and 1.4 GHz ($2'$) showing HI shell.		
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.		
Fesen & Gull 1983, PASP, 95, 196. Optical image.		
Leahy <i>et al.</i> 1985, ApJ, 294, 183. Einstein observations.		
Landecker <i>et al.</i> 1987, AJ, 94, 111. DRAO at 408 MHz ($3.5'\times4':S=75\pm15$ Jy), plus review of flux densities.		
Routledge <i>et al.</i> 1991, A&A, 247, 529. DRAO at 1.4 GHz ($1.0'\times1.1'$) for HI, plus CO observations.		
Fesen <i>et al.</i> 1995, AJ, 110, 2876. Optical imaging and spectroscopy, DRAO at 408 MHz ($3.5'\times4'$) and 1.4 GHz ($1.0'\times1.1'$).		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
Koralesky <i>et al.</i> 1998, AJ, 116, 1323. VLA detection of compact OH emission.		
Reich <i>et al.</i> 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14.5':S=51.5\pm3.5$ Jy).		
Tian & Leahy 2005, A&A, 436, 187. CGPS at 408 MHz ($3.4'\times3.8':S=29.4\pm2.7$ Jy), including review of flux densities.		
see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.		
Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim3':S=61\pm9$ Jy) and 1420 MHz ($\sim1':S=29.4\pm2.7$ Jy), including review of flux densities.		
Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.		
Green 2007, BASI, 35, 77. Review of radio spectrum.		
Shi <i>et al.</i> 2008, A&A, 487, 601. Urumqi 25-m at 4.8 GHz ($9.5':S=2.5\pm0.5$ Jy), plus other survey observations for spectral studies.		
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.		
Kilpatrick <i>et al.</i> 2016, ApJ, 816, 1. CO observations, including broad lines.		
Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.		
Katagiri <i>et al.</i> 2016, ApJ, 818, 114. γ -ray observations.		
Zhou <i>et al.</i> 2016, ApJ, 833, 4. CO observations of region.		
G152.4–2.1		
RA: 04 ^h 07 ^m 50 ^s	1-GHz flux/Jy: 3.5?	Size/arcmin: 100×95
Dec: +49°11'	Spectral index: 0.7?	Type: S
Radio: Bilateral shell.		
References:		
Foster <i>et al.</i> 2013, A&A, 549, A107. Effelsberg 100-m at 2.7 GHz, including polarisation, plus various radio survey observations.		
G156.2+5.7		
RA: 04 ^h 58 ^m 40 ^s	1-GHz flux/Jy: 5	Size/arcmin: 110
Dec: +51°50'	Spectral index: 0.5	Type: S
Radio: Faint shell, brighter in E and W.		
Optical: Filamentary ring and smaller patchy ring.		
X-ray: Faint shell.		
Distance: Optical/X-ray observations imply >1.7 kpc.		
References:		
Pfeffermann <i>et al.</i> 1991, A&A, 246, L28. ROSAT detection.		
Reich <i>et al.</i> 1992, A&A, 256, 214. Effelsberg 100-m at 1.4 ($9':S=4.2\pm1.0$ Jy) and 2.7 GHz ($4.3':S=3.0\pm1.0$ Jy), plus HI and IRAS.		
Yamauchi <i>et al.</i> 1993, PASJ, 45, 795. Hard X-ray observations.		
Lorimer <i>et al.</i> 1998, A&A, 331, 1002. Pulsar search.		
Yamauchi <i>et al.</i> 1999, PASJ, 51, 13. ASCA observations of some regions.		
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz, including polarisation.		
Pannuti & Allen 2004, AdSpR, 33, 434. ASCA and RXTE observations.		
Kaplan <i>et al.</i> 2006, ApJS, 163, 344. X-ray upper limit on compact sources.		
Kothes <i>et al.</i> 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim3':S=8.1\pm1.3$ Jy), including review of flux densities.		
Xu <i>et al.</i> 2007, A&A, 470, 969. Urumqi 25-m at 4.8 GHz ($9.5':S=2.5\pm0.5$ Jy), including polarisation.		
Gerardy & Fesen 2007, MNRAS, 376, 929. Optical observations.		
Katsuda <i>et al.</i> 2009, PASJ, 61, S155. Suzaku observations.		
Uchida <i>et al.</i> 2012, PASJ, 64, 61. Suzaku observations.		
Katsuda <i>et al.</i> 2016, ApJ, 826, 108. Multi-epoch H α observations for expansion.		

G159.6+7.3

RA: 05^h20^m00^s
Dec: +50°00'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 240×180?
Type: S

Optical: Large, faint shell.

X-ray: Possible emission.

References:

Fesen & Milisavljevic 2010, AJ, 140, 1163. H α and other observations.

G160.9+2.6

RA: 05^h01^m00^s
Dec: +46°40'

1-GHz flux/Jy: 110
Spectral index: 0.64

HB9

Size/arcmin: 140×120
Type: S

Has been called G160.5+2.8 and G160.4+2.8.

Radio: Large, filamentary shell.

Optical: Incomplete shell.

X-ray: Centrally brightened.

Point sources: Pulsar within boundary of the remnant, plus several nearby compact radio sources.

Distance: Various observations suggests less than 4 kpc.

References:

- d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.
Damashek *et al.* 1978, ApJ, 225, L31. Pulsar.
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
Dwarakanath *et al.* 1982, JApA, 3, 207. Radio observations at 34.5 MHz (26'×40' : $S = 750 \pm 150$ Jy), plus review of flux densities.
van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT HI absorption to nearby point source.
Sequaist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
Leahy 1987, ApJ, 322, 917. Einstein observations.
Leahy & Roger 1991, AJ, 101, 1033. DRAO at 408 MHz (3.5'×4.8') and 1.4 GHz (1.0'×1.4'), including HI and discussion of distance.
Yamauchi & Koyama 1993, PASJ, 45, 545. Hard X-ray observations.
Leahy & Aschenbach 1995, A&A, 293, 853. ROSAT observations.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Leahy *et al.* 1998, A&A, 339, 601. 232 MHz (3.8'×5.2'), 151 MHz (4.2'×5.8') and Effelsberg 100-m at 4.7 GHz (2.5') for spectral index studies.
Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density ($S = 1130 \pm 340$ Jy).
Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14.5' : $S = 91 \pm 3$ Jy).
Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : $S = 102 \pm 7$ Jy) and 1420 MHz (~1' : $S = 54.0 \pm 2.9$ Jy), including polarisation and review of flux densities.
Leahy & Tian 2007, A&A, 461, 1013. CGPS at 408 MHz (2.8'×3.9' : $S = 117.8 \pm 5.3$ Jy) and 1.4 GHz (58''×80'' : $S = 65.9 \pm 3.4$ Jy).
Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5' : $S = 34 \pm 3$ Jy), including polarisation and review of flux densities.
Gosachinskii 2013, AstL, 39, 179. HI observations of region.
Araya *et al.* 2014, MNRAS, 444, 860. γ -ray detection.

G166.0+4.3

RA: 05^h26^m30^s
Dec: +42°56'

1-GHz flux/Jy: 7
Spectral index: 0.37

VRO 42.05.01

Size/arcmin: 55×35
Type: S

Radio: Two arcs of strikingly different radii.

Optical: Nearly complete ring.

X-ray: Predominantly in SW.

Distance: H_I indicates 4.5 kpc.

References:

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 5.2 \pm 1.0$ Jy).
 Lozinskaya 1979, AuJPh, 32, 113. H α interferometry.
 Landecker *et al.* 1982, ApJ, 261, L41. DRAO at 1.4 GHz ($1.0' \times 1.4'$), plus review of flux densities.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Pineault *et al.* 1985, A&A, 151, 52. VLA at 1.4 GHz ($16'' \times 20''$) of part of remnant, and optical observations.
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Braun & Strom 1986, A&AS, 63, 345. WSRT H_I Observations.
 Pineault *et al.* 1987, ApJ, 315, 580. DRAO and VLA combined at 1.4 GHz ($20''$).
 Landecker *et al.* 1989, MNRAS, 237, 277. DRAO at 1.4 GHz ($1.0' \times 1.4'$), including H_I.
 Burrows & Guo 1994, ApJ, 421, L19. ROSAT images and spec-

tra.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Guo & Burrows 1997, ApJ, 480, L51. ASCA observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Leahy & Tian 2005, A&A, 440, 929. CGPS at 408 MHz ($3.4' \times 5.0'$) and 1.4 GHz ($1.0' \times 1.4'$), for spectral index studies.
 see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 8.1 \pm 0.9$ Jy) and 1420 MHz ($\sim 1' : S = 5.1 \pm 0.4$ Jy), including polarisation and review of flux densities.
 Bocchino *et al.* 2009, A&A, 498, 139. XMM-Newton observations.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9.5' : S = 3.3 \pm 0.3$ Jy), including polarisation and review of flux densities.
 Araya 2013, MNRAS, 434, 2202. Fermi observations.

G178.2–4.2

RA: 05^h25^m05^s
Dec: +28°11'

1-GHz flux/Jy: 2
Spectral index: 0.5

Size/arcmin: 72×62
Type: S

Radio: Faint shell, brighter in NE.

References:

Gao *et al.* 2011, A&A, 532, A144. Urumqi 25-m at 5 GHz ($9.5' : S = 1.0 \pm 0.1$ Jy), plus other observations.

G179.0+2.6

RA: 05^h53^m40^s
Dec: +31°05'

1-GHz flux/Jy: 7
Spectral index: 0.4

Size/arcmin: 70
Type: S?

Radio: Thick shell, with background extragalactic sources near centre.

References:

Fürst & Reich 1986, A&A, 154, 303. Effelsberg 100-m at 1.4 ($9.4'$), 2.7 ($4.3'$) and 4.75 GHz ($2.4'$).
 Fürst *et al.* 1989, A&A, 223, 66. Observations of central, extra-galactic source.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9.5' : S = 3.2 \pm 0.3$ Jy), including polarisation and review of flux densities.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G180.0–1.7		S147
RA: 05 ^h 39 ^m 00 ^s	1-GHz flux/Jy: 65	Size/arcmin: 180
Dec: +27°50'	Spectral index: varies	Type: S
Radio: Large faint shell, with spectral break.		
Optical: Wispy ring.		
X-ray: Possible detection.		
Point sources: Pulsar within boundary, with faint wind nebula.		
Distance: Optical absorption towards stars indicates > 0.36 and < 0.88 kpc.		
References:		
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical observations.	863 MHz (14'5 : $S = 77 \pm 10$ Jy).	
Sofue <i>et al.</i> 1980, PASJ, 32, 1. Effelsberg 100-m at 5 GHz (2'6 of parts).	Romani & Ng 2003, ApJ, 585, L41. Chandra of pulsar.	
Kundu <i>et al.</i> 1980, A&A, 92, 225. Effelsberg 100-m at 2.7 GHz (5'5 : $S = 34.9 \pm 4$ Jy) and 1.6 GHz (10' : $S = 60.2 \pm 6$ Jy).	Kramer <i>et al.</i> 2003, ApJ, 593, L31. Pulsar observations.	
Angerhofer & Kundu 1981, AJ, 86, 1003. Arecibo at 430 MHz (9' : $S = 97 \pm 20$ Jy).	Sallmen & Welsh 2004, A&A, 426, 555. Optical absorption towards background stars.	
Fürst <i>et al.</i> 1982, A&A, 115, 428. Observations of compact radio sources near the remnant.	Drew <i>et al.</i> 2005, MNRAS, 362, 753. H α imaging.	
Fesen <i>et al.</i> 1985, ApJ, 292, 29. Optical spectra.	Ng <i>et al.</i> 2007, ApJ, 654, 487. Pulsar and wind nebula observations.	
Fürst & Reich 1986, A&A, 163, 185. Effelsberg 100-m at 1.4, 2.7 and 4.7/5.0 GHz (9'4, 4'3 and 2'4/2'6).	Xiao <i>et al.</i> 2008, A&A, 482, 783. Urumqi 25-m at 4.8 GHz (9'5 : $S = 15.4 \pm 3.0$ Jy) and Effelsberg 100-m at 2.6 GHz (4'4 : $S = 34.6 \pm 4.0$ Jy).	
Sauvageot <i>et al.</i> 1990, A&A, 227, 183. EXOSAT possible detection.	Jeong <i>et al.</i> 2012, Ap&SS, 342, 389. CO observations of region.	
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.	Katsuta <i>et al.</i> 2012, ApJ, 752, 135. Fermi observations.	
Anderson <i>et al.</i> 1996, ApJ, 468, L55. Pulsar detection.	Dinçel <i>et al.</i> 2015, MNRAS, 448, 3196. Identification of OB runaway star near centre.	
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.	
Reich <i>et al.</i> 2003, A&A, 408, 961. Effelsberg 100-m at		

G182.4+4.3		
RA: 06 ^h 08 ^m 10 ^s	1-GHz flux/Jy: 0.5	Size/arcmin: 50
Dec: +29°00'	Spectral index: 0.4	Type: S
Radio: Incomplete shell.		
Optical: Brighter in S and NW.		
X-ray: Diffuse emission.		
References:		
Kothes <i>et al.</i> 1998, A&A, 331, 661. Effelsberg 100-m at 1.4, 2.7, 4.9 and 10.5 GHz (9'4 : $S = 0.36 \pm 0.08$ Jy, 4'4 : $S = 0.25 \pm 0.04$ Jy, 2'5 : $S = 0.20 \pm 0.02$ Jy and 1'2 : $S = 0.15 \pm 0.03$ Jy), plus X-ray upper limit.	$S = 0.26 \pm 0.5$ Jy) including polarisation and review of flux densities.	
Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz and 4.9 GHz (3').	Sezer <i>et al.</i> 2012, MNRAS, 427, 1168. Optical and XMM-Newton observations.	
Sun <i>et al.</i> 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 :	Jeong <i>et al.</i> 2012, Ap&SS, 342, 389. CO observations of region.	

G184.6–5.8		Crab Nebula, 3C144, SN1054
RA: 05 ^h 34 ^m 31 ^s	1-GHz flux/Jy: 960	Size/arcmin: 7×5
Dec: +22°01'		
Spectral index: 0.30		
This is the remnant of the SN of AD1054.		
Radio: Filled-centre, central pulsar, with faint ‘jet’ (or tube) extending from the N edge; shows secular decline.		
Optical: Strongly polarised filaments, diffuse synchrotron emission, with ‘jet’ faintly visible.		
X-ray: Central ‘torus’ around the pulsar.		
Point sources: Pulsar powering the remnant.		
Distance: Proper motions and radial velocities give 2 kpc.		
References:		
Velusamy 1984, Nature, 308, 251. VLA at 1.4 GHz, radio detection of ‘jet’.		
Velusamy 1985, MNRAS, 212, 359. VLA at 1.4 GHz (15'').		
Aller & Reynolds 1985, ApJ, 293, L73. Radio flux density decrease.		
Fesen & Gull 1986, ApJ, 306, 259. Deep [OIII] imagery of ‘jet’.		
Marcelin <i>et al.</i> 1990, A&A, 228, 471. Optical of ‘jet’.		
Bietenholz & Kronberg 1990, ApJ, 357, L13. VLA at 1.4 GHz (1''.8).		
Hester <i>et al.</i> 1990, ApJ, 357, 539. Optical and IR images.		
Hickson & van den Bergh 1990, ApJ, 365, 224. Optical polarisation.		
Bietenholz & Kronberg 1992, ApJ, 393, 206. VLA at 1.5 and 5 GHz (1''.8) and 1.5 and 14 GHz (6''.5) for spectral studies.		
Fesen & Staker 1993, MNRAS, 263, 69. [OIII] imaging of ‘jet’, and proper motion studies.		
Kassim <i>et al.</i> 1993, AJ, 106, 2218. VLA at 74 MHz (20'').		
Véron-Cetty & Woltjer 1993, A&A, 270, 370. Continuum and [OIII] photometry.		
Wallace <i>et al.</i> 1994, A&A, 286, 565. H I of surroundings.		
Frail <i>et al.</i> 1995, ApJ, 454, L129. VLA at 333 MHz (20'') for limits on shell.		
Bietenholz <i>et al.</i> 1997, ApJ, 490, 291. Comparison of VLA observations at 74 MHz, 327 MHz, 1.5 GHz and 5 GHz for spectral index studies.		
Blair <i>et al.</i> 1997, ApJS, 109, 473. HST imaging.		
Fesen <i>et al.</i> 1997, AJ, 113, 354. Limits on Hα halo.		
van der Meulen <i>et al.</i> 1998, A&A, 330, 321. γ-ray observations.		
Tanimori <i>et al.</i> 1998, ApJ, 492, L33. γ-ray observations.		
Nugent 1998, PASP, 110, 831. Optical expansion.		
Sankrit <i>et al.</i> 1998, ApJ, 504, 344. HST images.		
Wallace <i>et al.</i> 1999, ApJS, 124, 181. DRAO at 1.4 GHz (1''.0×2''.8), plus Effelsberg 100-m, for H I studies.		
Greindl & Aschenbach 1999, ApJ, 510, 305. X-ray variability of torus.		
Weisskopf <i>et al.</i> 2000, ApJ, 536, L81. Chandra observations.		
Sollerman <i>et al.</i> 2000, ApJ, 537, 861. HST observations.		
Aharonian <i>et al.</i> 2000, ApJ, 539, 317. γ-ray observations.		
Oser <i>et al.</i> 2001, ApJ, 547, 949. γ-ray observations.		
Willingale <i>et al.</i> 2001, A&A, 365, L212. XMM-Newton observations.		
Douvion <i>et al.</i> 2001, A&A, 373, 281. ISO observations.		
Bietenholz <i>et al.</i> 2001, ApJ, 560, 254. Multi-epoch VLA observations, showing variations near the pulsar.		
Bandiera <i>et al.</i> 2002, A&A, 386, 1044. 1.3 mm observations.		
Hester <i>et al.</i> 2002, ApJ, 577, L49. HST and Chandra multi-epoch observations.		
Atkins <i>et al.</i> 2003, ApJ, 595, 803. γ-ray observations.		
Green <i>et al.</i> 2004, MNRAS, 355, 1315. Sub-mm and ISO observations.		
Mori <i>et al.</i> 2004, ApJ, 609, 186. Chandra observations.		
Čadež <i>et al.</i> 2004, ApJ, 609, 797. Optical observations.		
Aharonian <i>et al.</i> 2004, ApJ, 614, 897. γ-ray observations.		
Bietenholz <i>et al.</i> 2004, ApJ, 615, 794. VLA at 5 GHz (1''.4) and HST multi-epoch observation for proper motion studies.		
Melatos <i>et al.</i> 2005, ApJ, 633, 931. Multi-epoch near-IR observations central region.		
Seward <i>et al.</i> 2006, ApJ, 636, 873. Chandra observations of scattering halo.		
Temim <i>et al.</i> 2006, AJ, 132, 1610. Spitzer observations. see also: Temim <i>et al.</i> 2009, AJ, 137, 5155. Erratum.		
Aharonian <i>et al.</i> 2006, A&A, 457, 899. γ-ray observations.		
Seward <i>et al.</i> 2006, ApJ, 652, 1277. Chandra observations.		
MacAlpine <i>et al.</i> 2007, AJ, 133, 81. Optical spectroscopy.		
Kaplan <i>et al.</i> 2008, ApJ, 677, 1201. HST proper motion of pulsar.		
Rudie <i>et al.</i> 2008, MNRAS, 384, 1200. [OIII] observations of ‘jet’, for proper motion.		
Hurley-Walker <i>et al.</i> 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.		
Tziampzis <i>et al.</i> 2009, A&A, 497, 167. Limits on Hα halo.		
Carlebois <i>et al.</i> 2010, AJ, 139, 2083. Optical imaging spectroscopy.		
Aumont <i>et al.</i> 2010, A&A, 514, A70. IRAM 30-m at 150 GHz (16''.7:S=244±24 Jy) and GBT at 90 GHz (9''.3).		
Satterfield <i>et al.</i> 2012, AJ, 144, 27. Optical spectroscopy.		
Temim <i>et al.</i> 2012, ApJ, 753, 72. Spitzer observations of dust.		
Gomez <i>et al.</i> 2012, ApJ, 760, 96. Herschel IR and sub-mm observations of dust.		
Loh <i>et al.</i> 2012, MNRAS, 421, 789. IR observations of H ₂ .		
Lundqvist & Tziampzis 2012, MNRAS, 423, 1571. Optical limits on outer shell.		
Loll <i>et al.</i> 2013, ApJ, 765, 152. HST optical line and continuum observations.		
Barlow <i>et al.</i> 2013, Science, 342, 1343. Herschel far-IR spectroscopy.		
Black & Fesen 2015, MNRAS, 447, 2540. [OIII] observations of ‘jet’.		
Owen & Barlow 2015, ApJ, 801, 141. Herschel and Spitzer observations.		
Bietenholz & Nugent 2015, MNRAS, 454, 2416. Multi-epoch VLA and optical observations for expansion study.		
Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux densities at 9 frequencies between 30 and 857 GHz.		

G189.1+3.0		IC443, 3C157
RA: 06 ^h 17 ^m 00 ^s	1-GHz flux/Jy: 165	Size/arcmin: 45
Dec: +22°34'	Spectral index: 0.36	Type: C
Radio: Limb-brightened to NE, with faint extension to the E.		
Optical: Brightest to the NE, with faint filaments outside the NE boundary.		
X-ray: Shell, brightest to the NE, plus compact source with nebula.		
Point sources: X-ray source and nebula in S.		
Distance: Mean optical velocity suggests 0.7–1.5 kpc, association with S249 gives 1.5–2 kpc.		
References:		
Fesen 1984, ApJ, 281, 658. Optical of filament to far NE.	Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.	
Erickson & Mahoney 1985, ApJ, 290, 596. TPT at 4 frequencies between 31 and 74 MHz (13'×11' to 5.4×4.7'), plus review of flux densities.	Lee <i>et al.</i> 2008, AJ, 135, 796. VLA at 1.4 GHz (39''×42'') including H _i .	
Braun & Strom 1986, A&A, 164, 193. WSRT at 327 MHz (72''×185'') and 1.4 GHz (17''×43''), plus H _i and IRAS.	Bocchino <i>et al.</i> 2008, AdSpR, 41, 396. INTEGRAL observations.	
Green 1986, MNRAS, 221, 473. 151 MHz observations (1.2×3.1) and Half-Mile Telescope at 1.4 GHz (2.1×5.4).	Bocchino <i>et al.</i> 2009, A&A, 498, 139. XMM-Newton observations.	
Mufson <i>et al.</i> 1986, AJ, 92, 1349. Radio, IR, optical, UV and X-ray comparison, including VLA at 1.6 GHz (3''.8×3''.3 and 40'').	Hirschauer <i>et al.</i> 2009, ApJ, 696, 1533. Optical absorption spectroscopy.	
Petre <i>et al.</i> 1988, ApJ, 335, 215. Einstein and other X-ray observations.	Acciari <i>et al.</i> 2009, ApJ, 698, L133. γ -ray observations.	
Dickel <i>et al.</i> 1989, AJ, 98, 1363. VLA at 1.4 GHz (1''.1×1''.2) of NE.	Yamaguchi <i>et al.</i> 2009, ApJ, 705, L6. Suzaku observations.	
Wood <i>et al.</i> 1991, AJ, 102, 224. VLA at 5 GHz (3''.6×3''.8) of northeast, including polarisation.	Koo <i>et al.</i> 2010, AJ, 140, 262. H _i Zeeman splitting observations.	
van Dishoeck <i>et al.</i> 1993, A&A, 279, 541. Observations of shocked molecular species.	Tavani <i>et al.</i> 2010, ApJ, 710, L151. γ -ray observations.	
Asaoka & Aschenbach 1994, A&A, 284, 573. X-ray, including possible overlapping remnant.	Gao <i>et al.</i> 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5 : $S = 85 \pm 9$ Jy), including polarisation and review of flux densities.	
Tauber <i>et al.</i> 1994, ApJ, 421, 570. Observations of shocked molecular species.	Castelletti <i>et al.</i> 2011, A&A, 534, A21. VLA at 74 MHz (35' : $S = 470 \pm 51$ Jy) and 330 MHz (17' : $S = 248 \pm 15$ Jy), plus review of flux densities.	
Seta <i>et al.</i> 1998, ApJ, 505, 286. CO observations of surroundings.	Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.	
Claussen <i>et al.</i> 1999, ApJ, 522, 349. High resolution observations of OH masers.	Xu <i>et al.</i> 2011, ApJ, 727, 81. CO observations of region.	
Cesarsky <i>et al.</i> 1999, A&A, 348, 945. ISO observations of shocked molecular H ₂ .	Shinn <i>et al.</i> 2011, ApJ, 732, 124. Akari observations of H ₂ .	
Rho <i>et al.</i> 2001, ApJ, 547, 885. ISO and 2Mass IR spectroscopy and imaging.	Lee <i>et al.</i> 2012, ApJ, 749, 34. CO and HCO ⁺ of region.	
Olbert <i>et al.</i> 2001, ApJ, 554, L205. Chandra of compact X-ray source and surrounding nebula.	Taylor <i>et al.</i> 2012, ApJ, 750, L15. Optical absorption of background stars.	
Reich <i>et al.</i> 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14.5' : $S = 160 \pm 5$ Jy).	Yuan <i>et al.</i> 2012, ApJ, 753, 126. Spitzer spectroscopy.	
Welsh & Sallmen 2003, A&A, 408, 545. Optical absorption studies.	Shinn <i>et al.</i> 2012, ApJ, 759, 34. Akari observations of H ₂ .	
Leahy 2004, AJ, 127, 2277. DRAO at 408 MHz (3'.3×8'.6) and 1.4 GHz (1'.0×2'.6), for spectral index studies. see also: Leahy 2004, AJ, 128, 1478. Addendum.	Ackermann <i>et al.</i> 2013, Science, 339, 807. Fermi observations.	
Snell <i>et al.</i> 2005, ApJ, 620, 758. Molecular line and IR observations of shocked material.	Hezareh <i>et al.</i> 2013, A&A, 558, A45. CO observations, including polarisation, of region.	
Bykov <i>et al.</i> 2005, ApJ, 624, L41. Chandra observations of compact source.	Kokusho <i>et al.</i> 2013, ApJ, 768, L8. IR observations of [FeII], plus Akari and Spitzer observations.	
Kawasaki <i>et al.</i> 2005, ApJ, 631, 935. ASCA observations.	Pihlström <i>et al.</i> 2014, AJ, 147, 73. VLA search for methanol masers.	
Gaensler <i>et al.</i> 2006, ApJ, 648, 1037. Chandra of X-ray source and nebula.	Ohnishi <i>et al.</i> 2014, ApJ, 784, 74. Suzaku observations.	
Hewitt <i>et al.</i> 2006, ApJ, 652, 1288. GBT at 1.6 and 1.7 GHz (7.2') for OH, and VLA at 330 MHz (74''×64'').	Yamaguchi <i>et al.</i> 2014, ApJ, 785, L27. Suzaku observations.	
Neufeld <i>et al.</i> 2007, ApJ, 664, 890. Spitzer observations.	Su <i>et al.</i> 2014, ApJ, 788, 122. CO observations.	
Rosado <i>et al.</i> 2007, AJ, 133, 89. Observations of shocked H ₂ .	Gusdorf <i>et al.</i> 2014, IAUS, 296, 178. CO observations.	
Albert <i>et al.</i> 2007, ApJ, 664, L87. γ -ray observations.	Mitra <i>et al.</i> 2014, IAUS, 296, 376. Combined GMRT and CLFST observations at 150 MHz (19''×24'').	
Troja <i>et al.</i> 2008, A&A, 485, 777. XMM-Newton observations.	Swartz <i>et al.</i> 2015, ApJ, 808, 84. Chandra observations of compact source and nebula.	
Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.	Kokusho <i>et al.</i> 2015, P&SS, 116, 92. IR [FeII] observations.	
Bykov <i>et al.</i> 2008, ApJ, 676, 1050. XMM-Newton, Chandra and Spitzer observations.	Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux densities at 9 frequencies between 30 and 857 GHz.	
	Kilpatrick <i>et al.</i> 2016, ApJ, 816, 1. CO observations, including broad lines.	
	Koo <i>et al.</i> 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).	
	McEwen <i>et al.</i> 2016, ApJ, 826, 189. NH ₃ and CH ₃ OH observations.	
	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.	

G190.9–2.2

RA: 06^h01^m55^s
Dec: +18°24'

1-GHz flux/Jy: 1.3?
Spectral index: 0.7?

Size/arcmin: 70×60
Type: S

Radio: Incomplete shell.

References:

Foster *et al.* 2013, A&A, 549, A107. Various radio survey obser-

G205.5+0.5

RA: 06^h39^m00^s
Dec: +06°30'

1-GHz flux/Jy: 140
Spectral index: 0.4

Monoceros Nebula
Size/arcmin: 220
Type: S

Radio: In complex region, parts may be HII regions.

Optical: Large ring, near Rosette nebula.

X-ray: Possibly detected.

Distance: Mean optical velocity suggests 0.8 kpc, low frequency radio absorption suggests 1.6 kpc.

References:

Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (9').
Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5'), part only.
Dicke & DeNoyer 1975, AJ, 80, 437. Arecibo at 111 MHz (1': S=462±180 Jy) and S_{610 MHz}=245 Jy.
Davies *et al.* 1978, A&AS, 31, 271. Deep optical plates.
Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
Graham *et al.* 1982, A&A, 109, 145. Effelsberg 100-m at 2.7 GHz (4.4': S=97.6±12.5 Jy), plus review of flux densities.
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
Leahy *et al.* 1986, MNRAS, 220, 501. Einstein observations.
Odegard 1986, ApJ, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24', 19' and 16').
Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ-ray

emission.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Oliver *et al.* 1996, A&A, 315, 578. CO observations of some of surroundings.
Jaffe *et al.* 1997, ApJ, 484, L129. γ-ray detection.
Aharonian *et al.* 2004, A&A, 417, 973. γ-ray limits.
Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
Casandjian & Grenier 2008, A&A, 489, 849. γ-ray observations.
Xiao & Zhu 2012, A&A, 545, A86. Review of radio, HI and Hα observations.
Dirks & Meyer 2016, ApJ, 819, 45. Time variation of optical line absorption.
Katagiri *et al.* 2016, ApJ, 831, 106. γ-ray observations.
Acero *et al.* 2016, ApJS, 224, 8. γ-ray observations.

G206.9+2.3

RA: 06^h48^m40^s
Dec: +06°26'

1-GHz flux/Jy: 6
Spectral index: 0.5

PKS 0646+06

Size/arcmin: 60×40
Type: S?

Radio: Diffuse source near the Monoceros Nebula.

Optical: Filaments detected.

X-ray: Possibly detected.

References:

Davies & Meaburn 1978, A&A, 69, 443. Optical observations.
Nousek *et al.* 1981, ApJ, 248, 152. HEAO-1 X-ray limit.
Graham *et al.* 1982, A&A, 109, 145. Effelsberg 100-m at 2.7 GHz (4.4': S=4.1±0.6 Jy), plus review of flux densities.
Rosado 1982, RMxAA, 5, 127. Optical observations.
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
Leahy 1986, A&A, 156, 191. Einstein observations.
Odegard 1986, ApJ, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz

(24', 19' and 16').
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9.5': S=2.9±0.3 Jy), including polarisation and review of flux densities.
Ambrocio-Cruz *et al.* 2014, RMxAA, 50, 323. [SII] spectroscopy.

G213.0–0.6

RA: 06^h50^m50^s
Dec: −00°30'

1-GHz flux/Jy: 21
Spectral index: 0.4

Size/arcmin: 160×140?
Type: S

Has also been called G213.3–0.4.

Radio: Large, faint shell.

Optical: Filamentary shell.

Point sources: Central X-ray source

References:

- Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5 : $S = 22.0 \pm 3.7$ Jy).
 Stupar & Parker 2012, MNRAS, 419, 1413. H α , radio survey and other observations.

G260.4–3.4

RA: 08^h22^m10^s
Dec: −43°00'

1-GHz flux/Jy: 130
Spectral index: 0.5

Puppis A, MSH 08–44
Size/arcmin: 60×50
Type: S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Angular shell, brightest to the E, poorly defined to the W.

Optical: Nebulosity and wisps.

X-ray: Brightest to the E.

Point sources: Central possible pulsating X-ray source.

Distance: Association with H α gives 2.2 kpc.

References:

- Green 1971, AuJPh, 24, 773. Molonglo at 408 MHz (3' : $S = 198 \pm 20$ Jy).
 Goudis & Meaburn 1978, A&A, 62, 283. H α +[NII] optical image.
 Petre *et al.* 1982, ApJ, 258, 22. Einstein observations.
 Milne *et al.* 1983, MNRAS, 204, 237. FIRST at 1415 MHz (50''), large scale emission missing.
 Teske & Petre 1987, ApJ, 318, 370. Coronal optical line emission.
 Dubner & Arnal 1988, A&AS, 75, 363. H α and CO observations of surroundings.
 Arendt *et al.* 1990, ApJ, 350, 266. MOST at 843 MHz (44''×65''), with large scale emission added, plus IR, optical and X-ray observations.
 Dubner *et al.* 1991, AJ, 101, 1466. VLA at 327 MHz (43''×73'') and 1.5 GHz (43''×77'').
 Arendt *et al.* 1991, ApJ, 368, 474. IR observations.
 Milne *et al.* 1993, MNRAS, 261, 366. Parkes 64-m at 4.75 (4.5' : $S = 59 \pm 5$ Jy) and 8.4 GHz (3' : $S = 38 \pm 4$ Jy), plus polarisation.
 Berthiaume *et al.* 1994, ApJ, 425, 132. X-ray spectroscopy.
 Sutherland & Dopita 1995, ApJ, 439, 365. Spectrophotometry.
 Reynolds *et al.* 1995, AJ, 110, 318. VLA at 1.4 GHz (90'') including neutral hydrogen.
 Blair *et al.* 1995, ApJ, 454, L35. Far UV spectroscopy.
 Petre *et al.* 1996, ApJ, 465, L43. ROSAT of central source.
 Bock *et al.* 1998, AJ, 116, 1886. MOST at 843 MHz (43''×60'').
 Pavlov *et al.* 1999, ApJ, 511, L45. Possible pulsation detection from central X-ray source.
 Zavlin *et al.* 1999, ApJ, 525, 959. X-ray observations of central source.
 Bocchino *et al.* 2000, A&A, 359, 316. Optical studies of selected filaments in N.
 Woermann *et al.* 2000, MNRAS, 317, 421. OH observations.
 Gaensler *et al.* 2000, ApJ, 537, L35. Radio limit for nebula around possible pulsar.
 Reynolds *et al.* 2003, MNRAS, 345, 671. ATCA at 1.4 GHz (90'') for H α near central X-ray source.
 Hwang *et al.* 2005, ApJ, 635, 355. Chandra observations of E edge.
 Hui & Becker 2006, A&A, 454, 543. XMM-Newton and Chandra observations.
 Hui & Becker 2006, A&A, 457, L33. Chandra proper motion study of central source.
 Castelletti *et al.* 2006, A&A, 459, 535. VLA at 1.4 GHz (16''×34'' : $S = 114 \pm 8$ Jy) and 327 MHz (45''×90'' : $S = 263 \pm 20$ Jy).
 Winkler & Petre 2007, ApJ, 670, 635. Chandra proper motion study of central source.
 Paron *et al.* 2008, A&A, 480, 439. CO observations of E.
 Hwang *et al.* 2008, ApJ, 676, 378. Suzaku observations.
 Katsuda *et al.* 2008, ApJ, 678, 297. XMM-Newton observations.
 Mignani *et al.* 2009, A&A, 500, 1211. Optical limits for compact X-ray source.
 Katsuda *et al.* 2010, ApJ, 714, 1725. Chandra and XMM-Newton observations.
 Arendt *et al.* 2010, ApJ, 725, 585. Spitzer observations.
 Becker *et al.* 2012, ApJ, 755, 141. Chandra proper motion study of central X-ray source.
 Katsuda *et al.* 2012, ApJ, 756, 49. XMM-Newton spectroscopy.
 Hewitt *et al.* 2012, ApJ, 759, 89. Fermi and WMAP 23 to 90 GHz observations.
 Dubner *et al.* 2013, A&A, 555, A9. XMM-Newton and Chandra observations.
 Katsuda *et al.* 2013, ApJ, 768, 182. XMM-Newton observations.
 Abramowski *et al.* 2015, A&A, 575, A81. High energy γ -ray limit.
 Reynolds & Walsh 2015, MNRAS, 451, 3044. ATCA at 1.4 GHz (51''×82'') and 1.7 GHz for spectral index study.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 4 frequencies between 30 and 100 GHz.
 Luna *et al.* 2016, A&A, 590, A70. XMM-Newton and Chandra observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G261.9+5.5

RA: 09^h04^m20^s
Dec: -38°42'

1-GHz flux/Jy: 10?
Spectral index: 0.4?

Size/arcmin: 40×30
Type: S

Radio: Faint shell with little limb brightening.

References:

- Hill 1967, AuJPh, 20, 297. Parkes 64-m at 2650 MHz ($7'.5 : S = 7$ Jy) also $S_{1410 \text{ MHz}} = 8$ Jy, $S_{81.5 \text{ MHz}} = 25$ Jy.
 Colomb & Dubner 1980, A&A, 82, 244. Argentine 30-m dish at 1.4 GHz, for H_I possibly associated with remnant.
- Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 71''$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G263.9–3.3		Vela (XYZ)
RA: 08 ^h 34 ^m 00 ^s	1-GHz flux/Jy: 1750	Size/arcmin: 255
Dec: −45°50'	Spectral index: varies	Type: C
This refers to the whole Vela XYZ complex, of which X has at times been classified as a separate (filled-centre) remnant. This remnant is overlapped by G260.4–3.4 and G266.2–1.2.		
Radio: Large shell, with flatter spectrum component (Vela X), and pulsar nebula.		
Optical: Filaments.		
X-ray: Patchy shell, with extensions, central nebula and pulsar.		
Point sources: Pulsar within Vela X, with one-sided ‘jet’.		
Distance: Vela pulsar parallax gives 0.3 kpc, optical spectra and H _I studies suggest 0.25 kpc.		
References:		
Milne 1968, <i>AuJPh</i> , 21, 201. Parkes 64-m at various frequencies, including 408 MHz ($48' : S = 2300 \pm 300$ Jy), 635 MHz ($31' : S = 2360 \pm 300$ Jy), 1410 MHz ($14' : S = 1640 \pm 300$ Jy) and 2650 MHz ($7.5' : S = 1400 \pm 250$ Jy), plus discussion of the distance.	Cha <i>et al.</i> 1999, <i>ApJ</i> , 515, L25. Optical spectra, for distance.	
Milne 1980, <i>A&A</i> , 81, 293. Maps of Vela X with Parkes 64-m at 1.66, 2.7 and 5 GHz ($12', 8.4'$ and $4.4'$).	Lu & Aschenbach 2000, <i>A&A</i> , 362, 1083. ROSAT spatially resolved spectroscopy.	
Weiler & Panagia 1980, <i>A&A</i> , 90, 269. Clarification of notation of this region and review previous observations.	Chadwick <i>et al.</i> 2000, <i>ApJ</i> , 537, 414. Limit on high energy γ -rays from pulsar.	
Bignami & Caraveo 1988, <i>ApJ</i> , 325, L5. Pulsar proper motion from optical observations.	Sankrit <i>et al.</i> 2001, <i>ApJ</i> , 549, 416. Far-UV observations of selected region.	
Dwarkanath 1991, <i>JApA</i> , 12, 199. Gauribidanur ‘T’ array at 34.5 MHz ($26' \times 84' : S = 1800$ Jy for Vela X and $S = 3900$ Jy for Vela YZ), plus review of flux densities.	Pavlov <i>et al.</i> 2001, <i>ApJ</i> , 554, L189. Chandra two epoch observations of pulsar nebula.	
Bietenholz <i>et al.</i> 1991, <i>ApJ</i> , 376, L41. VLA at 5 GHz in vicinity of pulsar.	Helfand <i>et al.</i> 2001, <i>ApJ</i> , 556, 380. Chandra observations of pulsar and its nebula.	
Willmore <i>et al.</i> 1992, <i>MNRAS</i> , 254, 139. Hard X-ray observations.	Miyata <i>et al.</i> 2001, <i>ApJ</i> , 559, L45. Chandra of X-ray extension.	
Dubner <i>et al.</i> 1992, <i>A&AS</i> , 96, 505. Argentine 30-m at 1.4 GHz ($30'$) of surrounding H _I .	Alvarez <i>et al.</i> 2001, <i>A&A</i> , 372, 636. Radio spectral index studies. see also: Alvarez <i>et al.</i> 2001, <i>A&A</i> , 379, 323. Erratum.	
Strom <i>et al.</i> 1995, <i>Nature</i> , 373, 590. Radio of X-ray extensions.	Caraveo <i>et al.</i> 2001, <i>ApJ</i> , 561, 930. HST parallax observations of pulsar.	
Markwardt & Ögelman 1995, <i>Nature</i> , 375, 40. X-ray jet from pulsar.	Moriguchi <i>et al.</i> 2001, <i>PASJ</i> , 53, 1025. CO observations.	
Jenkins & Wallerstein 1995, <i>ApJ</i> , 440, 227. Optical absorption of associated neutral carbon cloud.	Dodson <i>et al.</i> 2003, <i>MNRAS</i> , 343, 116. ATCA at 1.4, 2.4, 5.2 and 8.5 GHz ($6.2' \times 8.1'$, $26' \times 36''$, $12.1' \times 10.5''$ and $11.2' \times 10.6''$) of pulsar nebula.	
Milne 1995, <i>MNRAS</i> , 277, 1435. Parkes 64-m at 8.4 GHz ($3'$), including polarisation, of Vela X.	Sankrit <i>et al.</i> 2003, <i>ApJ</i> , 589, 242. Optical nebulosity to NE of G266.2–1.2.	
Danks & Sembach 1995, <i>AJ</i> , 109, 2627. Optical spectroscopy of background stars.	Nichols & Slavin 2004, <i>ApJ</i> , 610, 285. UV absorption toward background sources.	
Blair <i>et al.</i> 1995, <i>AJ</i> , 110, 312. UV spectroscopy.	Hales <i>et al.</i> 2004, <i>ApJ</i> , 613, 977. Vela X at 31 GHz ($4.1'$).	
Duncan <i>et al.</i> 1996, <i>MNRAS</i> , 280, 252. Parkes 64-m at 2.4 GHz ($8.9'$).	Mongano <i>et al.</i> 2005, <i>A&A</i> , 436, 917. XMM-Newton and other X-ray observations of pulsar nebula.	
Whiteoak & Green 1996, <i>A&AS</i> , 118, 329. MOST at 843 MHz ($43' \times 62'$) of part.	Miceli <i>et al.</i> 2005, <i>A&A</i> , 442, 513. XMM-Newton observations of N rim.	
Green <i>et al.</i> 1997, <i>AJ</i> , 114, 2058. Parkes 64-m and ATCA OH observations.	Katsuda & Tsunemi 2005, <i>PASJ</i> , 57, 621. XMM-Newton observations of E.	
Frai <i>et al.</i> 1997, <i>ApJ</i> , 475, 224. VLA at 330 MHz ($1' \times 1.8'$) and comparison with ROSAT observations of Vela X.	Aharonian <i>et al.</i> 2006, <i>A&A</i> , 448, L43. γ -ray observations.	
Markwardt & Ögelman 1997, <i>ApJ</i> , 480, L13. ASCA observations of pulsar ‘jet’.	McConnell <i>et al.</i> 2006, <i>AJ</i> , 131, 648. ATCA at 4.9 GHz ($12'$) including polarisation.	
Raymond <i>et al.</i> 1997, <i>ApJ</i> , 482, 881. UV spectroscopy.	Katsuda & Tsunemi 2006, <i>ApJ</i> , 642, 917. XMM-Newton observations of NE.	
Yoshikoshi <i>et al.</i> 1997, <i>ApJ</i> , 487, L65. γ -ray observations of pulsar.	Nishikida <i>et al.</i> 2006, <i>ApJ</i> , 644, L171. Far UV observations.	
Jenkins <i>et al.</i> 1998, <i>ApJ</i> , 492, L147. UV absorption to background star.	Miceli <i>et al.</i> 2008, <i>ApJ</i> , 676, 1064. XMM-Newton observations.	
Dubner <i>et al.</i> 1998, <i>AJ</i> , 116, 813. Parkes 64-m at 1.4 GHz ($15'$) for H _I studies.	LaMassa <i>et al.</i> 2008, <i>ApJ</i> , 689, L121. XMM-Newton observations of Vela X.	
Bock <i>et al.</i> 1998, <i>AJ</i> , 116, 1886. MOST at 843 MHz ($43' \times 60''$).	Grondin <i>et al.</i> 2013, <i>ApJ</i> , 774, 110. Fermi observations of Vela X.	
	Arnaud <i>et al.</i> 2016, <i>A&A</i> , 586, A134. Planck flux densities at 30, 44 and 70 GHz.	
	Rao <i>et al.</i> 2016, <i>MNRAS</i> , 455, 2529. Time variation of optical line absorption.	

G266.2–1.2	RX J0852.0–4622
RA: 08 ^h 52 ^m 00 ^s	1-GHz flux/Jy: 50?
Dec: −46°20'	Spectral index: 0.3?
This remnant overlaps the Vela SNR (G263.9–3.3).	
Radio: Incomplete shell, confused by the Vela SNR.	
Optical: Nebulosity offset to NE.	
X-ray: Non-thermal shell, confused by the Vela SNR, with central source, and possible associated pulsar.	
Point sources: Central X-ray source, with optical nebula, and possible associated pulsar.	
Distance: X-ray data suggest an upper limit of 1 kpc.	
References:	
Aschenbach 1998, <i>Nature</i> , 396, 141. ROSAT identification.	Redman & Meaburn 2005, <i>MNRAS</i> , 356, 969. Possible pulsar association.
Iyudin <i>et al.</i> 1998, <i>Nature</i> , 396, 142. γ-ray observations.	Iyudin <i>et al.</i> 2005, <i>A&A</i> , 429, 225. XMM-Newton observations.
Combi <i>et al.</i> 1999, <i>ApJ</i> , 519, L177. Radio observations.	Aharonian <i>et al.</i> 2005, <i>A&A</i> , 437, L7. γ-ray observations.
Duncan & Green 2000, <i>A&A</i> , 364, 732. Parkes 64-m at 1.4 GHz (14''), and comparison with other observations.	Katagiri <i>et al.</i> 2005, <i>ApJ</i> , 619, L163. γ-ray observations.
Redman <i>et al.</i> 2000, <i>ApJ</i> , 543, L153. Optical of nearly nebulosity.	Bamba <i>et al.</i> 2005, <i>ApJ</i> , 632, 294. Chandra of NW rim.
Tsunemi <i>et al.</i> 2000, <i>PASJ</i> , 52, 887. ASCA spectral observations.	Reynoso <i>et al.</i> 2006, <i>A&A</i> , 449, 243. ATCA at 1.38 GHz (32'' × 37'').
Slane <i>et al.</i> 2001, <i>ApJ</i> , 548, 814. ASCA observations.	Enomoto <i>et al.</i> 2006, <i>ApJ</i> , 652, 1268. γ-ray observations.
Mereghetti <i>et al.</i> 2001, <i>ApJ</i> , 548, L213. BeppoSAX observations of central sources.	Mignani <i>et al.</i> 2007, <i>A&A</i> , 473, 883. Deep optical observations of compact X-ray source.
Pavlov <i>et al.</i> 2001, <i>ApJ</i> , 559, L131. Chandra of central X-ray source.	Aharonian <i>et al.</i> 2007, <i>ApJ</i> , 661, 236. γ-ray observations.
Moriguchi <i>et al.</i> 2001, <i>PASJ</i> , 53, 1025. CO observations.	Katsuda <i>et al.</i> 2008, <i>ApJ</i> , 678, L35. XMM-Newton proper motion study.
Pellizzoni <i>et al.</i> 2002, <i>A&A</i> , 393, L65. Optical observations of central source.	Pannuti <i>et al.</i> 2010, <i>ApJ</i> , 721, 1492. Chandra observations of NW.
Redman <i>et al.</i> 2002, <i>MNRAS</i> , 336, 1093. Optical nebulosity to NE.	Allen <i>et al.</i> 2015, <i>ApJ</i> , 798, 82. Two epoch Chandra observations for expansion.
Kargaltsev <i>et al.</i> 2002, <i>ApJ</i> , 580, 1060. Chandra observations of central source.	Acero <i>et al.</i> 2016, <i>ApJS</i> , 224, 8. γ-ray observations.
Sankrit <i>et al.</i> 2003, <i>ApJ</i> , 589, 242. Optical nebulosity to NE.	Takeda <i>et al.</i> 2016, <i>PASJ</i> , 68, S10. Suzaku observations.

G272.2–3.2	Size/arcmin: 15?
RA: 09 ^h 06 ^m 50 ^s	Type: S?
Dec: −52°07'	
1-GHz flux/Jy: 0.4	
Spectral index: 0.6	

Radio: Diffuse shell.	
Optical: Detected.	
X-ray: Centrally brightened.	
References:	
Greiner <i>et al.</i> 1994, <i>A&A</i> , 286, L35. ROSAT observations, plus optical observations.	(45'' × 70'': $S = 0.45 \pm 0.10$ Jy), and ATCA at 2.4 GHz (37'' × 52''), plus ROSAT observations.
Duncan <i>et al.</i> 1997, <i>MNRAS</i> , 289, 97. Parkes 64-m at 1.4 GHz (18'': $S = 0.38 \pm 0.09$ Jy) 2.4 GHz (10'': $S = 0.25 \pm 0.04$ Jy) and 4.8 GHz (5.7'': $S = 0.17 \pm 0.02$ Jy), MOST at 843 MHz	Harrus <i>et al.</i> 2001, <i>ApJ</i> , 552, 614. ASCA and ROSAT observations, plus review of earlier observations.
	Kamitsukasa <i>et al.</i> 2016, <i>PASJ</i> , 68, S7. Suzaku observations.

G279.0+1.1	Size/arcmin: 95
RA: 09 ^h 57 ^m 40 ^s	Type: S
Dec: −53°15'	
1-GHz flux/Jy: 30?	
Spectral index: 0.6?	

Radio: Faint, incomplete shell.	
Optical: Detected.	
Point sources: Pulsar nearby.	
References:	
Woermann & Jonas 1988, <i>MNRAS</i> , 234, 971. Hartesbeesthoek 26-m at 1.6 (30'': $S = 25.2 \pm 4$ Jy) and 2.3 GHz (20'': $S = 20.7 \pm 3$ Jy).	Whiteoak & Green 1996, <i>A&AS</i> , 118, 329. MOST at 843 MHz (43'' × 53'') of part.
Duncan <i>et al.</i> 1995, <i>MNRAS</i> , 277, 319. Parkes 64-m at 1.4 (18'': $S = 28 \pm 3$ Jy) and 2.4 GHz (11'': $S = 20 \pm 2$ Jy), including polarisation.	Green <i>et al.</i> 1997, <i>AJ</i> , 114, 2058. Parkes 64-m OH observations.
	Stupar & Parker 2011, <i>MNRAS</i> , 414, 2282. Hα observations.

G284.3–1.8	MSH 10–53
RA: $10^{\text{h}}18^{\text{m}}15^{\text{s}}$	1-GHz flux/Jy: 11?
Dec: $-59^{\circ}00'$	Spectral index: 0.3?
Has been called G284.2–1.8.	
Radio: Incomplete, poorly defined shell.	
Optical: Faint filament near edge.	
Point sources: Central X-/γ-ray source.	
References:	
Ruiz & May 1986, ApJ, 309, 667. CO and optical observations.	Camilo <i>et al.</i> 2001, ApJ, 557, L51. Observations of nearby pulsar.
Milne <i>et al.</i> 1989, PASA, 8, 187. MOST at 843 MHz ($43'' \times 50''$) and Parkes 64-m at 8.4 GHz ($3' : S = 5.4 \pm 0.8$ Jy) including polarisation, plus earlier flux densities.	Camilo <i>et al.</i> 2004, ApJ, 616, 1118. Chandra observations of pulsar and nebula.
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.	Abramowski <i>et al.</i> 2012, A&A, 541, A5. γ-ray observations.
	Williams <i>et al.</i> 2015, ApJ, 808, L19. Chandra observations.
G286.5–1.2	Size/arcmin: 26×6
RA: $10^{\text{h}}35^{\text{m}}40^{\text{s}}$	Type: S?
Dec: $-59^{\circ}42'$	1-GHz flux/Jy: 1.4?
Optical: Detected.	Spectral index: ?
References:	
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 1.6$ Jy).	
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.	
Stupar & Parker 2011, MNRAS, 414, 2282. Hα observations.	
G289.7–0.3	Size/arcmin: 18×14
RA: $11^{\text{h}}01^{\text{m}}15^{\text{s}}$	Type: S
Dec: $-60^{\circ}18'$	1-GHz flux/Jy: 6.2
Point sources: Compact radio source near centre.	Spectral index: 0.2?
References:	
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 6.4 \pm 0.5$ Jy), plus Parkes 64-m at 4.5 GHz ($S = 7.5 \pm 2.5$ Jy) and 8.55 GHz ($S = 3.6 \pm 0.9$ Jy).	

G290.1–0.8		MSH 11–61A
RA: 11 ^h 03 ^m 05 ^s	1-GHz flux/Jy: 42	Size/arcmin: 19×14
Dec: −60°56'	Spectral index: 0.4	Type: S
Radio: Elongated, clumpy shell.		
Optical: Filaments detected.		
X-ray: Centrally brightened.		
Point sources: Pulsar nearby, with PWN and extended ‘jet’ in X-rays.		
Distance: H _I absorption indicates 7±1 kpc.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').	Gotthelf & Kaspi 1998, ApJ, 497, L29. ASCA observations of pulsar.	
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4.4': $S = 20.2$ Jy).	Slane <i>et al.</i> 2002, ApJ, 564, 284. ASCA observations.	
Elliott & Malin 1979, MNRAS, 186, 45P. Optical image and spectra.	Filipović <i>et al.</i> 2005, SerAJ, 170, 47. ATCA at 1.4 GHz (21''), plus other observations, including CO of surroundings.	
Kirshner & Winkler 1979, ApJ, 227, 853. Optical observations.	Reynoso <i>et al.</i> 2006, MNRAS, 369, 416. ATCA at 1.4 GHz (22'5×25'0) including H _I .	
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×50'').	Pavan <i>et al.</i> 2014, A&A, 562, A122. Chandra and ATCA observations at 2 GHz (4''.1×5''.2) of PWN.	
Milne <i>et al.</i> 1989, PASA, 8, 187. MOST at 843 MHz (43''×49'': $S = 45 \pm 11$ Jy), and Parkes 64-m at 8.4 GHz (3': $S = 19.5 \pm 1.0$ Jy), including polarisation.	Halpern <i>et al.</i> 2014, ApJ, 795, L27. XMM-Newton observations of PWN.	
Seward 1990, ApJS, 73, 781. Einstein observations.	Auchettl <i>et al.</i> 2015, ApJ, 810, 43. γ -ray and Suzaku observations.	
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'': $S = 43$ Jy).	Kamitsukasa <i>et al.</i> 2015, PASJ, 67, 16. Suzaku observations.	
Rosado <i>et al.</i> 1996, A&A, 315, 243. Optical kinematics.	Pavan <i>et al.</i> 2016, A&A, 591, A91. Chandra observations, including PWN and ‘jet’.	
Kaspi <i>et al.</i> 1997, ApJ, 485, 820. Pulsar detection.		

G291.0–0.1		(MSH 11–62)
RA: 11 ^h 11 ^m 54 ^s	1-GHz flux/Jy: 16	Size/arcmin: 15×13
Dec: −60°38'	Spectral index: 0.29	Type: C
Radio: Centrally brightened core, with surrounding arcs.		
Optical: Detected.		
X-ray: Centrally brightened.		
Point sources: Central compact X-ray source.		
References:		
Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').	Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'': $S = 12.7$ Jy).	
Roger <i>et al.</i> 1986, MNRAS, 219, 815. MOST at 843 MHz (43''×50'': $S = 17.2 \pm 1.0$ Jy), and Parkes 64-m at 5 and 8.4 GHz (4'.6 and 3': $S = 10.4 \pm 0.4$ Jy and 9'.1±0.2), with polarisation.	Harris <i>et al.</i> 1998, ApJ, 499, 273. ASCA observations.	
Wilson 1986, ApJ, 302, 718. Einstein observations.	Stupar & Parker 2011, MNRAS, 414, 2282. H _α observations.	
	Slane <i>et al.</i> 2012, ApJ, 749, 131. Chandra, XMM-Newton and Fermi observations.	
	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.	

G292.0+1.8

RA: 11^h24^m36^s
Dec: -59°16'

1-GHz flux/Jy: 15
Spectral index: 0.4

MSH 11-54

Size/arcmin: 12×8
Type: C

Radio: Centrally brightened source surrounded by a plateau of faint emission.

Optical: Oxygen rich.

X-ray: Ring of emission, with diffuse central nebula and pulsar.

Point sources: Central pulsar.

Distance: H_I absorption implies 6.0 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Lockhart *et al.* 1977, MNRAS, 179, 147. Fleurs at 1415 MHz (50'': $S = 13.0$ Jy).
Goss *et al.* 1979, MNRAS, 188, 357. Optical spectra.
Clark *et al.* 1980, MNRAS, 193, 129. X-ray spectrum.
Tuohy *et al.* 1982, ApJ, 260, L65. Einstein observations.
Dopita & Tuohy 1984, ApJ, 282, 135. Optical spectra.
Braun *et al.* 1986, A&A, 162, 259. MOST at 843 MHz (0'8) and IRAS.
Hughes & Singh 1994, ApJ, 422, 126. EXOSAT spectrum.
Sutherland & Dopita 1995, ApJ, 439, 365. Spectrophotometry.
Hughes *et al.* 2001, ApJ, 559, L153. Chandra observations, including central nebula.
Park *et al.* 2002, ApJ, 564, L39. Chandra observations.
Camilo *et al.* 2002, ApJ, 567, L71. Pulsar detection.
Gonzalez & Safi-Harb 2003, ApJ, 583, L91. Chandra observations.
Hughes *et al.* 2003, ApJ, 591, L139. Chandra observations of pulsar.
Gaensler & Wallace 2003, ApJ, 594, 326. ATCA at 1.4, 2.3 and 5.2 GHz (9''.6×8''.0, 7''.2×6''.2 and 5''.5×4''.8 : $S = 11.9 \pm 0.1, 11.4 \pm 0.1$ and 8.8 ± 0.1), plus H_I observations.
Park *et al.* 2004, ApJ, 602, L33. Chandra observations.

- Vink *et al.* 2004, NuPhS, 132, 62. XMM-Newton observations.
Ghavamian *et al.* 2005, ApJ, 635, 365. Optical imaging spectroscopy.
Winkler & Long 2006, AJ, 132, 360. Optical observations.
Park *et al.* 2007, ApJ, 670, L121. Chandra observations.
Zharikov *et al.* 2008, A&A, 492, 805. Possible optical counterpart to pulsar and nebula.
Winkler *et al.* 2009, ApJ, 692, 1489. Optical proper motion studies.
Ghavamian *et al.* 2009, ApJ, 696, 1307. Spitzer spectroscopy.
Lee *et al.* 2009, ApJ, 706, 441. IR observations.
Lee *et al.* 2010, ApJ, 711, 861. Chandra observations.
Ghavamian *et al.* 2012, ApJ, 750, 39. Spitzer observations.
Zharikov *et al.* 2013, A&A, 554, A120. IR of pulsar wind nebula.
Yamaguchi *et al.* 2014, ApJ, 785, L27. Suzaku observations.
Kamitsukasa *et al.* 2014, PASJ, 66, 64. Suzaku observations.
Bhalerao *et al.* 2015, ApJ, 800, 65. Chandra observations.
Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
Ghavamian & Williams 2016, ApJ, 831, 188. Spitzer observations.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G292.2-0.5

RA: 11^h19^m20^s
Dec: -61°28'

1-GHz flux/Jy: 7
Spectral index: 0.5

Size/arcmin: 20×15
Type: S

Radio: Shell.

X-ray: Shell, brighter to W, with central nebula.

Point sources: Central, young pulsar.

Distance: H_I absorption indicates 8.4 kpc.

References:

- Camilo *et al.* 2000, ApJ, 541, 367. Pulsar detection.
Crawford *et al.* 2001, ApJ, 554, 152. ATCA at 1.4 GHz (29''×25'': $S = 5.6 \times 0.3$ Jy) and 2.5 GHz (21''×20'').
Pivovaroff *et al.* 2001, ApJ, 554, 161. ROSAT and ASCA observations.
Gonzalez & Safi-Harb 2003, ApJ, 591, L143. Chandra observations of pulsar.
Caswell *et al.* 2004, MNRAS, 352, 1405. ATCA at 5 GHz (1' : $S = 2.8$ Jy), including polarisation, and 1.4 GHz for H_I absorption.

- Gonzalez & Safi-Harb 2005, ApJ, 619, 856. Chandra observations.
Gonzalez *et al.* 2005, ApJ, 630, 489. XMM-Newton observations of pulsar.
Safi-Harb *et al.* 2008, ApJ, 684, 532. Chandra observations of pulsar and nebula.
Kumar *et al.* 2012, ApJ, 754, 96. Chandra and XMM-Newton observations.
Ng *et al.* 2012, ApJ, 761, 65. XMM-Newton observations.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G293.8+0.6

RA: 11^h35^m00^s
Dec: -60°54'

1-GHz flux/Jy: 5?
Spectral index: 0.6?

Size/arcmin: 20
Type: C

Radio: Central source, with faint extended plateau.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 9.0$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 2.1$ Jy).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz

($44'' \times 51''$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
($43'' \times 49'' : S = 2.6$ Jy).

G294.1–0.0

RA: 11^h36^m10^s
Dec: -61°38'

1-GHz flux/Jy: >2?
Spectral index: ?

Size/arcmin: 40
Type: S

Radio: Faint shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
($43'' \times 49'' : S > 2$).

G296.1–0.5

RA: 11^h51^m10^s
Dec: -62°34'

1-GHz flux/Jy: 8?
Spectral index: 0.6?

Size/arcmin: 37×25
Type: S

Incorporates the previously catalogued remnant G296.1–0.7. Has been called G296.05–0.50.

Radio: Irregular shell, with nearby HII regions.

Optical: Detected.

X-ray: Irregular, incomplete shell.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 6.9$ Jy) and Parkes 64-m at 5 GHz ($4' : S > 0.74$).
Longmore *et al.* 1977, MNRAS, 181, 541. Optical spectra.
van den Bergh 1978, ApJS, 38, 119. Optical observations.
Markert *et al.* 1981, ApJ, 248, L17. Einstein observations.
Caswell & Barnes 1983, ApJ, 271, L55. Molonglo at 408 MHz
($3' : S = 12.4$ Jy).
Bignami *et al.* 1986, ApJ, 302, 606. EXOSAT and Einstein ob-

servations.
Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
($43'' \times 48'' : S > 2.4$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Castro *et al.* 2011, ApJ, 734, 86. XMM-Newton observations.
Gök & Sezer 2012, MNRAS, 419, 1603. Suzaku observations.

G296.5+10.0

RA: 12^h09^m40^s
Dec: -52°25'

1-GHz flux/Jy: 48
Spectral index: 0.5

Has been called G296.5+9.7.

Radio: Shell with two bright limbs.

Optical: Detected.

X-ray: Incomplete shell, with central pulsar.

Point sources: Central pulsar.

References:

- Irvine & Irvine 1974, ApJ, 192, L111. Optical observations.
Danziger & Dennefeld 1976, PASP, 88, 44. Optical spectra.
Dickey & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
Tuohy *et al.* 1979, ApJ, 230, L27. X-ray detection with HEAO-1 A2 experiment.
Ruiz 1983, AJ, 88, 1210. Optical spectra.
Dubner *et al.* 1986, AJ, 91, 343. Argentine 30-m dish at 1.4 GHz (34'), plus HI.
Klelett *et al.* 1987, MNRAS, 225, 199. EXOSAT of the west of the remnant, including the compact source.
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×56'').
Matsui *et al.* 1988, ApJ, 329, 838. Einstein observations, including compact source.
Roger *et al.* 1988, ApJ, 332, 940. MOST at 843 MHz (44''×56'').
Bignami *et al.* 1992, ApJ, 389, L67. Optical in vicinity of X-ray source.
Milne & Haynes 1994, MNRAS, 270, 106. Parkes 64-m at 2.4 GHz (8'.3 : $S = 33 \pm 3$ Jy), 4.8 GHz (4'.5 : $S = 23.3 \pm 3$ Jy) and 8.4 GHz (3'.0 : 18.8 ± 3), including polarisation and review of flux densities.

PKS 1209–51/52

Size/arcmin: 90×65
Type: S

G296.7–0.9

RA: 11^h55^m30^s
Dec: -63°08'

1-GHz flux/Jy: 3
Spectral index: 0.5

Size/arcmin: 15×8
Type: S

Radio: Bilateral shell.

X-ray: Brighter to SE.

References:

- Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT observations.
Robbins *et al.* 2012, MNRAS, 419, 2623. ATCA at 1.4 GHz (33''×45'' : $S = 2.5 \pm 0.2$ Jy), plus MOST at 843 MHz (43''×49''), plus other observations.

- Prinz & Becker 2013, A&A, 550, A33. XMM-Newton observations.
Green *et al.* 2014, PASA, 31, 42. Radio observations at 843 MHz (45''×50'').

G296.8–0.3

RA: 11^h58^m30^s
Dec: -62°35'

1-GHz flux/Jy: 9
Spectral index: 0.6

1156–62

Size/arcmin: 20×14
Type: S

Radio: Shell, brighter to the NW.

X-ray: Detected.

Distance: HI absorption gives 9.6 kpc.

References:

- Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S = 15.0$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 3.2$ Jy).
Dickey & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

- (43''×48'' : $S = 9.2$ Jy).
Gaensler *et al.* 1998, MNRAS, 296, 813. ATCA at 1.3 GHz (22''×24'' : $S = 7.0 \pm 0.3$ Jy), including polarisation and HI observations, plus review of flux densities.
Sánchez-Ayaso *et al.* 2012, Ap&SS, 337, 573. XMM-Newton and IR observations.

G298.5–0.3

RA: 12^h12^m40^s
Dec: −62°52'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 5?
Type: ?

Radio: Not well resolved, may be part of a larger ring?

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'': $S=1.8$ Jy).
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G298.6–0.0

RA: 12^h13^m41^s
Dec: −62°37'

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 12×9
Type: S

Has been called G298.6–0.1.

Radio: Incomplete shell, in complex region.

X-ray: Centrally brightened.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×50'').
Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'': $S=7.4$ Jy).
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.
Bamba *et al.* 2016, PASJ, 68, S5. Suzaku observations.

G299.2–2.9

RA: 12^h15^m13^s
Dec: −65°30'

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 18×11
Type: S

Radio: Faint source.

Optical: Filaments in W.

X-ray: Centrally brightened with shell at higher energies.

References:

Busser *et al.* 1996, A&A, 310, L1. ROSAT detection, plus optical studies.
Slane *et al.* 1996, ApJ, 465, 840. Einstein, IRAS and radio observations.

Bai & Wang 2000, ApJ, 539, 760. ASCA observations.
Park *et al.* 2007, ApJ, 665, 1173. Chandra observations.
Part *et al.* 2014, ApJ, 792, L20. Chandra observations.

G299.6–0.5

RA: 12^h21^m45^s
Dec: −63°09'

1-GHz flux/Jy: 1.0?
Spectral index: ?

Size/arcmin: 13
Type: S

Radio: Faint shell, brightest to E.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'': $S=1.1$ Jy).

G301.4–1.0

RA: 12^h37^m55^s
Dec: −63°49'

1-GHz flux/Jy: 2.1?
Spectral index: ?

Size/arcmin: 37×23
Type: S

Radio: Faint, incomplete shell, with possible extension to southwest.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×48'': $S=2.3$ Jy).

G302.3+0.7

RA: 12^h45^m55^s
Dec: -62°08'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 17
Type: S

Radio: Distorted shell, in complex region, with possibly associated filament.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 7.5$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 3.0$ Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 49'' : S = 3.2$ Jy).
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

G304.6+0.1

RA: 13^h05^m59^s
Dec: -62°42'

1-GHz flux/Jy: 14
Spectral index: 0.5

Kes 17

Size/arcmin: 8
Type: S

Radio: Incomplete shell.

X-ray: Detected.

Distance: Possible limit of > 9.7 kpc from H α absorption.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ($3'$) and Parkes 64-m at 5 GHz ($4'$).
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ($4.4' : S = 6.9$ Jy).
Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 48'' : S = 18$ Jy).
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
Combi *et al.* 2010, A&A, 523, A76. XMM-Newton observations

Lee *et al.* 2011, ApJ, 740, 31. Akari and Spitzer observations.
Wu *et al.* 2011, ApJ, 740, L12. Fermi observations.
Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
Gök & Sezer 2012, MNRAS, 423, 1215. Suzaku observations.
Gelfand *et al.* 2013, ApJ, 777, 148. ATCA at 1.4 GHz ($8'' \times 23'' : S = 10.9 \pm 0.4$ Jy), plus other observations.
Pannuti *et al.* 2014, AJ, 147, 55. ASCA and XMM-Newton observations.
Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.
Washino *et al.* 2016, PASJ, 68, S4. Suzaku observations.

G306.3-0.9

RA: 13^h21^m50^s
Dec: -63°34'

1-GHz flux/Jy: 0.16?
Spectral index: 0.5?

Size/arcmin: 4
Type: S?

Radio: Diffuse emission.

X-ray: Partial shell.

References:

Reynolds *et al.* 2013, ApJ, 766, 112. Chandra observations, and ATCA at 5.5 GHz ($23'' \times 26''$), and 9 GHz.
Combi *et al.* 2016, A&A, 592, A125. XMM-Newton and Chandra observations.

G308.1-0.7

RA: 13^h37^m37^s
Dec: -63°04'

1-GHz flux/Jy: 1.2?
Spectral index: ?

Size/arcmin: 13
Type: S

Radio: Faint shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 48'' : S = 1.3$ Jy).

G308.4–1.4

RA: 13^h41^m30^s
Dec: −63°44'

1-GHz flux/Jy: 0.4?
Spectral index: ?

Size/arcmin: 12×6?
Type: S?

W part has been called G308.3–1.4.

Radio: Complex structure, with multiple arcs.

X-ray: Limb brightened partial shell in W.

References:

Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT observations.
Prinz & Becker 2012, A&A, 544, A7. ATCA at 1.4 GHz
($53'' \times 64'' : S = 0.33$ Jy) and 2.5 GHz ($29'' \times 35'' : S = 0.24$ Jy),
plus Chandra and other observations.
Hui *et al.* 2012, ApJ, 750, 7. XMM-Newton and other observa-

tions.
De Horta *et al.* 2013, MNRAS, 428, 1980. ATCA at 1.4 GHz
($54'' \times 65''$) and 2.5 GHz ($29'' \times 35''$), plus other observations.
Green *et al.* 2014, PASA, 31, 42. Radio observations at
843 MHz ($45'' \times 50''$).

G308.8–0.1

RA: 13^h42^m30^s
Dec: −62°23'

1-GHz flux/Jy: 15?
Spectral index: 0.4?

Size/arcmin: 30×20?
Type: C?

Incorporates previous catalogued remnant G308.7+0.0.

Radio: Bright ridge in north, and arc to south.

Point sources: Pulsar near centre of remnant.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 16.7$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.0$ Jy).
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz
($4.4' : S = 6.5$ Jy).
Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz
($50''$).
Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.

Caswell *et al.* 1992, ApJ, 399, L151. MOST at 843 MHz
($43'' \times 49''$).
Kaspi *et al.* 1992, ApJ, 399, L155. Pulsar observations.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-
tions.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G309.2–0.6

RA: 13^h46^m31^s
Dec: −62°54'

1-GHz flux/Jy: 7?
Spectral index: 0.4?

Size/arcmin: 15×12
Type: S

Has been called G309.2–0.7.

Radio: Distorted shell.

X-ray: Extended emission, with unrelated central source.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 10.0$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 3.9$ Jy).
Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz
($1.5'$).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz
($44'' \times 50''$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
($43'' \times 48'' : S = 6$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
Gaensler *et al.* 1998, MNRAS, 299, 812. ATCA at 1.3 GHz
($24'' \times 23'' : S = 5.2 \pm 0.2$ Jy).
Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT obser-
vations.
Rakowski *et al.* 2006, ApJ, 649, L111. Observations of central
source.
Safi-Harb *et al.* 2007, ApJ, 659, 407. Observations of central
source.

G309.8+0.0

RA: 13^h50^m30^s
Dec: −62°05'

1-GHz flux/Jy: 17
Spectral index: 0.5

Size/arcmin: 25×19
Type: S

Radio: Distorted shell.

Point sources: Steep radio spectrum source near the centre of the remnant.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 26.4$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.4$ Jy).
Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz
($1'$).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
($43'' \times 49'' : S > 8.8$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-
tions.

G310.6–1.6

RA: 14^h00^m45^s
Dec: −63°26'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 2.5
Type: C?

X-ray: Bright central nebula, with faint shell.

Point sources: X-ray pulsar, and extended emission.

References:

Tomsick *et al.* 2009, ApJ, 701, 811. Chandra observations.
 Renaud *et al.* 2010, ApJ, 716, 663. Chandra and other observations.

G310.6–0.3

RA: 13^h58^m00^s
Dec: −62°09'

1-GHz flux/Jy: 5?
Spectral index: ?

Kes 20B
Size/arcmin: 8
Type: S

Radio: Asymmetric shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 49'': S=5.4 \text{ Jy})$.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G310.8–0.4

RA: 14^h00^m00^s
Dec: −62°17'

1-GHz flux/Jy: 6?
Spectral index: ?

Kes 20A
Size/arcmin: 12
Type: S

Radio: Arc in E, in complex region.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 49'': S=6.9 \text{ Jy})$.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.

G311.5–0.3

RA: 14^h05^m38^s
Dec: −61°58'

1-GHz flux/Jy: 3?
Spectral index: 0.5

Size/arcmin: 5
Type: S

Radio: Shell, not well resolved.

X-ray: Detected.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz
 $(3')$ and Parkes 64-m at 5 GHz $(4')$.
 Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at
 408 MHz $(3')$.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

$(43'' \times 49'': S=2.9 \text{ Jy})$.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Pannuti *et al.* 2014, AJ, 147, 55. ASCA observations.

G312.4–0.4

RA: 14^h13^m00^s
Dec: −61°44'

1-GHz flux/Jy: 45
Spectral index: 0.36

Size/arcmin: 38
Type: S

Radio: Irregular, incomplete shell.

X-ray: Weak emission in W.

Point sources: Nearby γ-ray sources and pulsars.

Distance: H_I absorption suggests >6 kpc and possibly >14 kpc.

References:

Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3': $S=56$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'' : $S > 19$), plus Parkes 64-m at 4.5 GHz ($S = 30 \pm 2$ Jy) and 8.55 GHz ($S=17 \pm 4$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Roberts *et al.* 1999, ApJ, 515, 712. MOST at 843 MHz (43''×49'').
 Case & Bhattacharya 1999, ApJ, 521, 246. Nearby γ-ray sources.
 Doherty *et al.* 2003, MNRAS, 339, 1048. ATCA at 1.4 GHz (25'') plus H_I absorption, and Chandra observations.

G312.5–3.0

RA: 14^h21^m00^s
Dec: −64°12'

1-GHz flux/Jy: 3.5?
Spectral index: ?

Size/arcmin: 20×18
Type: S

Radio: Distorted shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10'.4)
 Kane & Vaughan 2003, MNRAS, 344, 625. ATCA at 1.4 GHz (129''×116'') and 2.4 GHz (75''×67'').

G315.1+2.7

RA: 14^h24^m30^s
Dec: −57°50'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 190×150
Type: S

Radio: Poorly defined shell.

Optical: Filaments, brighter in NE.

References:

Duncan *et al.* 1997, MNRAS, 287, 722. Parkes 64-m at 2.4 GHz (11': $S=19 \pm 3$ Jy).
 Combi *et al.* 1998, A&A, 333, 298. Radio survey observations.

Stupar *et al.* 2007, MNRAS, 374, 1441. Optical and various radio observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. Hα observations.

G315.4–2.3		RCW 86, MSH 14–63
RA: 14 ^h 43 ^m 00 ^s	1-GHz flux/Jy: 49	Size/arcmin: 42
Dec: −62°30'	Spectral index: 0.6	Type: S
Possibly the remnant of the SN of AD185?		
Radio: Shell, brightest to the SW.		
Optical: Bright, radiative filaments, with some faint Balmer dominated filaments.		
X-ray: Partial shell, with thermal and non-thermal emission.		
Point sources: Several X-ray sources.		
Distance: Optical observations imply 2.3 kpc.		
References:		
van den Bergh <i>et al.</i> 1973, ApJS, 26, 19. Optical observations.	Rho <i>et al.</i> 2002, ApJ, 581, 1116. Chandra observations.	
Caswell <i>et al.</i> 1975, AuJPA, 37, 39. Molonglo at 408 MHz ($3' : S = 86$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 18.2$ Jy).	Gvaramadze & Vikhlinin 2003, A&A, 401, 625. Chandra point source search.	
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8.4'$) and 5 GHz ($4.4'$).	Sollerman <i>et al.</i> 2003, A&A, 407, 249. Optical spectroscopy.	
Winkler 1978, ApJ, 221, 220. X-ray detection.	Kaplan <i>et al.</i> 2004, ApJS, 153, 269. Chandra limits for any compact source.	
Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.	Bamba <i>et al.</i> 2005, ApJ, 621, 793. Chandra observations of rim.	
Pisarski <i>et al.</i> 1984, ApJ, 277, 710. Einstein observations	Vink <i>et al.</i> 2006, ApJ, 648, L33. Chandra and XMM-Newton observations.	
Nugent <i>et al.</i> 1984, ApJ, 284, 612. X-ray spectrum.	Ueno <i>et al.</i> 2007, PASJ, 59, S171. Suzaku observations.	
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 50''$).	Aharonian <i>et al.</i> 2009, ApJ, 692, 1500. γ -ray observations.	
Long & Blair 1990, ApJ, 358, L13. Balmer dominated optical filaments.	Yamaguchi <i>et al.</i> 2011, PASJ, 63, S837. Suzaku observations.	
Greidanus & Strom 1990, A&A, 240, 385. IRAS observations.	Helder <i>et al.</i> 2011, ApJ, 737, 85. XMM-Newton and optical observations.	
Kaastra <i>et al.</i> 1992, A&A, 264, 654. Ginga X-ray spectra.	Williams <i>et al.</i> 2011, ApJ, 741, 96. Spitzer observations.	
Strom 1994, MNRAS, 268, L5. Historical association	Mignani <i>et al.</i> 2012, MNRAS, 425, 2309. X-ray and optical observations of compact X-ray sources.	
Chin & Huang 1994, Nature, 371, 398. Questioning of historical association.	Lemoine-Goumard <i>et al.</i> 2012, A&A, 545, A28. γ -ray observations.	
Schaefer 1995, AJ, 110, 1793. Questioning of historical association.	Helder <i>et al.</i> 2013, MNRAS, 435, 910. Optical proper motion studies.	
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 48'' : S > 22$).	Castro <i>et al.</i> 2013, ApJ, 779, 49. Chandra of NW.	
Rosado <i>et al.</i> 1996, A&A, 315, 243. Optical kinematics.	Yuan <i>et al.</i> 2014, ApJ, 785, L22. γ -ray observations.	
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.	Tsubone <i>et al.</i> 2014, in EFXU, p72. Suzaku observations.	
Smith <i>et al.</i> 1997, AJ, 114, 2664. Observations of Balmer dominated filaments.	Broersen <i>et al.</i> 2014, MNRAS, 441, 3040. XMM-Newton observations.	
Vink <i>et al.</i> 1997, A&A, 328, 628. ASCA spectroscopy.	Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux densities at 30, 44 and 70 GHz.	
Bocchino <i>et al.</i> 2000, A&A, 360, 671. BeppoSAX observations of N and SW.	Ajello <i>et al.</i> 2016, ApJ, 819, 98. γ -ray observations, and ATCA H α of surroundings.	
Bamba <i>et al.</i> 2000, PASJ, 52, 1157. ASCA observations.	Yamaguchi <i>et al.</i> 2016, ApJ, 820, L3. Chandra observations of NE.	
Dickel <i>et al.</i> 2001, ApJ, 546, 447. ATCA at 1.34 GHz ($8''$), including polarisation.	Koo <i>et al.</i> 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).	
Ghavamian <i>et al.</i> 2001, ApJ, 547, 995. Optical spectroscopy.		
Borkowski <i>et al.</i> 2001, ApJ, 550, 334. ASCA observations.		

G315.4–0.3		Size/arcmin: 24×13
RA: 14 ^h 35 ^m 55 ^s	1-GHz flux/Jy: 8	Type: ?
Dec: −60°36'	Spectral index: 0.4	

Radio: Irregular non-thermal emission, with H α region superposed in E.

Optical: Detected.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 15.9$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 4.9$ Jy). Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
- Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz ($50'' : S = 6.25$ Jy), re-assessment of earlier flux densities. Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G315.9–0.0

RA: 14^h38^m25^s
Dec: −60°11'

1-GHz flux/Jy: 0.8?
Spectral index: ?

Size/arcmin: 25×14
Type: S

Has been called G315.8–0.0.

Radio: Faint, distorted shell, with elongated trail to pulsar.

Point sources: Pulsar at end of radio trail.

References:

- Kesteven *et al.* 1987, *AuJPh*, 40, 855. MOST at 843 MHz (44''×50'').
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×50'': $S=0.9$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m and ATCA OH

observations.
 Camilo *et al.* 2009, *ApJ*, 703, L55. Pulsar detection.
 Ng *et al.* 2012, *ApJ*, 746, 105. ATCA at 1.4, 2.5, 5 and 9 GHz, including polarisation, of pulsar trail.

G316.3–0.0

RA: 14^h41^m30^s
Dec: −60°00'

1-GHz flux/Jy: 20?
Spectral index: 0.4

(MSH 14–57)

Size/arcmin: 29×14
Type: S

Radio: Distorted shell, with possible ‘blowout’.

X-ray: Detected.

Distance: HI absorption data suggests > 7.2 kpc.

References:

- Shaver & Goss 1970, *AuJPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1975, *A&A*, 45, 239. Parkes HI absorption.
 Milne & Dickel 1975, *AuJPh*, 28, 209. Parkes 64-m at 5 GHz (4.4': $S=16.7$ Jy).
 Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44''×51'').

Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×50'': $S=20$ Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
 Acero *et al.* 2016, *ApJS*, 224, 8. γ-ray observations.

G317.3–0.2

RA: 14^h49^m40^s
Dec: −59°46'

1-GHz flux/Jy: 4.7?
Spectral index: ?

Size/arcmin: 11
Type: S

Radio: Incomplete shell.

Optical: Detected.

References:

- Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×50'': $S=5.2$ Jy).
 Stupar & Parker 2011, *MNRAS*, 414, 2282. Hα observations.

G318.2+0.1

RA: 14^h54^m50^s
Dec: −59°04'

1-GHz flux/Jy: >3.9?
Spectral index: ?

Size/arcmin: 40×35
Type: S

Radio: Faint shell, with central HII region.

X-ray: Sources within remnant.

References:

- Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×50'': $S>4.3$).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

Bocchino *et al.* 2001, *A&A*, 367, 629. BeppoSAX and ROSAT observations.
 Onako *et al.* 2016, *ApJ*, 829, 106. Akari observations.

G318.9+0.4

RA: 14^h58^m30^s
Dec: -58°29'

1-GHz flux/Jy: 4?
Spectral index: 0.2?

Size/arcmin: 30×14
Type: C

May not be a SNR?

Radio: Complex arcs, with off-centre core.

References:

- Whiteoak 1990, Nature, 347, 157. MOST at 843 MHz (43''×51'').
 Whiteoak 1993, ApJ, 415, 701. MOST at 843 MHz (43''×51'': $S=4.8\pm0.6$ Jy), Parkes 64-m at 4.5 GHz (4'7:S=3.7±0.2 Jy) and 8.4 GHz (2'8:S=3.0±0.4 Jy) including polarisation, and ATCA at 1.4 GHz and 4.8 GHz (11''×13'') of core.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'': S=4.8 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G320.4–1.2

RA: 15^h14^m30^s
Dec: -59°08'

1-GHz flux/Jy: 60?
Spectral index: 0.4

MSH 15–52, RCW 89

Size/arcmin: 35
Type: C

Has been suggested as the remnant of the SN of AD185?

Radio: Ragged shell.

Optical: RCW 89 is the H α emitting region to the NW.

X-ray: Partial shell, central nebula and pulsar and ‘jet’.

Point sources: Radio and X-ray pulsar, with wind nebula.

Distance: HI absorption indicates 5.2 kpc.

References:

- Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8/4) and 5 GHz (4/4).
 Dopita *et al.* 1977, ApJ, 214, 179. Optical spectra.
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'').
 Seward & Harnden 1982, ApJ, 256, L45. X-ray observations of pulsar.
 Manchester *et al.* 1982, ApJ, 262, L31. Radio observations of pulsar.
 Seward *et al.* 1983, ApJ, 267, 698. X-ray, Optical and IR.
 van den Bergh & Kamper 1984, ApJ, 280, L51. Optical expansion.
 Seward *et al.* 1984, ApJ, 281, 650. X-ray observations of pulsar and nebulosity.
 Lorret *et al.* 1987, A&A, 180, 65. Optical observations.
 Trussoni *et al.* 1990, A&A, 234, 403. EXOSAT observations.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Arendt 1991, AJ, 101, 2160. IRAS observations, including compact source.
 Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz (4'5:S=37±7 Jy) and 8.4 GHz (3'0:S=24±4 Jy), including polarisation and review of flux densities.
 Strom 1994, MNRAS, 268, L5. Historical association.
 Chin & Huang 1994, Nature, 371, 398. Questioning of historical association.
 Matz *et al.* 1994, ApJ, 434, 288. X-ray observations of pulsar.
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.
 Du Plessis *et al.* 1995, ApJ, 453, 746. Hartesbeesthoek 26-m at 2.3, 5 and 8.5 GHz (S=42,35.6 and 14.5 Jy).
 Greiveldinger *et al.* 1995, ApJ, 454, 855. ROSAT observations.
 Trussoni *et al.* 1996, A&A, 306, 581. ROSAT observations.
 Tamura *et al.* 1996, PASJ, 48, L33. ASCA observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'': S=62 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Marsden *et al.* 1997, ApJ, 491, L39. X-ray spectroscopy.
 Brazier & Becker 1997, MNRAS, 284, 335. ROSAT observations.
 Gaensler *et al.* 1999, MNRAS, 305, 724. ATCA at 1.4 GHz (24''×21''), plus HI observations, and 5.3 GHz (15''×10'').
 Sako *et al.* 2000, ApJ, 537, 422. Possible high energy γ-ray detection of pulsar.
 Mineo *et al.* 2001, A&A, 380, 695. BeppoSAX observations.
 Gaensler *et al.* 2002, ApJ, 569, 878. Chandra observations of pulsar and nebula.
 Dubner *et al.* 2002, AJ, 123, 337. ATCA at 1.4 GHz (4'0×2'7), plus HI observations.
 Aharonian *et al.* 2005, A&A, 435, L17. γ-ray detection.
 Yatsu *et al.* 2005, ApJ, 631, 312. Chandra observations of pulsar and jet.
 DeLaney *et al.* 2006, ApJ, 640, 929. Chandra and ROSAT multi-epoch observations of pulsar wind nebula.
 Kaplan & Moon 2006, ApJ, 644, 1056. Possible IR detection of pulsar.
 Forot *et al.* 2006, ApJ, 651, L45. X-ray observations of pulsar and nebula.
 Nakamori *et al.* 2008, ApJ, 677, 297. γ-ray observations.
 Yatsu *et al.* 2009, PASJ, 61, 129. Chandra observations of pulsar wind nebula.
 Koo *et al.* 2011, ApJ, 732, 6. Akari and Spitzer observations.
 An *et al.* 2014, ApJ, 793, 90. NuSTAR observations of PWN.

G320.6–1.6

RA: $15^{\text{h}}17^{\text{m}}50^{\text{s}}$
Dec: $-59^{\circ}16'$

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 60×30
Type: S

Radio: Faint shell, overlapping G320.4–1.2 in W.

Optical: Detected.

References:

Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz ($4'5$) and 8.4 GHz ($3'0$), including polarisation.
Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz ($10'4$).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S > 9.3$).
Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G321.9–1.1

RA: $15^{\text{h}}23^{\text{m}}45^{\text{s}}$
Dec: $-58^{\circ}13'$

1-GHz flux/Jy: >3.4 ?
Spectral index: ?

Size/arcmin: 28
Type: S

Radio: Faint shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 51'' : S > 3.8$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G321.9–0.3

RA: $15^{\text{h}}20^{\text{m}}40^{\text{s}}$
Dec: $-57^{\circ}34'$

1-GHz flux/Jy: 13
Spectral index: 0.3

Size/arcmin: 31×23
Type: S

Radio: Shell brighter to the W, with Cir X-1 to N.

Point sources: Compact, probably thermal source at S edge.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 18.3$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.8$ Jy).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 52''$).
Stewart *et al.* 1993, MNRAS, 261, 593. ATCA at 1.5 GHz ($21''$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

($43'' \times 51'' : S > 8.3$).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Mignani *et al.* 2002, A&A, 386, 487. HST proper motion study of Cir X-1.
Tudose *et al.* 2006, MNRAS, 372, 417. MOST at 843 MHz.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G322.1+0.0

RA: $15^{\text{h}}20^{\text{m}}49^{\text{s}}$
Dec: $-57^{\circ}10'$

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 8×4.5 ?
Type: S?

Radio: Circular shell, with extension to S.

X-ray: Diffuse emission.

Point sources: Cir X-1 HMXB at centre.

References:

Heinz *et al.* 2013, ApJ, 779, 171. Chandra observations and ATCA at 1.1–3.1 GHz ($4''0 \times 4''9$).

G322.5–0.1

RA: $15^{\text{h}}23^{\text{m}}23^{\text{s}}$
Dec: $-57^{\circ}06'$

1-GHz flux/Jy: 1.5
Spectral index: 0.4

Size/arcmin: 15
Type: C

Radio: Shell with central extended source.

Point sources: PN Pe 2-8 within boundary.

References:

Whiteoak 1992, MNRAS, 256, 121. MOST at 843 MHz
 $(43'' \times 51' : S = 2.0 \pm 0.3 \text{ Jy})$.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 51' : S = 1.7 \text{ Jy})$, plus Parkes 64-m at 4.5 GHz

$(=0.89 \pm 0.13)$.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G323.5+0.1

RA: $15^{\text{h}}28^{\text{m}}42^{\text{s}}$
Dec: $-56^{\circ}21'$

1-GHz flux/Jy: 3?
Spectral index: 0.4?

Size/arcmin: 13
Type: S

Radio: Distorted shell, confused with thermal emission.

Point sources: Compact, probably thermal source near centre.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 4.2 \text{ Jy}$) and Parkes 64-m at 5 GHz ($4' : S = 1.5 \text{ Jy}$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 52' : S = 4.2 \text{ Jy})$.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G326.3–1.8

RA: $15^{\text{h}}53^{\text{m}}00^{\text{s}}$
Dec: $-56^{\circ}10'$

1-GHz flux/Jy: 145
Spectral index: varies

MSH 15–56

Size/arcmin: 38
Type: C

Has been called G326.2–1.7.

Radio: Shell, with elongated, flat-spectrum core.

Optical: Emission around the shell.

X-ray: Shell, with central extended emission.

Point sources: Compact X-ray source.

References:

Clark *et al.* 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 180 \text{ Jy}$).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.
 Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz
 $(0.8' : S > 95)$ and Parkes 64-m at 14.7 GHz ($2.2' : S = 69 \pm 8 \text{ Jy}$).
 van den Bergh 1979, ApJ, 227, 497. Optical observations.
 Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.
 Dennefeld 1980, PASP, 92, 603. Optical spectra.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz
 $(44'' \times 53'')$.
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz
 $(43'' \times 52' : S = 153 \pm 40 \text{ Jy})$, and Parkes 64-m at 8.4 GHz
 $(3' : S = 68 \pm 5 \text{ Jy})$, including polarisation.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Kassim *et al.* 1993, ApJ, 419, 733. ROSAT image.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 52' : S > 130)$.
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Dickel *et al.* 2000, ApJ, 543, 840. ATCA at 1.34 GHz
 $(8.5'' \times 6.4' : S > 60 \text{ Jy})$, 4.80 GHz ($3''/8 : S = 25 \text{ Jy}$ for core
 only) and 8.64 GHz ($3''/2 : S = 15 \text{ Jy}$ for core only).
 Temiim *et al.* 2013, ApJ, 768, 61. XMM-Newton and Chandra
 observations.
 Yatsu *et al.* 2013, ApJ, 773, 25. XMM-Newton and Chandra
 observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 5
 frequencies between 30 and 143 GHz.
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G327.1–1.1

RA: 15^h54^m25^s
Dec: −55°09'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 18
Type: C

Radio: Shell, with off-centre core.

X-ray: Diffuse, with core.

References:

- Milne & Dickel 1974, *AuJPh*, 27, 549. Parkes 64-m at 2.7 GHz (8' : $S=10\pm15$ Jy).
Clark *et al.* 1975, *AuJPA*, 37, 1. Molonglo at 408 MHz (3' : $S=10.6$ Jy) and Parkes 64-m at 5 GHz (4' : $S=4.3$ Jy).
Lamb & Markert 1981, *ApJ*, 244, 94. Einstein observations.
Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 52'' : $S=7.6$ Jy).
Seward *et al.* 1996, *ApJ*, 471, 887. ROSAT observations.
Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
Sun *et al.* 1999, *ApJ*, 511, 274. ASCA and ROSAT observa-

- tions.
Bocchino & Bandiera 2003, *A&A*, 398, 195. BeppoSAX observations.
Temim *et al.* 2009, *ApJ*, 691, 895. Chandra and XMM-Newton observations.
Acero *et al.* 2011, *ICRC*, 7, 185. γ -ray observations.
Temim *et al.* 2015, *ApJ*, 808, 100. Chandra observations.
Ma *et al.* 2016, *ApJ*, 820, 100. ATCA at 1.4 (19'' × 23''), 2.3 (11'' × 16''), 4.8 (13'' × 15'') and 8.6 GHz (8'' × 10'') of core, including polarisation.

G327.2–0.1

RA: 15^h50^m55^s
Dec: −54°18'

1-GHz flux/Jy: 0.4
Spectral index: ?

Size/arcmin: 5
Type: S

Has been called G327.24–0.13.

Radio: Shell, possibly with central emission.

Point sources: Central pulsar (magnetar).

References:

- Camilo *et al.* 2007, *ApJ*, 666, L93. Pulsar observations.
Gelfand & Gaensler 2007, *ApJ*, 667, 1111. Chandra and XMM-Newton observations, and 843 GHz (43'' × 53'' : $S=0.5\pm0.1$ Jy) and SGPS at 1.4 GHz ($S=0.3\pm0.1$ Jy).

G327.4+0.4

RA: 15^h48^m20^s
Dec: −53°49'

1-GHz flux/Jy: 30?
Spectral index: 0.6

Kes 27

Size/arcmin: 21
Type: S

Has been called G327.3+0.4 and G327.3+0.5.

Radio: Incomplete, multi-arc shell, brightest to the SE.

X-ray: Diffuse, brighter in E.

Distance: H_I absorption indicates 4.3 to 5.4 kpc.

References:

- Caswell *et al.* 1975, *AuJPA*, 37, 39. Molonglo at 408 MHz (3' : $S=58$ Jy) and Parkes 64-m at 5 GHz (4' : $S=12.4$ Jy).
Dickel & Milne 1976, *AuJPh*, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8.4') and 5 GHz (4.4').
Lamb & Markert 1981, *ApJ*, 244, 94. Einstein observations.
Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44'' × 55'').
Milne *et al.* 1989, *PASA*, 8, 187. MOST at 843 MHz (43'' × 53'' : $S=32.2\pm6$ Jy), and Parkes 64-m at 8.4 GHz (3' : $S=9.4\pm0.8$ Jy), including polarisation.
Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43'' × 53'' : $S=25$ Jy).

- Seward *et al.* 1996, *ApJ*, 471, 887. ROSAT observations.
Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m and ATCA OH observations.
McClure-Griffiths *et al.* 2001, *ApJ*, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'.0 × 1'.8), plus H_I.
Enoguchi *et al.* 2002, *PASJ*, 54, 229. ASCA observations.
Kawasaki *et al.* 2005, *ApJ*, 631, 935. ASCA observations.
Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer possible detection.
Chen *et al.* 2008, *ApJ*, 676, 1040. Chandra observations.
Minami *et al.* 2014, in EFXU, p48. Suzaku observations.
Xing *et al.* 2015, *ApJ*, 805, 19. γ -ray observations.

G327.4+1.0

RA: 15^h46^m48^s
Dec: -53°20'

1-GHz flux/Jy: 1.9?
Spectral index: ?

Size/arcmin: 14
Type: S

Radio: Asymmetric shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

(43''×54'': $S=2.1$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'0×1'8), plus H α .

G327.6+14.6		SN1006, PKS 1459–41
RA: 15 ^h 02 ^m 50 ^s	1-GHz flux/Jy: 19	Size/arcmin: 30
Dec: −41°56'	Spectral index: 0.6	Type: S
This is the remnant of the SN of AD1006.		
Radio: Shell, with two bright arcs.		
Optical: Filaments to the NW, with broad H α component.		
X-ray: Thermal shell, with non-thermal limb-brightened arcs.		
Point sources: The background Schweizer–Middleditch star is near the middle of the remnant.		
Distance: Optical spectra and proper motion indicate 2.2 kpc.		
References:		
van den Bergh 1976, ApJ, 208, L17. Optical observations.	Sollerman <i>et al.</i> 2003, A&A, 407, 249. Optical spectroscopy.	
Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).	Winkler <i>et al.</i> 2003, ApJ, 585, 324. Optical proper motion studies.	
Becker <i>et al.</i> 1980, ApJ, 240, L33. X-ray spectroscopy.	Vink <i>et al.</i> 2003, ApJ, 587, L31. XMM-Newton observations.	
Schweizer & Middleditch 1980, ApJ, 241, 1039. Possible stellar remnant.	Bamba <i>et al.</i> 2003, ApJ, 589, 827. Chandra observations of NE.	
Pye <i>et al.</i> 1981, MNRAS, 194, 569. Einstein observations.	Korreck <i>et al.</i> 2004, ApJ, 615, 280. Far-uv observations.	
Caswell <i>et al.</i> 1983, MNRAS, 204, 921. FIRST at 1415 MHz (77").	Aharonian <i>et al.</i> 2005, A&A, 437, 135. γ -ray limit.	
Reynolds & Gilmore 1986, AJ, 92, 1138. VLA at 1.37 and 1.67 GHz (16''. \times 20'').	Winkler <i>et al.</i> 2005, ApJ, 624, 189. HST absorption towards background sources.	
Kirshner <i>et al.</i> 1987, ApJ, 315, L135. Broad H α optical component.	Kalemci <i>et al.</i> 2006, ApJ, 640, L55. γ -ray limit upper limit.	
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''. \times 66'').	Kalemci <i>et al.</i> 2006, ApJ, 644, 274. INTEGRAL observations.	
Fesen <i>et al.</i> 1988, ApJ, 327, 164. UV absorption spectra of the Schweizer–Middleditch star.	Acero <i>et al.</i> 2007, A&A, 475, 883. XMM-Newton observations.	
Roger <i>et al.</i> 1988, ApJ, 332, 940. MOST at 843 MHz (44''. \times 66''). $S=17.5\pm1.5$ Jy.	Raymond <i>et al.</i> 2007, ApJ, 659, 1257. HST H α observations.	
Long <i>et al.</i> 1988, ApJ, 333, 749. Optical proper motion for distance.	Hamilton <i>et al.</i> 2007, MNRAS, 381, 771. UV spectroscopy of the Schweizer–Middleditch star.	
Jones & Pye 1989, MNRAS, 238, 567. EXOSAT observations.	Bamba <i>et al.</i> 2008, AdSpR, 41, 411. Suzaku observations.	
Leahy <i>et al.</i> 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.	Cassam-Chenaï <i>et al.</i> 2008, ApJ, 680, 1180. ATCA and VLA at 1.5 GHz (6''. \times 9'') plus Chandra observations.	
Wu <i>et al.</i> 1993, ApJ, 416, 247. UV spectra.	Yamaguchi <i>et al.</i> 2008, PASJ, 60, S141. Suzaku observations.	
Reynolds & Gilmore 1993, AJ, 106, 272. VLA at 1.37 and 1.67 GHz (24''), including polarisation.	Bamba 2008, PASJ, 60, S153. Suzaku observations.	
Moffett <i>et al.</i> 1993, AJ, 106, 1566. VLA at 1.37 and 1.67 GHz (10''. \times 14''), for proper motion studies.	Dyer <i>et al.</i> 2009, AJ, 137, 2956. GBT and VLA observations at 1.4 GHz.	
Ozaki <i>et al.</i> 1994, PASJ, 46, 367. X-ray observations.	Winkler <i>et al.</i> 2011, ApJ, 742, 80. Multi-epoch UV spectroscopy of the Schweizer–Middleditch star.	
Raymond <i>et al.</i> 1995, ApJ, 454, L31. Far UV spectroscopy.	Miceli <i>et al.</i> 2012, A&A, 546, A66. XMM-Newton observations.	
Koyama <i>et al.</i> 1995, Nature, 378, 255. ASCA observations.	Miceli <i>et al.</i> 2013, A&A, 550, A80. XMM-Newton observations.	
Willingale <i>et al.</i> 1996, MNRAS, 278, 749. ROSAT observations.	Broersen <i>et al.</i> 2013, A&A, 552, A9. XMM-Newton observations of NW knot.	
Blair <i>et al.</i> 1996, ApJ, 468, 871. Optical absorption studies.	Katsuda <i>et al.</i> 2013, ApJ, 763, 85. Chandra proper motion study of NW.	
Laming <i>et al.</i> 1996, ApJ, 472, 267. Modelling of optical spectra, including distance.	Winkler <i>et al.</i> 2013, ApJ, 764, 156. Spitzer observations.	
Wu <i>et al.</i> 1997, ApJ, 477, L53. Far UV absorption spectra of the Schweizer–Middleditch star.	Uchida <i>et al.</i> 2013, ApJ, 771, 56. Suzaku observations.	
Winkler & Long 1997, ApJ, 486, L137. UV absorption spectra of background quasar.	Reynoso <i>et al.</i> 2013, AJ, 145, 104. VLA and ATCA at 1.4 GHz (10''), including polarisation.	
Winkler & Long 1997, ApJ, 491, 829. ROSAT and optical images.	Nikolić <i>et al.</i> 2013, Science, 340, 45. Optical spectroscopy.	
Tanimori <i>et al.</i> 1998, ApJ, 497, L25. γ -ray detection.	Winkler <i>et al.</i> 2014, ApJ, 781, 65. Chandra observations and H α spectroscopy.	
Vink <i>et al.</i> 2000, A&A, 354, 931. X-ray spectroscopy.	Miceli <i>et al.</i> 2014, ApJ, 782, L33. XMM-Newton observations of SW.	
Burleigh <i>et al.</i> 2000, A&A, 356, 585. Optical spectroscopy of the Schweizer–Middleditch star.	Acero <i>et al.</i> 2015, A&A, 580, A74. γ -ray limit.	
Allen <i>et al.</i> 2001, ApJ, 558, 739. ASCA, ROSAT and other X-ray observations.	Sparls <i>et al.</i> 2015, ApJ, 815, L9. Polarised optical line observations.	
Dubner <i>et al.</i> 2002, A&A, 387, 1047. ATCA at 1.4 GHz (4'.7 \times 3'.0) for H α , plus CO observations.	Li <i>et al.</i> 2015, MNRAS, 453, 3953. XMM-Newton observations.	
Ghavamian <i>et al.</i> 2002, ApJ, 572, 888. Optical of filaments in NW.	Arnaud <i>et al.</i> 2016, A&A, 586, A134. Planck flux densities at 4 frequencies between 30 and 100 GHz.	
	Xing <i>et al.</i> 2016, ApJ, 823, 44. γ -ray observations.	
	Li <i>et al.</i> 2016, MNRAS, 462, 158. XMM-Newton observations.	

G328.4+0.2		(MSH 15–57)
RA: 15 ^h 55 ^m 30 ^s Dec: −53°17'	1-GHz flux/Jy: 15 Spectral index: 0.0	Size/arcmin: 5 Type: F

Radio: Amorphous emission, with central bar.

X-ray: Detected at high energies.

Distance: HI absorption indicates > 17.4 kpc.

References:

- Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50'').
 Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×54'' : $S = 15$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Gaensler *et al.* 2000, ApJ, 542, 380. ATCA at 1.4 GHz

- (16''5×19''5 : $S = 14.3 \pm 0.1$ Jy) and 4.5 GHz (2''0×1''5 : $S = 12.5 \pm 0.2$ Jy).
 Hughes *et al.* 2000, ApJ, 542, 386. ASCA observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1'8×2'0), plus HI.
 Johnston *et al.* 2004, MNRAS, 348, L19. ATCA at 19 GHz (6''1×7''7), including polarisation.
 Gelfand *et al.* 2007, ApJ, 663, 468. ATCA at 1.4 GHz (5''8×7''0 : $S = 13.8 \pm 0.4$ Jy), plus XMM-Newton observations.

G329.7+0.4

RA: 16 ^h 01 ^m 20 ^s Dec: −52°18'	1-GHz flux/Jy: >34? Spectral index: ?	Size/arcmin: 40×33 Type: S
---	--	---

Radio: Diffuse shell, in complex region.

References:

- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×53'' : $S > 38$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

- McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (2'0×1'8), plus HI.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G330.0+15.0

Lupus Loop

RA: 15 ^h 10 ^m 00 ^s Dec: −40°00'	1-GHz flux/Jy: 350? Spectral index: 0.5?	Size/arcmin: 180? Type: S
---	---	--

Radio: Low surface brightness loop with HI shell.

X-ray: Detected, with central source.

Point sources: Central, possibly pulsating, X-ray source.

References:

- Milne 1971, AuJPh, 24, 757. Parkes 64-m at 408 MHz (48'), 635 MHz (31') and 1410 MHz (15').
 Milne & Dickel 1974, AuJPh, 27, 549. Parkes 64-m at 2.7 GHz (8'4 : $S = 120 \pm 30\%$ Jy).
 Toor 1980, A&A, 85, 184. X-ray image and spectrum.
 Colomb & Dubner 1982, A&A, 112, 141. Argentine 30-m dish

- at 1.42 GHz (30'), HI observations.
 Leahy *et al.* 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.
 Ozaki *et al.* 1994, PASJ, 46, 367. X-ray observations.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Shinn *et al.* 2006, ApJ, 644, L189. Far UV observations.

G330.2+1.0

RA: $16^{\text{h}}01^{\text{m}}06^{\text{s}}$
Dec: $-51^{\circ}34'$

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 11
Type: S?

Radio: Clumpy non-thermal emission, possibly a distorted shell.

X-ray: Shell.

Point sources: Central compact X-ray source.

Distance: H α absorption indicates >4.9 kpc.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 8.6$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 4.0$ Jy).
Caswell *et al.* 1983, MNRAS, 204, 915. FIRST at 1415 MHz ($52'' \times 47''$), and MOST at 843 MHz ($43'' \times 55''$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 55'' : S = 4.7$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ($2'.0 \times 1'.8$), plus H α .
Torii *et al.* 2006, PASJ, 58, L11. ASCA detection.
Park *et al.* 2009, ApJ, 695, 431. XMM-Newton and Chandra observations.
Abramowski *et al.* 2014, MNRAS, 441, 790. γ -ray upper limit.

G332.0+0.2

RA: $16^{\text{h}}13^{\text{m}}17^{\text{s}}$
Dec: $-50^{\circ}53'$

1-GHz flux/Jy: 8?
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Incomplete shell.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ($3'$) and Parkes 64-m at 5 GHz ($4'$).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 57''$).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 55'' : S = 8.9$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ($2'.0 \times 1'.8$), plus H α .
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G332.4–0.4		RCW 103
RA: $16^{\text{h}}17^{\text{m}}33^{\text{s}}$	1-GHz flux/Jy: 28	Size/arcmin: 10
Dec: $-51^{\circ}02'$	Spectral index: 0.5	Type: S
Radio: Shell, brightest to the S.		
Optical: Filaments correspond well to the radio shell, brightest in SE.		
X-ray: Brightest to NW, with point source near centre.		
Point sources: Central, variable X-ray source, and nearby pulsar.		
Distance: H α absorption indicates 3.1 kpc.		
References:		
Shaver & Goss 1970, <i>AuJPA</i> , 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').	Kaspi <i>et al.</i> 1998, <i>ApJ</i> , 503, L161. Pulsar observations.	
Caswell <i>et al.</i> 1975, <i>A&A</i> , 45, 239. Parkes H α absorption.	Gotthelf <i>et al.</i> 1999, <i>ApJ</i> , 514, L107. X-ray variability of central source.	
Tuohy & Garmire 1980, <i>ApJ</i> , 239, L107. Einstein detection of X-ray point source.	Oliva <i>et al.</i> 1999, <i>A&A</i> , 343, 943. ISO spectroscopy.	
Caswell <i>et al.</i> 1980, <i>MNRAS</i> , 190, 881. FIRST at 1415 MHz (50'').	Torii <i>et al.</i> 2000, <i>ApJ</i> , 534, L71. X-ray timing on pulsar, including glitch.	
Lamb & Markert 1981, <i>ApJ</i> , 244, 94. Einstein observations.	Reynoso <i>et al.</i> 2004, <i>PASA</i> , 21, 82. ATCA at 1.4 GHz (50''), including H α absorption to central source.	
Ruiz 1983, <i>AJ</i> , 88, 1210. Optical spectra.	Russeil <i>et al.</i> 2005, <i>A&A</i> , 429, 497. H α observations.	
Tuohy <i>et al.</i> 1983, <i>ApJ</i> , 268, 778. X-ray observations.	Reach <i>et al.</i> 2006, <i>AJ</i> , 131, 1479. Spitzer observations.	
Leibowitz & Danziger 1983, <i>MNRAS</i> , 204, 273. Optical spectra.	Paron 2006, <i>PASA</i> , 23, 69. CO and HCO $^{+}$ observations of surroundings.	
Nugent <i>et al.</i> 1984, <i>ApJ</i> , 284, 612. X-ray spectrum (and Einstein image from Tuohy, private communication).	De Luca 2006, <i>Science</i> , 313, 814. XMM-Newton observations of central source.	
Meaburn & Allen 1986, <i>MNRAS</i> , 222, 593. Optical spectra.	De Luca 2007, <i>Ap&SS</i> , 308, 231. XMM-Newton observation of periodicity of central source.	
Kesteven & Caswell 1987, <i>A&A</i> , 183, 118. MOST at 843 MHz (44'' \times 57'').	Matsumoto <i>et al.</i> 2007, <i>PASJ</i> , 59, S199. Suzaku observations.	
Oliva <i>et al.</i> 1990, <i>A&A</i> , 240, 453. IR spectroscopy.	De Luca 2008, <i>ApJ</i> , 682, 1185. IR observations of central source.	
Dickel <i>et al.</i> 1996, <i>AJ</i> , 111, 340. ATCA at 1.36 (8'') and 2.37 GHz (4.5''), including polarisation.	Anderson <i>et al.</i> 2011, <i>ApJ</i> , 742, 7. Spitzer observations.	
Whiteoak & Green 1996, <i>A&AS</i> , 118, 329. MOST at 843 MHz (43'' \times 55'': S=34 Jy).	Xing <i>et al.</i> 2014, <i>ApJ</i> , 781, 64. γ -ray observations.	
Fral <i>et al.</i> 1996, <i>AJ</i> , 111, 1651. OH emission near remnant.	Frank <i>et al.</i> 2015, <i>ApJ</i> , 810, 113. Chandra observations.	
Gotthelf <i>et al.</i> 1997, <i>ApJ</i> , 487, L175. ASCA observations of compact X-ray source.	Koo <i>et al.</i> 2016, <i>ApJ</i> , 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).	
Carter <i>et al.</i> 1997, <i>PASP</i> , 109, 990. Optical expansion.	Rea <i>et al.</i> 2016, <i>ApJ</i> , 828, L13. Chandra, NuSTAR and Swift observations.	
Torii <i>et al.</i> 1998, <i>ApJ</i> , 494, L207. ASCA detection of nearby pulsar.		

G332.4+0.1		MSH 16–51, Kes 32
RA: $16^{\text{h}}15^{\text{m}}20^{\text{s}}$	1-GHz flux/Jy: 26	Size/arcmin: 15
Dec: $-50^{\circ}42'$	Spectral index: 0.5	Type: S
Has been called G332.4+0.2.		
Radio: Distorted shell, with thermal jet and plume adjacent.		
Optical: Detected.		
X-ray: Shell, brightest to NW.		
Point sources: Pulsar nearby.		
References:		
Shaver & Goss 1970, <i>AuJPA</i> , 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').	Fral <i>et al.</i> 1996, <i>AJ</i> , 111, 1651. OH emission near remnant.	
Roger <i>et al.</i> 1985, <i>Nature</i> , 316, 44. MOST at 843 MHz (44'' \times 57'').	Brinkmann <i>et al.</i> 1999, <i>A&A</i> , 346, 599. ROSAT image of remnant and nearby pulsar.	
Caraveo 1993, <i>ApJ</i> , 415, L111. Nearby pulsar.	Vink <i>et al.</i> 2004, <i>ApJ</i> , 604, 693. Chandra observations.	
Whiteoak & Green 1996, <i>A&AS</i> , 118, 329. MOST at 843 MHz (43'' \times 56'': S=29 Jy).	Reach <i>et al.</i> 2006, <i>AJ</i> , 131, 1479. Spitzer possible detection.	
	Stupar & Parker 2011, <i>MNRAS</i> , 414, 2282. H α observations.	

G332.5–5.6

RA: 16^h43^m20^s
Dec: −54°30'

1-GHz flux/Jy: 2?
Spectral index: 0.7?

Size/arcmin: 35
Type: S

Radio: Bipolar shell, with central emission also.

Optical: Patchy filaments.

X-ray: Emission from centre.

Distance: Optical extinction suggests 3 kpc.

References:

Reynoso & Green 2007, MNRAS, 375, 92. ATCA at 1.4 GHz (40'' : $S = 1.90 \pm 0.15$ Jy) and 2.4 GHz (90'' : $S = 1.3 \pm 0.2$ Jy) including polarisation, and 843 MHz (43'' × 53'') plus ROSAT observations.
 Stupar *et al.* 2007, MNRAS, 381, 377. Optical observations, plus ATCA at 1.4 GHz (58'') and 2.4 GHz (95'') including

polarisation, and other observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Zhu *et al.* 2015, MNRAS, 452, 3470. Suzaku observations, and distance estimate.
 Suárez *et al.* 2015, A&A, 583, A84. XMM-Newton observations of central region.

G335.2+0.1

RA: 16^h27^m45^s
Dec: −48°47'

1-GHz flux/Jy: 16
Spectral index: 0.5

Size/arcmin: 21
Type: S

Radio: Well defined shell.

Point sources: Old pulsar within remnant boundary.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 27.1$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 8.6$ Jy).
 Kaspi *et al.* 1996, AJ, 111, 2028. Pulsar observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

(43'' × 57'' : $S = 16$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G336.7+0.5

RA: 16^h32^m11^s
Dec: −47°19'

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 14 × 10
Type: S

Radio: Irregular shell.

Optical: Detected.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' × 60'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

(43'' × 58'' : $S = 6.1$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G337.0–0.1

RA: 16^h35^m57^s
Dec: −47°36'

1-GHz flux/Jy: 1.5
Spectral index: 0.6?

Size/arcmin: 1.5
Type: S

(CTB 33)

This entry refers to a small (1.5') SNR, not the larger previously catalogued G337.0–0.1. Has mistakenly been called G337.7–0.1.

Radio: Shell, in a complex region.

Point sources: Associated with a soft gamma repeater.

Distance: Association with CTB 33 gives 11 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' × 58'' : $S = 21$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Sarma *et al.* 1997, ApJ, 483, 335. ATCA at 1.4 GHz (12'') including H α , and 8.9 GHz (13'' × 15'') for recombination lines,

clarifying extent of the remnant.
 Woods *et al.* 1999, ApJ, 519, L139. Soft gamma repeater observations.
 Hurley *et al.* 2000, ApJ, 528, L21. ASCA observations of soft gamma repeater.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Castro *et al.* 2013, ApJ, 774, 36. Fermi observations.

G337.2–0.7

RA: $16^{\text{h}}39^{\text{m}}28^{\text{s}}$
Dec: $-47^{\circ}51'$

1-GHz flux/Jy: 1.5
Spectral index: 0.4

Size/arcmin: 6
Type: S

Radio: Shell, brighter in S.

X-ray: Extended emission.

Distance: HI absorption suggests 2.0 to 9.3 kpc.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 3.8$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 0.70$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 58'' : S = 2.0$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT obser-

vations.
 Rakowski *et al.* 2006, ApJ, 646, 982. ATCA observations at 1.3 GHz ($15'' : S = 1.55 \pm 0.05$ Jy) and 5 GHz ($15'' : S = 0.93 \pm 0.02$ Jy), plus HI and Chandra observations.
 Yamaguchi *et al.* 2014, ApJ, 785, L27. Suzaku observations.
 Takata *et al.* 2016, PASJ, 68, S3. Suzaku observations.

G337.2+0.1

RA: $16^{\text{h}}35^{\text{m}}55^{\text{s}}$
Dec: $-47^{\circ}20'$

1-GHz flux/Jy: 1.5?
Spectral index: ?

Size/arcmin: 3×2
Type: ?

Radio: Not well defined.

X-ray: Detected.

Distance: Association with HI hole gives 14 kpc.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59'' : S = 1.6$ Jy)
 Combi *et al.* 2005, A&A, 431, L9. ASCA and other observations.

Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations of nearby source.
 Combi *et al.* 2006, ApJ, 653, L41. XMM-Newton observations.

G337.3+1.0

RA: $16^{\text{h}}32^{\text{m}}39^{\text{s}}$
Dec: $-46^{\circ}36'$

1-GHz flux/Jy: 16
Spectral index: 0.55

Kes 40

Size/arcmin: 15×12
Type: S

Radio: Nearly complete shell.

References:

Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ($3' : S = 24.6$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.2$ Jy).
 Dickel & Milne 1976, AuJPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8/4) and 5 GHz (4/4).
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 61''$).
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz

($43'' \times 59'' : S = 14.8 \pm 3.0$ Jy), and Parkes 64-m at 8.4 GHz ($3' : S = 5.1 \pm 0.6$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59'' : S = 20$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G337.8–0.1

RA: $16^{\text{h}}39^{\text{m}}01^{\text{s}}$
Dec: $-46^{\circ}59'$

1-GHz flux/Jy: 18
Spectral index: 0.5

Kes 41

Size/arcmin: 9×6
Type: S

Radio: Distorted shell.

X-ray: Centrally brightened.

Distance: HI absorption suggests 11 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz ($3'$) and Parkes 64-m at 5 GHz ($4'$).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59'' : S = 18$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
 Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz, for associated OH masers.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including HI.
 Combi *et al.* 2008, A&A, 488, L25. XMM-Newton observations.
 Zhang *et al.* 2015, ApJ, 799, 103. XMM-Newton, Chandra and CO observations.
 Liu *et al.* 2015, ApJ, 809, 102. γ -ray observations.
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.
 Qiao *et al.* 2016, ApJS, 227, 26. OH maser observations.

G338.1+0.4

RA: $16^{\text{h}}37^{\text{m}}59^{\text{s}}$
Dec: $-46^{\circ}24'$

1-GHz flux/Jy: 4?
Spectral index: 0.4

Size/arcmin: 15?
Type: S

Radio: Arc in NE, merging with thermal emission in S.

Optical: Detected.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz
 $(3')$ and Parkes 64-m at 5 GHz $(4')$.
Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

$(43'' \times 59'': S=3.8 \text{ Jy})$.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G338.3–0.0

RA: $16^{\text{h}}41^{\text{m}}00^{\text{s}}$
Dec: $-46^{\circ}34'$

1-GHz flux/Jy: 7?
Spectral index: ?

Size/arcmin: 8
Type: C?

Radio: Irregular shell, in complex region.

X-ray: Central X-ray source and nebula.

Point sources: Central pulsar.

Distance: H α observations suggest 8 to 13 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz
 $(3')$ and Parkes 64-m at 5 GHz $(4')$.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 59'': S=7.4 \text{ Jy})$.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Aharonian *et al.* 2005, Science, 307, 1938. γ -ray detection.
Aharonian *et al.* 2006, ApJ, 636, 777. γ -ray observations.
Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
Funk *et al.* 2007, ApJ, 662, 517. XMM-Newton observations.
Lemiere *et al.* 2009, ApJ, 706, 1269. Chandra observations of central source and nebula.

Slane *et al.* 2010, ApJ, 720, 266. Fermi observations.
Castelletti *et al.* 2011, A&A, 536, A98. GMRT at 235 MHz
 $(10'' \times 26'')$, 610 MHz $(5''.0 \times 12''.6)$ and 1280 MHz $(5''.0 \times 6''.4)$, plus ATCA at 2.3 GHz $(4''.3 \times 4''.8)$.
Gotthelf *et al.* 2014, ApJ, 788, 155. NuSTAR observations.
Lemoine-Goumard *et al.* 2014, ApJ, 794, L16. γ -ray observations.
Abramowski *et al.* 2014, MNRAS, 439, 2828. γ -ray observations.
see also: Abramowski *et al.* 2014, MNRAS, 441, 3640. Erratum.
Supan *et al.* 2016, A&A, 589, A51. SGPS H α and other observations.

G338.5+0.1

RA: $16^{\text{h}}41^{\text{m}}09^{\text{s}}$
Dec: $-46^{\circ}19'$

1-GHz flux/Jy: 12?
Spectral index: ?

Size/arcmin: 9
Type: ?

Radio: Circle of non-thermal emission in complex region, not well defined.

Distance: H α absorption suggests 11 kpc.

References:

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz
 $(3')$ and Parkes 64-m at 5 GHz $(4')$.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 59'': S=13 \text{ Jy})$.
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including H α .
Abramowski *et al.* 2014, ApJ, 794, L1. γ -ray observations.

G340.4+0.4

RA: 16^h46^m31^s
Dec: -44°39'

1-GHz flux/Jy: 5
Spectral index: 0.4

Size/arcmin: 10×7
Type: S

Radio: Distorted shell, elongated east–west.

Optical: Detected.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 8.2$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 2.9$ Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz ($50''$).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz ($75'' \times 48'' : S = 9.8 \pm 0.9$ Jy) and 1.4 GHz ($27'' \times 9'' : S = 5.8 \pm 0.1$ Jy).

3.6 ± 0.1 Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 61'' : S = 5.9$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G340.6+0.3

RA: 16^h47^m41^s
Dec: -44°34'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 6
Type: S

Radio: Incomplete shell.

Optical: Possible associated filaments.

Distance: H α absorption suggests 15 kpc.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 7.0$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 2.8$ Jy).
 Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz ($50''$).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz ($75'' \times 48'' : S = 9.2 \pm 0.9$ Jy) and 1.4 GHz ($27'' \times 9'' : S = 5.8 \pm 0.1$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

($43'' \times 61'' : S = 4.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including H α .

G341.2+0.9

RA: 16^h47^m35^s
Dec: -43°47'

1-GHz flux/Jy: 1.5?
Spectral index: 0.6?

Size/arcmin: 22×16
Type: C

Radio: Incomplete shell, with extension to SW.

Point sources: Pulsar in W, with wind nebula.

References:

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz ($54'' \times 116'' : S = 3.0 \pm 0.1$ Jy) and 1.4 GHz ($21'' \times 25'' : S = 12.5 \pm 0.05$ Jy).
 Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4 and 4.9 GHz ($25''$) of pulsar wind nebula.

G341.9-0.3

RA: 16^h55^m01^s
Dec: -44°01'

1-GHz flux/Jy: 2.5
Spectral index: 0.5

Size/arcmin: 7
Type: S

Radio: Incomplete shell, brightest to NE.

References:

Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz ($3' : S = 7.4$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 1.7$ Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz ($50''$), revision of previous flux densities.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ($35'' \times 11'' : S = 2.2 \pm 0.1$ Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 62'' : S = 2.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.0–0.2

RA: $16^{\text{h}}54^{\text{m}}50^{\text{s}}$
Dec: $-43^{\circ}53'$

1-GHz flux/Jy: 3.5?
Spectral index: 0.4?

Size/arcmin: 12×9
Type: S

Radio: Distorted shell.

References:

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz ($50''$), estimate $S_{408 \text{ MHz}} = 5 \text{ Jy}$, $S_{5 \text{ GHz}} = 2 \text{ Jy}$ from previous maps.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz ($35'' \times 11''$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz

($43'' \times 62''$: $S = 3.5 \text{ Jy}$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.1+0.9

RA: $16^{\text{h}}50^{\text{m}}43^{\text{s}}$
Dec: $-43^{\circ}04'$

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 10×9
Type: S

Radio: Incomplete shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 63''$: $S = 0.6 \text{ Jy}$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G343.0–6.0

RA: $17^{\text{h}}25^{\text{m}}00^{\text{s}}$
Dec: $-46^{\circ}30'$

1-GHz flux/Jy: ?
Spectral index: ?

RCW 114

Size/arcmin: 250
Type: S

Radio: Faint, poorly defined.

Optical: Filamentary shell.

References:

Walker & Zealey 2001, MNRAS, 325, 287. Optical observations, and review of earlier observations.
 Welsh *et al.* 2003, A&A, 403, 605. Optical spectroscopy.

Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Kim *et al.* 2010, ApJ, 709, 823. UV observations.

G343.1–2.3

RA: $17^{\text{h}}08^{\text{m}}00^{\text{s}}$
Dec: $-44^{\circ}16'$

1-GHz flux/Jy: 8?
Spectral index: 0.5?

Size/arcmin: 32?
Type: C?

Radio: Incomplete shell?

X-ray: Pulsar wind nebula.

Point sources: Pulsar near edge, with wind nebula.

References:

McAdam *et al.* 1993, Nature, 361, 516. MOST at 843 MHz (smoothed to $2''$).
 Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz ($56'' \times 111''$: $S = 10.6 \text{ Jy}$) and 1.4 GHz ($22'' \times 27''$) near pulsar.
 Becker *et al.* 1995, A&A, 298, 528. ROSAT of pulsar, and limit for remnant.
 Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4, 4.9 and

8.5 GHz ($25''$) of pulsar wind nebula.
 Gotthelf *et al.* 2002, ApJ, 567, L125. Pulsar x-ray detection.
 Dodson & Golap 2002, MNRAS, 334, L1. ATCA at 1.4 GHz ($70'' \times 47''$) including polarisation, and Chandra observations of pulsar wind nebula.
 Aharonian *et al.* 2005, A&A, 432, L9. γ -ray limit.
 Abramowski *et al.* 2011, A&A, 528, A143. γ -ray observations.

G343.1–0.7

RA: 17^h00^m25^s
Dec: −43°14'

1-GHz flux/Jy: 7.8
Spectral index: 0.55

Size/arcmin: 27×21
Type: S

Radio: Shell, with smaller thermal shell adjacent.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×63'': S=8.5±0.6), plus Parkes 64-m at 4.5 GHz (S=3.9±0.6 Jy) and 8.55 GHz (S=2.4±0.5 Jy)
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G344.7–0.1

RA: 17^h03^m51^s
Dec: −41°42'

1-GHz flux/Jy: 2.5?
Spectral index: 0.3?

Size/arcmin: 8
Type: C?

Radio: Asymmetric shell, with possible core.

X-ray: Detected.

Distance: H_I absorption and association with features suggests 6.3 kpc.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S=4.7 Jy) and Parkes 64-m at 5 GHz (4': S=1.3 Jy).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (30''×43'': S=1.7±0.1 Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×65'': S=2.5 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Yamauchi *et al.* 2005, PASJ, 57, 459. ASCA observations.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Giacani *et al.* 2011, A&A, 531, A138. VLA and ATCA at 1.4 GHz (5''6×8''), ATCA at 5 GHz (10''×13'') and 8.4 GHz (5''×10''), plus XMM-Newton observations and H_I from SGPS.
Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
Yamaguchi *et al.* 2012, ApJ, 749, 137. Suzaku observations.
Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.

G345.7–0.2

RA: 17^h07^m20^s
Dec: −40°53'

1-GHz flux/Jy: 0.6?
Spectral index: ?

Size/arcmin: 6
Type: S

Radio: Poorly defined diffuse shell.

Point sources: Old pulsar nearby.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×66'': S=0.7 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G346.6–0.2

RA: 17^h10^m19^s
Dec: −40°11'

1-GHz flux/Jy: 8?
Spectral index: 0.5?

Size/arcmin: 8
Type: S

Radio: Irregular shell.

X-ray: Centrally brightened.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': S=14.9 Jy) and Parkes 64-m at 5 GHz (4': S=4.3 Jy).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (31''×43'': S=8.1±0.9 Jy).
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×67'': S=8.7 Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact

OH emission.
Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
Sezer *et al.* 2011, MNRAS, 415, 301. Suzaku observations.
Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
Yamauchi *et al.* 2013, PASJ, 65, 6. Suzaku observations.
Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
Pannuti *et al.* 2014, AJ, 147, 55. ASCA observations.

G347.3–0.5		RX J1713.7–3946			
RA: 17 ^h 13 ^m 50 ^s	1-GHz flux/Jy: 30?	Size/arcmin: 65×55	Type: S?		
Dec: −39°45'	Spectral index: ?				
Radio: Faint emission.					
X-ray: Non-thermal, limb-brightened to W, with central source.					
Point sources: Central X-ray source.					
Distance: Association with molecular clouds and X-ray observations imply 1.3 kpc.					
References:					
Koyama <i>et al.</i> 1997, PASJ, 49, L7. ASCA of NW.					
Slane <i>et al.</i> 1999, ApJ, 525, 357. ASCA and other observations.					
Muraishi <i>et al.</i> 2000, A&A, 354, L57. High energy γ-ray detection.					
Butt <i>et al.</i> 2001, ApJ, 562, L167. Associated γ-ray emission.					
Ellison <i>et al.</i> 2001, ApJ, 563, 191. ATCA at 1.4 GHz (46''×36''), and ASCA observations.					
Enomoto <i>et al.</i> 2002, Nature, 416, 823. γ-ray observations.					
Uchiyama <i>et al.</i> 2002, PASJ, 54, L73. ASCA observations.					
Uchiyama <i>et al.</i> 2003, A&A, 400, 567. Chandra spectroscopy.					
Pannuti <i>et al.</i> 2003, ApJ, 593, 377. ROSAT, ASCA and RXTE observations.					
Lazendic <i>et al.</i> 2003, ApJ, 593, L27. Chandra, XMM-Newton and other X-ray observations of central source.					
Fukui <i>et al.</i> 2003, PASJ, 55, L61. CO observations of surroundings.					
Cassam-Chenaï <i>et al.</i> 2004, A&A, 427, 199. XMM-Newton and other observations.					
Aharonian <i>et al.</i> 2004, Nature, 432, 75. γ-ray detection.					
Lazendic <i>et al.</i> 2004, ApJ, 602, 271. Chandra observations of parts, and ATCA at 1.4 GHz (36''×46'').					
Hiraga <i>et al.</i> 2005, A&A, 431, 953. XMM-Newton observations,					
Morlighem <i>et al.</i> 2005, ApJ, 631, 947. CO observations of surroundings.					
Aharonian <i>et al.</i> 2007, A&A, 464, 235. γ-ray observations. see also: Aharonian <i>et al.</i> 2011, A&A, 531, C1. Erratum.					
Uchiyama <i>et al.</i> 2007, Nature, 449, 576. Chandra multi-epoch observations for study of small-scale variability.					
Tanaka <i>et al.</i> 2008, ApJ, 685, 988. Suzaku observations.					
Takahashi <i>et al.</i> 2008, PASJ, 60, S131. Suzaku observations of SW.					
Mignani <i>et al.</i> 2008, A&A, 484, 457. Optical and IR observations of central source.					
Acero <i>et al.</i> 2009, A&A, 505, 157. XMM-Newton observations, plus other radio, IR and γ-ray data.					
Sano <i>et al.</i> 2010, ApJ, 724, 59. CO observations of region.					
Maxted <i>et al.</i> 2012, MNRAS, 422, 2230. Molecular line observations. see also: Maxted <i>et al.</i> 2013, MNRAS, 430, 2511. Erratum.					
Sano <i>et al.</i> 2013, ApJ, 778, 59. CO, SGPS HI and Suzaku observations.					
Federici <i>et al.</i> 2015, A&A, 577, A12. γ-ray detection.					
Sano <i>et al.</i> 2015, ApJ, 799, 175. Suzaku observations.					
Katsuda <i>et al.</i> 2015, ApJ, 814, 29. XMM-Newton observations.					
Acero <i>et al.</i> 2016, ApJS, 224, 8. γ-ray observations.					
Tsujii & Uchiyama 2016, PASJ, 68, 108. Chandra observations of NW.					

G348.5–0.0		Size/arcmin: 10?	
RA: 17 ^h 15 ^m 26 ^s	1-GHz flux/Jy: 10?	Type: S?	
Dec: −38°28'	Spectral index: 0.4?		
Radio: Arc, overlapping G348.5+0.1.			
References:			
Kassim <i>et al.</i> 1991, ApJ, 374, 212. VLA at 333 MHz (46''×53''), 1.4 GHz (18''×33'') and part at 5 GHz (2''.5×3''.9).			
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×69'': $S=10.2$ Jy).			
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant, including masers.			
Green <i>et al.</i> 1997, AJ, 114, 2058. Parkes 64-m OH observations.			
Koralesky <i>et al.</i> 1998, AJ, 116, 1323. VLA search for OH emis-			
sion.			
Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer observations.			
Hewitt <i>et al.</i> 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.			
Hewitt <i>et al.</i> 2009, ApJ, 694, 1266. Spitzer spectroscopy.			
Anderson <i>et al.</i> 2011, ApJ, 742, 7. Spitzer observations.			
Pihlström <i>et al.</i> 2014, AJ, 147, 73. VLA search for methanol masers.			
Pannuti <i>et al.</i> 2014, AJ, 147, 55. XMM-Newton observations for upper limit.			

G348.5+0.1			CTB 37A
RA: $17^{\text{h}}14^{\text{m}}06^{\text{s}}$	1-GHz flux/Jy: 72	Size/arcmin: 15	
Dec: $-38^{\circ}32'$	Spectral index: 0.3	Type: S	
Radio: Shell, poorly define to S and W, overlapping G348.5–0.0 in E.			
X-ray: Brighter to W.			
Distance: H α absorption indicates 8.0 kpc.			
References:			
Clark <i>et al.</i> 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 97$ Jy).	Zeeman splitting.		
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ($4'4 : S = 43$ Jy).	Reynoso & Mangum 2000, ApJ, 545, 874. CO observations.		
Caswell <i>et al.</i> 1975, A&A, 45, 239. Parkes H α absorption.	Reach <i>et al.</i> 2006, AJ, 131, 1479. Spitzer observations.		
Milne <i>et al.</i> 1979, MNRAS, 188, 437. FIRST at 1415 MHz ($0'8 : S > 50$) and Parkes 64-m at 14.7 GHz ($2'2 : S = 18 \pm 5$ Jy).	Aharonian <i>et al.</i> 2008, A&A, 490, 685. γ -ray observations.		
Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz ($20'' \times 45''$).	Castro & Slane 2010, ApJ, 717, 372. Fermi observations.		
Kassim <i>et al.</i> 1991, ApJ, 374, 212. VLA at 333 MHz ($46'' \times 53''$), 1.4 GHz ($18'' \times 33''$) and part at 5 GHz ($2'5 \times 3''9$).	Anderson <i>et al.</i> 2011, ApJ, 742, 7. Spitzer observations.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 69'' : S = 71$ Jy).	Sezer <i>et al.</i> 2011, MNRAS, 417, 1387. Suzaku observations.		
Brogan <i>et al.</i> 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH	Tian & Leahy 2012, MNRAS, 421, 2593. SGPS H α absorption observations.		
	Maxted <i>et al.</i> 2013, MNRAS, 434, 2188. Molecular line observations of region.		
	Pannuti <i>et al.</i> 2014, AJ, 147, 55. XMM-Newton and Chandra observations.		
	Yamauchi <i>et al.</i> 2014, PASJ, 66, 2. Suzaku observations.		
	Acero <i>et al.</i> 2016, ApJS, 224, 8. γ -ray observations.		

G348.7+0.3			CTB 37B
RA: $17^{\text{h}}13^{\text{m}}55^{\text{s}}$	1-GHz flux/Jy: 26	Size/arcmin: 17?	
Dec: $-38^{\circ}11'$	Spectral index: 0.3	Type: S	
Radio: Incomplete shell with faint eastern extensions.			
X-ray: Diffuse emission.			
Point sources: X-ray pulsar.			
Distance: H α absorption suggests 13 kpc.			
References:			
Clark <i>et al.</i> 1975, AuJPA, 37, 75. Molonglo at 408 MHz ($3' : S = 34$ Jy).	Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant.		
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz ($4'4 : S = 32$ Jy).	Aharonian <i>et al.</i> 2006, ApJ, 636, 777. γ -ray detection.		
Caswell <i>et al.</i> 1975, A&A, 45, 239. Parkes H α absorption.	Aharonian <i>et al.</i> 2008, A&A, 486, 829. γ -ray and Chandra observations.		
Milne <i>et al.</i> 1979, MNRAS, 188, 437. FIRST at 1415 MHz ($0'8 : S > 20$) and Parkes 64-m at 14.7 GHz ($2'2 : S = 8 \pm 3$ Jy).	Nakamura <i>et al.</i> 2009, PASJ, 61, S197. Suzaku and Chandra observations.		
Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz ($20'' \times 45''$).	Halpern & Gotthelf 2010, ApJ, 710, 941. Chandra detection of pulsar.		
Kassim <i>et al.</i> 1991, ApJ, 374, 212. VLA at 333 MHz ($46'' \times 53''$).	Halpern & Gotthelf 2010, ApJ, 725, 1384. Chandra observations of pulsar.		
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 69'' : S = 33$ Jy).	Tian & Leahy 2012, MNRAS, 421, 2593. SGPS H α observations.		
	Xin <i>et al.</i> 2016, ApJ, 817, 64. γ -ray observations.		

G349.2–0.1			
RA: $17^{\text{h}}17^{\text{m}}15^{\text{s}}$	1-GHz flux/Jy: 1.4?	Size/arcmin: 9×6	
Dec: $-38^{\circ}04'$	Spectral index: ?	Type: S	
Radio: Elongated shell, adjacent to bright H α region.			
References:			
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 70'' : S = 1.6$ Jy).			
Frail <i>et al.</i> 1996, AJ, 111, 1651. OH emission near remnant.			

G349.7+0.2

RA: 17^h17^m59^s
Dec: −37°26'

1-GHz flux/Jy: 20
Spectral index: 0.5

Size/arcmin: 2.5×2
Type: S

Radio: Incomplete clumpy shell, with enhancement to the S.

X-ray: Irregular shell, brighter to S and E.

Distance: HI absorption indicates 11.5 kpc.

References:

- Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').
 Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3' : $S = 31.0$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 9.1$ Jy), no maps.
 Caswell *et al.* 1975, A&A, 45, 239. Parkes HI absorption.
 Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz (3'4×14'5).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×71'' : $S = 22$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Reynoso & Mangum 2001, AJ, 121, 347. CO observations of the vicinity.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Slane *et al.* 2002, ApJ, 580, 904. ASCA observations.
 Dubner *et al.* 2004, A&A, 426, 201. CO observations of surroundings.
 Lazendic *et al.* 2005, ApJ, 618, 733. Chandra observations.

- Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Castro & Slane 2010, ApJ, 717, 372. Fermi observations.
 Lazendic *et al.* 2010, MNRAS, 409, 371. OH, CO and other molecular line observations of region.
 Anderson *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 TIAN & Leahy 2014, ApJ, 783, L2. SGPS HI observations.
 Yasumi *et al.* 2014, PASJ, 66, 68. Suzaku observations.
 Abramowski *et al.* 2015, A&A, 574, A100. γ -ray observations.
 see also: Abramowski *et al.* 2015, A&A, 580, C1. Correction.
 Ergin *et al.* 2015, ApJ, 804, 124. γ -ray and Suzaku observations.
 Rho *et al.* 2015, ApJ, 812, 44. Herschel observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G350.0−2.0

RA: 17^h27^m50^s
Dec: −38°32'

1-GHz flux/Jy: 26
Spectral index: 0.4

Size/arcmin: 45
Type: S

Incorporates the previously catalogued G350.0−1.8 in the NW.

Radio: Shell, brightest in NW.

Optical: Detected.

X-ray: Brighter to NW.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 49.5$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 13.6$ Jy).
 Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4'4).
 Gaensler 1998, ApJ, 493, 781. VLA and Parkes 64-m at

- 1.4 GHz (18''×21'' : $S = 22.3 \pm 0.3$ Jy), clarifying extent of remnant.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Karpova *et al.* 2016, MNRAS, 462, 3845. XMM-Newton observations.

G350.1−0.3

RA: 17^h21^m05^s
Dec: −37°27'

1-GHz flux/Jy: 6?
Spectral index: 0.8?

Size/arcmin: 4?
Type: ?

Radio: Several clumps of emission.

X-ray: Diffuse emission, with compact source.

Point sources: X-ray source.

Distance: HI absorption indicates 4.5 to 10.7 kpc, possible interaction with molecular cloud indicates 4.5 kpc.

References:

- Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3' : $S = 10.7$) and Parkes 64-m at 5 GHz (4' : $S = 1.7$).
 Salter *et al.* 1986, A&A, 162, 217. VLA at 1.5 (4''×10'') and 4.8 GHz (15''×35'') and 15 GHz (5''×2'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×71'').
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observa-

- tions.
 Gaensler *et al.* 2008, ApJ, 680, L37. VLA at 4.8 GHz (11''4×5''.6) plus XMM-Newton observations.
 Lovchinsky *et al.* 2011, ApJ, 731, 70. Chandra and Spitzer observations.
 Yasumi *et al.* 2014, PASJ, 66, 68. Suzaku observations.

G351.0–5.4

RA: 17^h46^m00^s
Dec: −39°25'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 30
Type: S

Radio: Faint shell, brighter to E and W.

References:

de Gasperin *et al.* 2014, A&A, 568, A107. GMRT at 325 MHz
 $(89'' \times 141'')$, plus other observations.

G351.2+0.1

RA: 17^h22^m27^s
Dec: −36°11'

1-GHz flux/Jy: 5?
Spectral index: 0.4

Size/arcmin: 7
Type: C?

Has been called G351.3+0.2.

Radio: Distorted shell, with possible flat-spectrum core.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 8.1$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 3.1$ Jy).
 Becker & Helfand 1988, AJ, 95, 883. VLA at 5 GHz ($15''$), and
 at 15 GHz of core.
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz

$(32'' \times 36'') : S = 4.8 \pm 0.2$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 73'') : S = 5.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-
 tions.

G351.7+0.8

RA: 17^h21^m00^s
Dec: −35°27'

1-GHz flux/Jy: 10
Spectral index: 0.5?

Size/arcmin: 18×14
Type: S

Radio: Elongated shell, adjacent to bright HII region.

Point sources: Pulsar nearby.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 74'') : S = 11$ Jy).
 Tian *et al.* 2007, MNRAS, 378, 1283. SGPS at 1.4 GHz ($100'' : S = 8.4 \pm 0.7$ Jy) including HI.

G351.9–0.9

RA: 17^h28^m52^s
Dec: −36°16'

1-GHz flux/Jy: 1.8?
Spectral index: ?

Size/arcmin: 12×9
Type: S

Radio: Asymmetric shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz
 $(43'' \times 73'') : S = 2.0$ Jy).

G352.7–0.1

RA: $17^{\text{h}}27^{\text{m}}40^{\text{s}}$
Dec: $-35^{\circ}07'$

1-GHz flux/Jy: 4
Spectral index: 0.6

Size/arcmin: 8×6
Type: S

Radio: Distorted shell.

X-ray: Detected.

Distance: HI absorption indicates 6.8 to 8.4 kpc.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz ($3' : S = 9.6$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 2.3$ Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz ($1.1'$).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($34'' : S = 3.4 \pm 0.4$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 75'' : S = 4.4$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Kinugasa *et al.* 1998, PASJ, 50, 249. ASCA observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Giacani *et al.* 2009, A&A, 507, 841. VLA at 4.8 GHz ($9'' \times 12''$), plus HI and XMM-Newton observations.
 Pannuti *et al.* 2014, ApJ, 782, 102. XMM-Newton and Chandra observations.
 Sezer & Gök 2014, ApJ, 790, 81. Suzaku observations.

G353.6–0.7

RA: $17^{\text{h}}32^{\text{m}}00^{\text{s}}$
Dec: $-34^{\circ}44'$

1-GHz flux/Jy: 2.5?
Spectral index: ?

Size/arcmin: 30
Type: S

Has erroneously been called G353.6–0.37.

Radio: Shell, brighter to S.

X-ray: Detected.

Point sources: Central X-ray source.

References:

Tian *et al.* 2008, ApJ, 679, L85. SGPS at 1.4 GHz ($100''$), plus 843 MHz ($43''$) and X-ray observations.
 Halpern & Gotthelf 2010, ApJ, 710, 941. XMM-Newton observations.
 Tian *et al.* 2010, ApJ, 712, 790. XMM-Newton, Suzaku and CO observations.
 Abramowski *et al.* 2011, A&A, 531, A81. γ -ray observations.
 Bamba *et al.* 2012, ApJ, 756, 149. Suzaku observations.

see also: Bamba *et al.* 2012, ApJ, 761, 80. Erratum.
 Klochkov *et al.* 2013, A&A, 556, A41. XMM-Newton and other X-ray observations.
 Yang *et al.* 2014, A&A, 567, A23. γ -ray limit.
 Fukuda *et al.* 2014, ApJ, 788, 94. γ -ray, SGPS HI and CO observations.
 Acero *et al.* 2015, A&A, 580, A74. γ -ray limit.

G353.9–2.0

RA: $17^{\text{h}}38^{\text{m}}55^{\text{s}}$
Dec: $-35^{\circ}11'$

1-GHz flux/Jy: 1?
Spectral index: 0.5?

Size/arcmin: 13
Type: S

Radio: Shell, with central double source.

References:

Green 2001, MNRAS, 326, 283. VLA at 327 MHz ($3.0' \times 2.7'$) and 1.4 GHz ($42'' \times 36''$), plus 8.4 GHz ($8.4'' \times 6.1''$) of central source only.

G354.1+0.1

RA: $17^{\text{h}}30^{\text{m}}28^{\text{s}}$
Dec: $-33^{\circ}46'$

1-GHz flux/Jy: ?
Spectral index: varies

Size/arcmin: $15 \times 3?$
Type: C?

Is this a SNR?

Radio: Elongated N–S.

Point sources: Pulsar at S tip.

References:

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz ($47'' \times 99''$) and 1.4 GHz ($8.8'' \times 21''$).
 Ajello *et al.* 2016, ApJ, 819, 44. γ -ray observations.

G354.8–0.8

RA: 17^h36^m00^s
Dec: −33°42'

1-GHz flux/Jy: 2.8?
Spectral index: ?

Size/arcmin: 19
Type: S

Radio: Distorted shell.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×78'': $S=3.1$ Jy).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observa-

tions.
Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G355.4+0.7

RA: 17^h31^m20^s
Dec: −32°26'

1-GHz flux/Jy: 5?
Spectral index: ?

Size/arcmin: 25
Type: S

Radio: Faint, incomplete shell.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×77'').
Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1': $S=8.9\pm1.3$ Jy).
Ajello *et al.* 2016, ApJ, 819, 44. γ-ray observations.
Acero *et al.* 2016, ApJS, 224, 8. γ-ray observations.

G355.6–0.0

RA: 17^h35^m16^s
Dec: −32°38'

1-GHz flux/Jy: 3?
Spectral index: ?

Size/arcmin: 8×6
Type: S

Radio: Well defined shell.

X-ray: Centrally brightened.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×80'': $S=2.6$ Jy).
Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1': $S=3.3\pm0.5$ Jy).
Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.
Minami *et al.* 2013, PASJ, 65, 99. Suzaku observations.

G355.9–2.5

RA: 17^h45^m53^s
Dec: −33°43'

1-GHz flux/Jy: 8
Spectral index: 0.5

Size/arcmin: 13
Type: S

Radio: Distorted shell, brightest to SE.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=12.3$ Jy) and Parkes 64-m at 5 GHz (4': $S=3.4$ Jy).
Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (32''×34'': $S=5.0\pm0.3$ Jy).
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×77'').
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.

G356.2+4.5

RA: 17^h19^m00^s
Dec: −29°40'

1-GHz flux/Jy: 4
Spectral index: 0.7

Size/arcmin: 25
Type: S

Has been called G356.2+4.4.

Radio: Faint shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10'.4).
Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz (3'×1.5': $S=8.1\pm1.7$ Jy), and NVSS at 1.4 GHz.

G356.3–1.5

RA: 17^h42^m35^s
Dec: −32°52'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 20×15
Type: S

Radio: Double arc.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×79'': $S=2.8$ Jy).
Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (0.8''×1.7'': $S=5.7\pm0.2$).

G356.3–0.3

RA: 17^h37^m56^s
Dec: −32°16'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 11×7
Type: S

Has been suggested this part of a larger SNR.

Radio: Diffuse emission.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×81'': $S=2.6$ Jy).
Roy & Pramesh Rao 2002, MNRAS, 329, 775. GMRT at 330 MHz (2.7''×4.8').
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G357.7–0.1

RA: 17^h40^m29^s
Dec: −30°58'

1-GHz flux/Jy: 37
Spectral index: 0.4

MSH 17–39

Size/arcmin: 8×3?
Type: ?

Has been suggested that this is not a SNR.

Radio: Multiple arcs and filaments, brighter to NW ‘head’.

X-ray: Detected from NW ‘head’, and SW ‘tail’.

Distance: H_I absorption suggests beyond Galactic Centre.

References:

Caswell *et al.* 1975, AuJPA, 37, 39. Molonglo at 408 MHz (3': $S=54.2$ Jy) and Parkes 64-m at 5 GHz (4': $S=18.5$ Jy).
Milne & Dickel 1975, AuJPh, 28, 209. Parkes 64-m at 5 GHz (4.4': $S=14.6$ Jy).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6').
Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50'').
Weiler & Panagia 1980, A&A, 90, 269. Effelsberg 100-m at 9 GHz (1.5') (private communication from Baker).
Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz (3''.8×10''.9) and 5 GHz (12''.8×26'').
Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 GHz and 5 GHz.
Helfand & Becker 1985, Nature, 313, 118. Suggest it is not a SNR.
Shaver *et al.* 1985, A&A, 147, L23. Observations of peripheral compact source.
Caswell *et al.* 1989, PASA, 8, 184. MOST at 843 MHz (43''×83'').
Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×84'').
Stewart *et al.* 1994, ApJ, 432, L39. ATCA at 4.79 and 5.84 GHz (12''×22'') and Effelsberg 100-m at 10.6 GHz (1'), including polarisation.
Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43''×24'').
see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
Brogan & Goss 2003, AJ, 125, 272. VLA at 1.4 GHz (13''.6×11''.4), including H_I, and 8.3 GHz (14''.3×6''.8) recombination line observation of HII region.
Gaensler *et al.* 2003, ApJ, 594, L35. Chandra detection.
Lazendic *et al.* 2003, AN, 324 (No S1), 157. Molecular line observations.
Burton *et al.* 2004, MNRAS, 348, 638. IR and radio observations of HII region.
Lazendic *et al.* 2004, MNRAS, 354, 393. IR and molecular line observations.
Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
Phillips *et al.* 2009, MNRAS, 397, 1215. Observations of interactions with surroundings.
Castro *et al.* 2013, ApJ, 774, 36. Fermi observations.
Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
Acero *et al.* 2016, ApJS, 224, 8. γ -ray observations.

G357.7+0.3

RA: $17^{\text{h}}38^{\text{m}}35^{\text{s}}$
Dec: $-30^{\circ}44'$

1-GHz flux/Jy: 10
Spectral index: 0.4?

Size/arcmin: 24
Type: S

Radio: Non-thermal shell in complex region.

References:

Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz ($4.3:S=7\pm1.5$ Jy), $S_{5 \text{ GHz}} = 5.5\pm1.5$ Jy from surveys.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43''\times84''$).
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for

OH masers.
 Phillips & Marquez-Lugo 2010, MNRAS, 409, 701. Spitzer observations of region.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G358.0+3.8

RA: $17^{\text{h}}26^{\text{m}}00^{\text{s}}$
Dec: $-28^{\circ}36'$

1-GHz flux/Jy: 1.5?
Spectral index: ?

Size/arcmin: 38
Type: S

Radio: Faint shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz ($10'.4$).
 Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz ($2'.2\times1'.3:S=2.5\pm1.3$ Jy), and NVSS at 1.4 GHz.

G358.1+1.0

RA: $17^{\text{h}}37^{\text{m}}00^{\text{s}}$
Dec: $-29^{\circ}59'$

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 20
Type: S

Was erroneously called G358.1+0.1.

Radio: Faint shell.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43''\times77''$).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1':S=6.0\pm2.5$ Jy).

G358.5-0.9

RA: $17^{\text{h}}46^{\text{m}}10^{\text{s}}$
Dec: $-30^{\circ}40'$

1-GHz flux/Jy: 4?
Spectral index: ?

Size/arcmin: 17
Type: S

Radio: Shell, brighter to NE.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43''\times77''$).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1':S=8.0\pm2.5$ Jy).

G359.0–0.9

RA: 17^h46^m50^s
Dec: −30°16'

1-GHz flux/Jy: 23
Spectral index: 0.5

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Optical: Detected.

X-ray: Partial shell.

References:

- Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'.3).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×86'').
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43''×24'').
 see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.

- Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.4×12''.8) of part.
 Bamba *et al.* 2009, ApJ, 691, 1854. Suzaku observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. Hα observations.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G359.1–0.5

RA: 17^h45^m30^s
Dec: −29°57'

1-GHz flux/Jy: 14
Spectral index: 0.4?

Size/arcmin: 24
Type: S

Radio: Non-thermal shell in complex region, crossed by the ‘snake’.

Optical: Detected.

X-ray: Centrally brightened.

Point sources: Several compact radio sources near centre, OH masers around edge.

References:

- Downes *et al.* 1979, A&AS, 35, 1. From observations by Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6:S=13 Jy).
 Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz (4'.3 : S = 10±1.5 Jy) and 4.8 GHz (2'.4 : S = 8.1±0.5 Jy).
 Uchida *et al.* 1992, ApJ, 398, 128. VLA at 1.5 GHz (11''×10''), and observations of nearby molecular material.
 Uchida *et al.* 1992, AJ, 104, 1533. VLA at 1.4 GHz.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×85'').
 Yusef-Zadeh *et al.* 1995, Science, 270, 1801. VLA at 1.4 GHz (33''×31''), and 1.7 GHz for OH survey.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (43''×24'').
 see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.
 Lazendic *et al.* 2002, MNRAS, 331, 537. Observations of

- shocked molecular gas where the ‘snake’ crosses the remnant.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.4×12''.8).
 Aharonian *et al.* 2008, A&A, 483, 509. XMM-Newton and γ-ray observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Bamba *et al.* 2009, ApJ, 691, 1854. Suzaku observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. Hα observations.
 Ohnishi *et al.* 2011, PASJ, 63, 527. Suzaku observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Hui *et al.* 2016, MNRAS, 457, 4262. γ-ray observations of region.

G359.1+0.9

RA: 17^h39^m36^s
Dec: −29°11'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 12×11
Type: S

Radio: Shell, brightest in E.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×88'': S=4.3 Jy).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz

- (1':S=4.3±1.0 Jy).
 Law *et al.* 2008, ApJS, 177, 515. VLA at 1.4 GHz (10''.9×15''.9: S=1.3±0.5 Jy).