

G0.0+0.0

Sgr A East

RA: 17^h45^m44^s**Dec:** -29°00'**1-GHz flux/Jy:** 100?**Spectral index:** 0.8?**Size/arcmin:** 3.5×2.5**Type:** S

Radio: Non-thermal shell, in complex region, interacting with molecular material to the west.

X-ray: Diffuse emission, centrally peaked.

Point sources: Compact X-ray/radio source.

References:

- Ekers *et al.* 1983, A&A, 122, 143. VLA at 1.4 GHz and 5 GHz (both 5''×8''), $S_{1.4\text{ GHz}} = 77\text{ Jy}$, $S_{5.0\text{ GHz}} = 31\text{ Jy}$.
- Pedlar *et al.* 1989, ApJ, 342, 769. VLA at 332 MHz (12''), 1.4 GHz (1''3×2''5) and 5 GHz (1''3×2''5).
- Mezger *et al.* 1989, A&A, 209, 337. Nearby molecular material.
- Ho *et al.* 1991, Nature, 350, 309. VLA of NH₃ emission from surroundings.
- Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (17''×33'').
- Serabyn *et al.* 1992, ApJ, 395, 166. Nearby molecular material.
- Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×88'').
- Yusef-Zadeh & Mehringer 1995, ApJ, 452, L37. VLA of nearby H₂O masers.
- Yusef-Zadeh *et al.* 1999, ApJ, 512, 230. OH maser observations.
- Coil & Ho 2000, ApJ, 533, 245. NH₃ observations of surroundings.
- LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24''×43'').
see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
- Fatuzzo *et al.* 2001, ApJ, 549, 293. Electron-positron lines from the vicinity.
- Yusef-Zadeh *et al.* 2001, ApJ, 560, 749. Observations of nearby molecular hydrogen.
- Maeda *et al.* 2002, ApJ, 570, 671. Chandra observations.
- Sakano *et al.* 2003, AN, 324 (No S1), 197. XMM-Newton observations.
- Roy & Rao 2004, MNRAS, 349, L25. GMRT at 620 MHz (6'.6×11'.4).
- Sakano *et al.* 2004, MNRAS, 350, 129. XMM-Newton observations.
- Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (1''×2''.2).
- Park *et al.* 2005, ApJ, 631, 964. Chandra observations.
- Aharonian *et al.* 2006, ApJ, 636, 777. HESS detection.
- Koyama *et al.* 2007, PASJ, 59, S237. Suzaku observations.
- Lee *et al.* 2008, ApJ, 674, 247. Molecular H₂ observations of surroundings.
- Sjouwerman & Pihlström 2008, ApJ, 681, 1287. VLA at 1.7 GHz of OH masers.
- Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
- Tsuboi *et al.* 2012, PASJ, 64, 111. CO observations of SW.
- Minh *et al.* 2013, ApJ, 773, 31. NH₃ observations of region.
- Zhao *et al.* 2013, ApJ, 777, 146. VLA at 4.8 GHz (0''.5×0''.7), 5.5 GHz (0''.6×1''.6), and 8.3 GHz (1''.0×2''.0).
- Nynka *et al.* 2013, ApJ, 778, L31. NuSTAR of compact X-ray source.
- Pihlström *et al.* 2014, AJ, 147, 73. VLA observations of methanol masers.
- Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
- Tsuboi *et al.* 2015, PASJ, 67, 109. CO observations of region.
- Lau *et al.* 2015, Science, 348, 413. IR observations.
- Ajello *et al.* 2016, ApJ, 819, 44. Fermi observations.
- Yusef-Zadeh *et al.* 2016, ApJ, 819, 60. VLA at 1.5 GHz (0''.5×1''.4) and other frequencies.
- McEwen *et al.* 2016, ApJ, 832, 129. CH₃OH maser observations in region.
- Qiao *et al.* 2018, ApJS, 239, 15. OH maser observations.

G0.3+0.0**RA:** 17^h46^m15^s**Dec:** −28°38′**1-GHz flux/Jy:** 22**Spectral index:** 0.6**Size/arcmin:** 15×8**Type:** S

Has been called G0.33+0.04, G0.30+0.04 and G0.4+0.1.

Radio: Bilateral shell, near Galactic Centre.

X-ray: Diffuse emission.

References:

Kassim & Frail 1996, MNRAS, 283, L51. VLA at 333 MHz (23″×42″), plus review of flux densities and other observations.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (30″).

see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30″).

Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.

Ajello *et al.* 2016, ApJ, 819, 44. Fermi observations.

G0.9+0.1**RA:** 17^h47^m21^s**Dec:** −28°09′**1-GHz flux/Jy:** 18?**Spectral index:** varies**Size/arcmin:** 8**Type:** C

Radio: Flat spectrum core within steep spectrum shell.

X-ray: Central core, with non-thermal spectrum.

Point sources: Central pulsar.

References:

Helfand & Becker 1987, ApJ, 314, 203. VLA at 1.4 and 5 GHz, and Einstein observations.

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43″×91″).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Mereghetti *et al.* 1998, A&A, 331, L77. X-ray detection.

LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24″×43″).

see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.

Sidoli *et al.* 2000, A&A, 361, 719. BeppoSAX observations.

Gaensler *et al.* 2001, ApJ, 556, L107. Chandra observations.

Porquet *et al.* 2003, A&A, 401, 197. XMM-Newton observations.

Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8″.6×11″.3).

Aharonian *et al.* 2005, A&A, 432, L25. HESS detection.

Dubner *et al.* 2008, A&A, 487, 1033. ATCA and VLA at 1.4 GHz (1″.2×2″.5; $S=8.3\pm 0.7$ Jy), 5 GHz (1″.6×2″.5) and 8.3 GHz (0″.8×1″.5).

Camilo *et al.* 2009, ApJ, 700, L34. Pulsar detection.

Holler *et al.* 2012, A&A, 539, A24. Chandra and XMM-Newton observations.

Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.

Archer *et al.* 2016, ApJ, 821, 129. γ -ray observations.

Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.

G1.0–0.1

RA: 17^h48^m30^s
Dec: –28°09′

1-GHz flux/Jy: 15
Spectral index: 0.6?

Size/arcmin: 8
Type: S

Has been called G1.05–0.1 and G1.05–0.15.

Radio: Incomplete shell, to the S of Sgr D.

X-ray: Possibly detected.

References:

- Downes *et al.* 1979, A&AS, 35, 1. Review of flux densities.
 Anantharamaiah *et al.* 1991, MNRAS, 249, 262. VLA at 330 MHz (64''×100'' : S=12.3 Jy).
 Liszt 1992, ApJS, 82, 495. VLA at 1.6 GHz (13''×23'').
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×91'').
 Mehringer *et al.* 1998, ApJ, 493, 274. VLA at 1.6 GHz (15''×28'') and 5 GHz, including masers observations.
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24''×43'').
see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Sidoli *et al.* 2001, A&A, 372, 651. BeppoSAX possible detection.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (30'').
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Nobukawa *et al.* 2009, AdSpR, 43, 1045. Suzaku observations.
 Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G1.4–0.1

RA: 17^h49^m39^s
Dec: –27°46′

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 10
Type: S

Radio: Shell, brightest in E.

X-ray: Diffuse emission.

References:

- Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×92'' : S=2 Jy).
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (2'.4×2'.7 : S=4.2±0.5).
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8''.2×12''.2).
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA observations of methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 McEwen *et al.* 2016, ApJ, 826, 189. NH₃ and CH₃OH observations.

G1.9+0.3**RA:** 17^h48^m45^s**Dec:** -27°10'**1-GHz flux/Jy:** 0.6**Spectral index:** 0.6**Size/arcmin:** 1.5**Type:** S**Radio:** Shell, brighter to the N, brightening; shows secular increase.**X-ray:** Shell, with bright limbs to E and W.**Distance:** H_i absorption gives < 10 kpc.**References:**Green & Gull 1984, *Nature*, 312, 527. VLA at 5 GHz (2''×4''4).Gray 1994, *MNRAS*, 270, 835. MOST at 843 MHz (43''×94'').Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.Yusef-Zadeh *et al.* 2004, *ApJS*, 155, 421. VLA at 1.4 GHz (8''3×12'').Nord *et al.* 2004, *AJ*, 128, 1646. VLA at 330 MHz (7''×12'').Green 2004, *BASI*, 32, 335. VLA at 1.5 GHz (7''2×9''4).Reynolds *et al.* 2008, *ApJ*, 680, L41. Chandra observations.Green *et al.* 2008, *MNRAS*, 387, L54. VLA at 4.86 GHz (4''×10''), for expansion studies.Murphy *et al.* 2008, *MNRAS*, 389, L23. MOST at 843 MHz for flux increase.Gómez & Rodríguez 2009, *RMxAA*, 45, 91. VLA at 1.5 GHz (5''1×10''6).Reynolds *et al.* 2009, *ApJ*, 695, L149. Chandra spectroscopy.Borkowski *et al.* 2010, *ApJ*, 724, L161. Chandra observations.Carlton *et al.* 2011, *ApJ*, 737, L22. Chandra expansion studies.Borkowski *et al.* 2013, *ApJ*, 771, L9. Chandra observations.Abramowski *et al.* 2014, *MNRAS*, 441, 790. HESS observations.Borkowski *et al.* 2014, *ApJ*, 790, L18. Chandra expansion studies.Roy & Pal 2014, *IAUS*, 296, 197. GMRT H_i observations.De Horta *et al.* 2014, *SerAJ*, 189, 41. ATCA at 1.4 (5''4×10''4), 2.4 (2''9×6''1) and 5 GHz (1''2×2''8).Gök & Ergin 2015, *AdSpR*, 56, 1793. Suzaku and Fermi observations.Zoglauer *et al.* 2015, *ApJ*, 798, 98. NuSTAR observations.Borkowski *et al.* 2017, *ApJ*, 837, L7. Chandra expansion studies.**G3.7-0.2****RA:** 17^h55^m26^s**Dec:** -25°50'**1-GHz flux/Jy:** 2.3**Spectral index:** 0.65**Size/arcmin:** 14×11**Type:** S

Has been called G003.8-00.3.

Radio: Double arc.**References:**Gray 1994, *MNRAS*, 270, 847. MOST at 843 MHz (43''×99'': S=2.4 Jy).Gaensler 1998, *ApJ*, 493, 781. VLA at 1.4 GHz (9''×15'': S=1.7±0.1 Jy).Yusef-Zadeh *et al.* 2004, *ApJS*, 155, 421. VLA at 1.4 GHz (8''4×11''4).

G3.8+0.3

RA: 17^h52^m55^s
Dec: -25°28'

1-GHz flux/Jy: 3?
Spectral index: 0.6

Size/arcmin: 18
Type: S?

Radio: Incomplete shell.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×100'':*S*=3.5 Jy).
 Bhatnagar 2002, MNRAS, 332, 1. GMRT at 327 MHz (17''×27'':*S*=6.0±0.4).

G4.2-3.5

RA: 18^h08^m55^s
Dec: -27°03'

1-GHz flux/Jy: 3.2?
Spectral index: 0.6?

Size/arcmin: 28
Type: S

Radio: Elongated shell.

Optical: Detected.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G4.5+6.8

RA: 17^h30^m42^s
Dec: -21°29'

1-GHz flux/Jy: 19
Spectral index: 0.64

Kepler, SN1604, 3C358

Size/arcmin: 3
Type: S

This is the remnant of Kepler's SN of AD1604.

Radio: Incomplete shell, brighter to the N.

Optical: Faint filaments.

X-ray: Shell, brighter to the N.

Distance: Optical expansion and proper motion indicates about 2.9 kpc, H α observations suggest 3.4 to 6.4 kpc.

References:

van den Bergh & Kamper 1977, ApJ, 218, 617. Optical proper motions.
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
 White & Long 1983, ApJ, 264, 196. Einstein observations.
 Matsui *et al.* 1984, ApJ, 287, 295. VLA at 1.4 (2''5×3''2) and 5 GHz (3''2×4''8) and Einstein image (5'').
 Dickel *et al.* 1988, ApJ, 330, 254. VLA at 1.4 (1''2×2''3) and 5 GHz (0''6×1''0) at two epochs.
 Smith *et al.* 1989, ApJ, 347, 925. EXOSAT observations.
 Hatsukade *et al.* 1990, PASJ, 42, 279. X-ray spectrum.
 Blair *et al.* 1991, ApJ, 366, 484. Optical imaging and spectroscopy.
 Bandiera & van den Bergh 1991, ApJ, 374, 186. Optical changes.
 van den Bergh 1991, PASP, 103, 194. Optical imaging.
 Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Hughes 1999, ApJ, 527, 298. ROSAT and Einstein image comparison for expansion studies.

Reynoso & Goss 1999, *AJ*, 118, 926. VLA at 1.4 GHz (13'' \times 23'') for H_I studies.
 Kinugasa & Tsunemi 1999, *PASJ*, 51, 239. ASCA observations.
 Gerardy & Fesen 2001, *AJ*, 121, 2781. IR spectroscopy and imaging.
 DeLaney *et al.* 2002, *ApJ*, 580, 914. VLA at 1.3 to 1.5 GHz and 5 GHz (7'' \times 2) for spectral index studies.
 Morgan *et al.* 2003, *ApJ*, 597, L33. Sub-mm dust observations.
 Sollerman *et al.* 2003, *A&A*, 407, 249. Optical spectroscopy.
 Cassam-Chenāi *et al.* 2004, *A&A*, 414, 545. XMM-Newton observations.
 Bamba *et al.* 2005, *ApJ*, 621, 793. Chandra observations of rim.
 Riesgo & López 2005, *RMxAA*, 41, 57. Optical observations of filament (previously classified as PN, H 2-12).
 Blair *et al.* 2007, *ApJ*, 662, 998. Spitzer observations.
 Reynolds *et al.* 2007, *ApJ*, 668, L135. Chandra observations.
 Sankrit *et al.* 2008, *AJ*, 135, 538. HST observations.
 Aharonian *et al.* 2008, *A&A*, 488, 219. HESS upper limit.
 Enomoto *et al.* 2008, *ApJ*, 683, 383. γ -ray upper limit.
 Katsuda *et al.* 2008, *ApJ*, 689, 225. Chandra proper motion studies.
 Vink *et al.* 2008, *ApJ*, 689, 231. Chandra proper motion studies.
 Gomez *et al.* 2012, *MNRAS*, 420, 3557. Herschel IR dust observations.
 Williams *et al.* 2012, *ApJ*, 755, 3. Spitzer spectroscopy.
 Burke *et al.* 2013, *ApJ*, 764, 63. Chandra observations.
 Yang *et al.* 2013, *ApJ*, 766, 44. Suzaku spectroscopy.
 Park *et al.* 2013, *ApJ*, 767, L10. Suzaku observations.
 Katsuda *et al.* 2015, *ApJ*, 808, 49. XMM-Newton, Chandra and Suzaku observations.
 Sankrit *et al.* 2016, *ApJ*, 817, 36. HST for proper motion studies.
 Koo *et al.* 2016, *ApJ*, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Sato & Hughes 2017, *ApJ*, 845, 167. Chandra expansion studies.
 Kasuga *et al.* 2018, *PASJ*, 70, 88. Chandra observations.

G4.8+6.2

RA: 17^h33^m25^s
Dec: -21°34'

1-GHz flux/Jy: 3
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G4.5+6.2.

Radio: Faint shell.

References:

Duncan *et al.* 1995, *MNRAS*, 277, 36. Parkes 64-m at 2.4-GHz (10'4).
 Bhatnagar 2000, *MNRAS*, 317, 453. GMRT at 327 MHz (1'3 \times 2'2 : S = 5.5 \pm 1.2 Jy), and NVSS at 1.4 GHz.
 Hewitt & Yusef-Zadeh 2009, *ApJ*, 694, L16. OH maser search.

G5.2-2.6

RA: 18^h07^m30^s
Dec: -25°45'

1-GHz flux/Jy: 2.6?
Spectral index: 0.6?

Size/arcmin: 18
Type: S

Radio: Poorly resolved shell.

References:

Reich *et al.* 1988, *IAUCo*, 101, 293. Summary of parameters.
 Reich *et al.* 1990, *A&AS*, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).

Hewitt & Yusef-Zadeh 2009, Apj, 694, L16. OH maser search.

G5.4–1.2

Milne 56

RA: 18^h02^m10^s
Dec: –24°54′

1-GHz flux/Jy: 35?
Spectral index: 0.2?

Size/arcmin: 35
Type: C?

Part been called G5.3–1.0. Has been suggested that this is not a SNR.

Radio: Incomplete shell, including wide ‘v’ of emission to east with small flat-spectrum source at apex.

Optical: Detected.

X-ray: Pulsar detected, with faint extension.

Point sources: Pulsar nearby, in flat spectrum source.

Distance: H_i absorption suggests > 4.3 kpc.

References:

- Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′:S=38 Jy).
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′).
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′.8:S=21.9±2.4 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.
 Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 and 5 GHz.
 Helfand & Becker 1985, Nature, 313, 118. Suggesting it is not a SNR.
 Manchester *et al.* 1985, MNRAS, 212, 975. Pulsar detection.
 Caswell *et al.* 1987, MNRAS, 225, 329. MOST at 843 MHz (42″×110″).
 Frail & Kulkarni 1991, Nature, 352, 785. Pulsar and remnant association.
 Manchester *et al.* 1991, MNRAS, 253, 7P. Pulsar and remnant association.
 Milne *et al.* 1992, MNRAS, 255, 707. Parkes 64-m at 4.75 (4′.5:S=30.8±2.1 Jy) and 8.4 GHz (3′:S=24±3 Jy), including polarisation.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.0×4′.9:S=38 Jy).
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz (68″×73″), plus H_i absorption.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gaensler & Frail 2000, Nature, 406, 158. Pulsar observations, including proper motion.
 Kaspi *et al.* 2001, Apj, 562, L163. X-ray detection of pulsar, and upper limit for remnant.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.
 Blazek *et al.* 2006, Apj, 652, 1523. Proper motion study of pulsar.
 Zeiger *et al.* 2008, Apj, 674, 271. Proper motion study of pulsar.
 Liszt *et al.* 2009, A&A, 508, 1331. CO and IR observations of region.
 Hewitt & Yusef-Zadeh 2009, Apj, 694, L16. OH maser detection.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.

G5.5+0.3

RA: 17^h57^m04^s
Dec: -24°00'

1-GHz flux/Jy: 5.5
Spectral index: 0.7

Size/arcmin: 15×12
Type: S

Has been called G5.55+0.32.

Radio: Shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=14.3\pm 0.3$ Jy), plus other observations.
 Liszt *et al.* 2009, A&A, 508, 1331. CO and IR observations of region.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G5.9+3.1

RA: 17^h47^m20^s
Dec: -22°16'

1-GHz flux/Jy: 3.3?
Spectral index: 0.4?

Size/arcmin: 20
Type: S

Radio: Asymmetric shell.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.1+0.5

RA: 17^h57^m29^s
Dec: -23°25'

1-GHz flux/Jy: 4.5
Spectral index: 0.9

Size/arcmin: 18×12
Type: S

Has been called G6.10+0.53.

Radio: Partial shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=13.4\pm 0.2$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G6.1+1.2

RA: 17^h54^m55^s
Dec: -23°05'

1-GHz flux/Jy: 4.0?
Spectral index: 0.3?

Size/arcmin: 30×26
Type: F

Has been called G6.1+1.15.

Radio: Faint, diffuse emission.

References:

Reich *et al.* 1988, IAUco, 101, 293. Summary of parameters.
 Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.4–0.1

W28

RA: 18^h00^m30^s**1-GHz flux/Jy:** 310**Size/arcmin:** 48**Dec:** –23°26'**Spectral index:** varies**Type:** C

Has been called G6.6–0.2.

Radio: Several non-thermal sources in a ring, with flat spectrum core.

Optical: Diffuse emission.

X-ray: Diffuse emission from most of the remnant.

Point sources: Young pulsar near edge of remnant, but not thought to be related.

Distance: Hi observations suggest 1.9 kpc.

References:

- Kundu & Velusamy 1972, A&A, 20, 237. NRAO 140-ft at 10 GHz (3').
 van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4) and 5 GHz (4'4).
 Dopita *et al.* 1977, ApJ, 214, 179. Some optical line ratios.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Lozinskaya 1981, SvAl, 7, 17. Mean optical velocity.
 Andrews *et al.* 1983, ApJ, 266, 684. VLA at 1.4 GHz (8'') 4.9 GHz (3'') and 15 GHz (1'') of central region only, plus Einstein image of central region.
 Bohigas *et al.* 1983, RMxAA, 8, 155. Optical spectra.
 Andrews *et al.* 1985, AJ, 90, 310. VLA of central component.
 Long *et al.* 1991, ApJ, 373, 567. Einstein and optical observations.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'1×4'4 : S=660 Jy).
 Kaspi *et al.* 1993, ApJ, 409, L57. Pulsar association.
 Frail *et al.* 1993, Nature, 365, 136. VLA at 327 MHz (smoothed to 65''), plus pulsar association.
 Frail *et al.* 1994, ApJ, 424, L111. VLA of associated OH masers.
 Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Claussen *et al.* 1997, ApJ, 489, 143. VLA of OH masers.
 Claussen *et al.* 1999, ApJ, 522, 349. High resolution observations of OH masers.
 Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz (2'6×5'5) and 1.48 GHz (40''×65'').
 Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz (52''×97'' : S = 425±40 Jy) and 1415 MHz (48''×88'' : S=246±20 Jy), and comparison with other observations.
 Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
 Douvion *et al.* 2001, A&A, 373, 281. ISO observations.
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
 Velázquez *et al.* 2002, AJ, 124, 2145. Parkes 64-m at 1.4 GHz (15') for Hi.
 Rho & Borkowski 2002, ApJ, 575, 201. ROSAT and ASCA observations.
 Claussen *et al.* 2002, ApJ, 580, 909. Observations of nearby source.
 Yusef-Zadeh *et al.* 2003, ApJ, 583, 267. OH observations.
 Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz of associated OH masers.
 Mavromatakis *et al.* 2004, A&A, 426, 567. Optical observations.
 Reach *et al.* 2005, ApJ, 618, 297. Molecular lines and near IR observations.
 Hoffman *et al.* 2005, ApJ, 620, 257. OH maser observations.

Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
 Aharonian *et al.* 2008, A&A, 481, 401. HESS observations.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Giuliani *et al.* 2010, A&A, 516, L11. γ -ray observations.
 Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.
 Abdo *et al.* 2010, ApJ, 718, 348. Fermi observations.
 Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
 Sawada *et al.* 2012, PASJ, 64, 81. Suzaku observations.
 Nichols *et al.* 2012, MNRAS, 419, 251. CO observations of selected regions.
 Gusdorf *et al.* 2012, A&A, 542, L19. CO observations of regions in NE.
 Vaupre e *et al.* 2014, A&A, 568, A50. CO, HCO⁺ and DCO⁺ molecular line observations.
 Pihlstr om *et al.* 2014, AJ, 147, 73. VLA observations of methanol masers.
 Neufeld *et al.* 2014, ApJ, 781, 102. Herschel and Spitzer IR spectroscopy.
 Hanabata *et al.* 2014, ApJ, 786, 145. Fermi observations of region.
 Zhou *et al.* 2014, ApJ, 791, 87. XMM-Newton observations.
 Gusdorf *et al.* 2014, IAUS, 296, 178. CO observations.
 Nakamura *et al.* 2014, PASJ, 66, 62. XMM-Newton observations of NE.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Maxted *et al.* 2016, MNRAS, 462, 532. NH₃ observations of region.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Pannuti *et al.* 2017, ApJ, 839, 59. Optical and X-ray observations.
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.
 Okon *et al.* 2018, PASJ, 70, 35. Suzaku observations.
 Nobukawa *et al.* 2018, ApJ, 854, 87. Suzaku observations.
 Cui *et al.* 2018, ApJ, 860, 69. Fermi observations.
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.
 Ruiz-Lapuente *et al.* 2018, ApJ, 862, 124. HST search for progenitor companion.

G6.4 + 4.0

RA: 17^h45^m10^s
Dec: -21°22'

1-GHz flux/Jy: 1.3?
Spectral index: 0.4?

Size/arcmin: 31
Type: S

Radio: Faint asymmetric shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Effelsberg 100-m at 2.7 GHz (4'3).
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G6.5 - 0.4

RA: 18^h02^m11^s
Dec: -23°34'

1-GHz flux/Jy: 27
Spectral index: 0.6

Size/arcmin: 18
Type: S

Has been called G6.51-0.48, and part has been called G6.67-0.42.

Radio: Shell, overlapping G6.4-0.1.

Optical: Detected.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 330 MHz (2'6×5'5) and 1.4 GHz (0'7×1'1).
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 60.8±0.4 Jy), plus other observations.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Ajello *et al.* 2016, ApJ, 819, 44. Fermi observations.

G7.0–0.1

RA: 18^h01^m50^s
Dec: –22°54′

1-GHz flux/Jy: 2.5?
Spectral index: 0.5?

Size/arcmin: 15
Type: S

Has been called G7.06–0.12.

Radio: Double rim, brightest in W, confused by bright H II region M20 in SE.

References:

Yusef-Zadeh *et al.* 2000, ApJ, 540, 842. VLA at 327 MHz (2′.6×5′.5) and 1.48 GHz (40″×65″).
 Dubner *et al.* 2000, AJ, 120, 1933. VLA at 328 MHz (52″×97″) and 1415 MHz (48″×88″).

G7.2+0.2

RA: 18^h01^m07^s
Dec: –22°38′

1-GHz flux/Jy: 2.8
Spectral index: 0.6

Size/arcmin: 12
Type: S

Has been called G7.20+0.20.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: S=5.2±0.2 Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G7.7–3.7

RA: 18^h17^m25^s
Dec: –24°04′

1-GHz flux/Jy: 11
Spectral index: 0.32

1814–24
Size/arcmin: 22
Type: S

Has been associated with the SN of AD386.

Radio: Shell, with high polarisation.

X-ray: Arc in S.

References:

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Milne *et al.* 1986, MNRAS, 223, 487. MOST at 843 MHz (44″×108″) and Parkes 64-m at 8.4 GHz (3′: S=4.6±0.5 Jy), with polarisation, plus review of flux densities.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (35″×70″: S=9.9±0.1 Jy), including polarisation.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Zhou *et al.* 2018, ApJ, 865, L6. XMM-Newton observations.

G8.3–0.0**RA:** 18^h04^m34^s**Dec:** –21°49′**1-GHz flux/Jy:** 1.2**Spectral index:** 0.6**Size/arcmin:** 5×4**Type:** S

Has been called G8.31–0.09.

Radio: Shell.**References:**Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=2.3\pm 0.1$ Jy), plus other observations.Higashi *et al.* 2008, ApJ, 683, 957. γ -ray detection.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.**G8.7–5.0****RA:** 18^h24^m10^s**Dec:** –23°48′**1-GHz flux/Jy:** 4.4**Spectral index:** 0.3**Size/arcmin:** 26**Type:** S**Radio:** Asymmetric shell.**References:**Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′3).
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.**G8.7–0.1****RA:** 18^h05^m30^s**Dec:** –21°26′**1-GHz flux/Jy:** 80**Spectral index:** 0.5**Size/arcmin:** 45**Type:** S?

(W30)

Has been called G8.6–0.1.

Radio: Clumpy non-thermal shell, with low-frequency turnover.**X-ray:** Northern edge detected.**Point sources:** Pulsar inside western edge.**References:**Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7′2×9′7: $S=190\pm 50$ Jy).

Kassim & Weiler 1990, Nature, 343, 146. VLA at 327 MHz (3′0×3′7).

Kassim & Weiler 1990, ApJ, 360, 184. VLA at 327 MHz (2′8×4′1: $S=129\pm 11$ Jy), and part at 1.4 GHz (0′9×1′8), plus review of flux densities.Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz (37″×55″).

Finley & Ögelman 1994, ApJ, 434, L25. ROSAT observations, including pulsar.

Aharonian *et al.* 2005, Science, 307, 1938. HESS detection.Aharonian *et al.* 2006, ApJ, 636, 777. HESS observations.Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.Briskin *et al.* 2006, ApJ, 652, 554. Pulsar proper motion.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.

Castro & Slane 2010, ApJ, 717, 372. Fermi observations.

Ajello *et al.* 2012, ApJ, 744, 80. Fermi observations.Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G8.9+0.4

RA: 18^h03^m58^s
Dec: -21°03'

1-GHz flux/Jy: 9
Spectral index: 0.6

Size/arcmin: 24
Type: S

Has been called G8.90+0.40.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=18.2\pm 0.5$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G9.7-0.0

RA: 18^h07^m22^s
Dec: -20°35'

1-GHz flux/Jy: 3.7
Spectral index: 0.6

Size/arcmin: 15×11
Type: S

Has been called G9.7-0.1 and G9.70-0.06.

Radio: Shell.

References:

Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': $S=6.5\pm 0.2$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser detection.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Yeung *et al.* 2016, ApJ, 827, 41. Fermi observations.

G9.8+0.6

RA: 18^h05^m08^s
Dec: -20°14'

1-GHz flux/Jy: 3.9
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Asymmetric shell.

References:

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Caswell 1983, MNRAS, 204, 833. Molonglo at 408 MHz (3': $S=5.8\pm 0.6$ Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41''×63'': $S=3.5\pm 0.4$ Jy).
 Frail *et al.* 1994, AJ, 107, 1120. VLA at 327 MHz.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G9.9-0.8

RA: 18^h10^m41^s
Dec: -20°43'

1-GHz flux/Jy: 6.7
Spectral index: 0.4

Size/arcmin: 12
Type: S

Has been called G9.95-0.81.

Radio: Shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 11.0 \pm 0.3$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. $H\alpha$ observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H_2 IR observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G10.5–0.0

RA: $18^{\text{h}}09^{\text{m}}08^{\text{s}}$
Dec: $-19^{\circ}47'$

1-GHz flux/Jy: 0.9
Spectral index: 0.6

Size/arcmin: 6
Type: S

Has been called G10.59–0.04.

Radio: Partial shell.

X-ray: Possibly detected.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 1.4 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G11.0–0.0

RA: $18^{\text{h}}10^{\text{m}}04^{\text{s}}$
Dec: $-19^{\circ}25'$

1-GHz flux/Jy: 1.3
Spectral index: 0.6

Size/arcmin: 11×9
Type: S

Has been called G11.0+0.0 and G11.03–0.05.

Radio: Partial shell.

X-ray: Diffuse emission.

Distance: Optical absorption suggests 2.4 kpc.

References:

Bamba *et al.* 2003, ApJ, 589, 253. ASCA observations.
 Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz ($25''$), 1.5 GHz ($25''$), and 74 MHz.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 3.1 \pm 0.2$ Jy), plus other observations.
 Castelletti *et al.* 2016, A&A, 587, A71. VLA at 1.4 GHz ($4''.4 \times 8''.3$) and CO observations of region.
 Araya *et al.* 2018, ApJ, 859, 69. Fermi observations.
 Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G11.1–1.0

RA: $18^{\text{h}}14^{\text{m}}03^{\text{s}}$
Dec: $-19^{\circ}46'$

1-GHz flux/Jy: 5.8
Spectral index: 0.5

Size/arcmin: 18×12
Type: S

Has been called G11.2–1.1 and G11.17–1.04.

Radio: Shell.

Optical: Detected.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 11.0 \pm 0.3$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. $H\alpha$ observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S=3.40±0.25 Jy) including polarisation and review of flux densities.

G11.1–0.7

RA: 18^h12^m46^s
Dec: –19°38'

1-GHz flux/Jy: 1.0
Spectral index: 0.7

Size/arcmin: 11×7
Type: S

Has been called G11.15–0.71.

Radio: Partial shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'':S=2.3±0.1 Jy), plus other observations.

G11.1+0.1

RA: 18^h09^m47^s
Dec: –19°12'

1-GHz flux/Jy: 2.3
Spectral index: 0.4

Size/arcmin: 12×10
Type: S

Has been called G11.18+0.11.

Radio: Shell.

References:

Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25''), 1.5 GHz (25''), and 74 MHz.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'':S=3.5±0.2 Jy), plus other observations.

Castelletti *et al.* 2016, A&A, 587, A71. VLA at 1.4 GHz (4''4×8''3) and CO observations of region.

G11.2–0.3

RA: 18^h11^m27^s
Dec: –19°25'

1-GHz flux/Jy: 22
Spectral index: 0.5

Size/arcmin: 4
Type: C

Probably associated with the SN of AD386.

Radio: Symmetrical clumpy shell, with flatter spectrum core.

X-ray: Shell, with hard spectrum centrally brightened region around pulsar.

Point sources: Central pulsar.

Distance: H_i absorption indicates 4.4 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Radhakrishnan *et al.* 1972, ApJS, 24, 49. H_i absorption.

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3').

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Downes 1984, MNRAS, 210, 845. VLA at 1465 MHz (20''×25'') and Einstein observations, with review of flux densities.

Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz, plus H_i absorption, Einstein observations.

Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26''5:S=4.04±0.24 Jy).

Green *et al.* 1988, MNRAS, 231, 735. VLA at 1.4 and 5 GHz.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'2×4'1:S=39 Jy).

Reynolds *et al.* 1994, MNRAS, 271, L1. ROSAT image and spectra.

Vasisht *et al.* 1996, ApJ, 456, L59. ASCA observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Torii *et al.* 1997, ApJ, 489, L145. ASCA detection of pulsar.
 Torii *et al.* 1999, ApJ, 523, L69. X-ray timing observations of pulsar.
 Kothes & Reich 2001, A&A, 372, 627. Effelsberg 100-m at 4.25, 10.45, 14.9 and 32 GHz (2′.5, 1′.1, 0′.86 and 0′.45 : S = 9.6±0.5, 6.3±0.4, 5.7±0.4 and 3.8±0.4).
 Kaspi *et al.* 2001, ApJ, 560, 371. Chandra observations.
 Tam *et al.* 2002, ApJ, 572, 202. VLA at 1.4/1.5 GHz (1′′8×2′′6 : S = 16.6±0.9 Jy) and 5 GHz (1′′5×2′′1 : S = 8.4±0.9 Jy) for spectral studies.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 14.7 GHz.
 Roberts *et al.* 2003, ApJ, 588, 992. Chandra observations.
 Tam & Roberts 2003, ApJ, 598, L27. Multi-epoch VLA observations at 1.4/1.5 GHz and 5 GHz, for expansion studies.
 Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25′′), 1.5 GHz (25′′), and 74 MHz.
 Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (18′′).
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.
 Koo *et al.* 2007, ApJ, 657, 308. IR observations.
 Dean *et al.* 2008, MNRAS, 384, L29. INTEGRAL observations of pulsar and nebula.
 Moon *et al.* 2009, ApJ, 703, L81. IR spectroscopy.
 Mizuno *et al.* 2010, AJ, 139, 1542. Spitzer observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 9.0±0.5 Jy) including polarisation and review of flux densities.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Lee *et al.* 2013, ApJ, 770, 143. IR observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Borkowski *et al.* 2016, ApJ, 819, 160. Chandra observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

G11.4–0.1

RA: 18^h10^m47^s
Dec: –19°05′

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 8
Type: S?

Radio: Incomplete shell, possibly with central core.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : S = 9.4 Jy) contaminated by sidelobes of a nearby source, and Parkes 64-m at 5 GHz (4′ : S = 2.8 Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′.8 : S = 2.0±0.4 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.2×4′.1 : S = 18 Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (44′′×63′′ : S = 5.1±0.6 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Brogan *et al.* 2004, AJ, 127, 355. VLA at 330 MHz (25′′), 1.5 GHz (25′′), and 74 MHz.
 Rangelov *et al.* 2014, ApJ, 796, 34. X-ray upper limit.
 Castelletti *et al.* 2016, A&A, 587, A71. VLA at 1.4 GHz (4′.4×8′.3) and CO observations of region.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G11.8–0.2**RA:** 18^h12^m25^s**Dec:** –18°44′**1-GHz flux/Jy:** 0.7**Spectral index:** 0.3**Size/arcmin:** 4**Type:** S

Has been called G11.89–0.21.

Radio: Shell.

X-ray: Possibly detected.

References:

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=0.9\pm 0.1$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.0–0.1**RA:** 18^h12^m11^s**Dec:** –18°37′**1-GHz flux/Jy:** 3.5**Spectral index:** 0.7**Size/arcmin:** 7?**Type:** ?

Radio: Incomplete shell, defined in E only.

X-ray: Detected, including possible PWN.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=6.6$ Jy) and Parkes 64-m at 5 GHz (4′: $S=1.1$ Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′2×4′1).

Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (41″×61″: $S=0.7$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Yamauchi *et al.* 2014, PASJ, 66, 20. Suzaku observations of possible PWN.

G12.2+0.3**RA:** 18^h11^m17^s**Dec:** –18°10′**1-GHz flux/Jy:** 0.8**Spectral index:** 0.7**Size/arcmin:** 6×5**Type:** S

Has been called G12.26+0.30.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″: $S=1.5\pm 0.1$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G12.5+0.2**RA:** 18^h12^m14^s**Dec:** -17°55'**1-GHz flux/Jy:** 0.6**Spectral index:** 0.4**Size/arcmin:** 6×5**Type:** C?

Has been called G12.58+0.22.

Radio: Diffuse, central brightened.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 0.8±0.1 Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.7-0.0**RA:** 18^h13^m19^s**Dec:** -17°54'**1-GHz flux/Jy:** 0.8**Spectral index:** 0.8**Size/arcmin:** 6**Type:** S

Has been called G12.72-0.00.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 2.0±0.1 Jy), plus other observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G12.8-0.0**RA:** 18^h13^m37^s**Dec:** -17°49'**1-GHz flux/Jy:** 0.8**Spectral index:** 0.5**Size/arcmin:** 3**Type:** C?

Has been called G12.82-0.02 and G12.83-0.02.

Radio: Shell.

X-ray: Diffuse.

Point sources: Central X-ray pulsar.

References:

Brogan *et al.* 2005, ApJ, 629, L105. VLA at 330 MHz (19'' × 32''), plus other observations.
Ubertini *et al.* 2005, ApJ, 629, L109. INTEGRAL and other observations.
Aharonian *et al.* 2006, ApJ, 636, 777. HESS detection.
Albert *et al.* 2006, ApJ, 637, L41. γ -ray observations.
Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 1.2±0.1 Jy), plus other observations.
Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
Funk *et al.* 2007, A&A, 470, 249. XMM-Newton observations, CO observations of surroundings.
Helfand *et al.* 2007, ApJ, 665, 1297. Chandra observations.
Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
Gotthelf & Halpern 2009, ApJ, 700, L158. Pulsar detection.
Halpern *et al.* 2012, ApJ, 753, L14. Pulsar observations.
Araya *et al.* 2018, ApJ, 859, 69. Fermi observations.
Dzib *et al.* 2018, ApJ, 866, 100. VLA of central source.

G13.3–1.3**RA:** 18^h19^m20^s**Dec:** –18°00′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 70×40**Type:** S?**Radio:** Amorphous emission.**Optical:** Filaments in S.**X-ray:** Elongated emission.**Distance:** Absorption indicates 2--4 kpc.**References:**Seward *et al.* 1995, ApJ, 449, 681. ROSAT detection, optical studies and observations of CO.Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.**G13.5+0.2****RA:** 18^h14^m14^s**Dec:** –17°12′**1-GHz flux/Jy:** 3.5?**Spectral index:** 1.0?**Size/arcmin:** 5×4**Type:** S

Has been called G13.46+0.16.

Radio: Elongated, incomplete shell.**References:**Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ($S = 0.65 \pm 0.05$ Jy) and 1.4 GHz ($15'' : S = 2.67 \pm 0.5$ Jy).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.**G14.1–0.1****RA:** 18^h16^m40^s**Dec:** –16°41′**1-GHz flux/Jy:** 0.5**Spectral index:** 0.6**Size/arcmin:** 6×5**Type:** S

Has been called G14.18–0.12.

Radio: Shell.**References:**Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.9 \pm 0.1$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G14.3+0.1**RA:** 18^h15^m58^s**Dec:** –16°27′**1-GHz flux/Jy:** 0.6**Spectral index:** 0.4**Size/arcmin:** 5×4**Type:** S

Has been called G14.30+0.14. Has been suggested this is not an SNR.

Radio: Partial shell.**References:**Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 1.2 \pm 0.1$ Jy), plus other observations.

Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Pinheiro Gonçalves *et al.* 2011, AJ, 142, 47. IR observations.

G15.1 – 1.6

RA: 18^h24^m00^s
Dec: –16°34′

1-GHz flux/Jy: 5.5?
Spectral index: 0.0?

Size/arcmin: 30×24
Type: S?

Radio: Elongated, incomplete shell.

Optical: Diffuse shell.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).
 Boumis *et al.* 2008, A&A, 481, 705. Optical detection.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 4.8±0.3 Jy) including polarisation and review of flux densities.

G15.4 + 0.1

RA: 18^h18^m02^s
Dec: –15°27′

1-GHz flux/Jy: 5.6
Spectral index: 0.62

Size/arcmin: 15×14
Type: C?

Has been called G15.42+0.18.

Radio: Shell.

X-ray: Centrally brightened.

Distance: H α observations suggest 4.8 kpc.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42′′ : S = 10.9±0.3 Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 2.1±0.2 Jy) including polarisation and review of flux densities.
 Castelletti *et al.* 2013, A&A, 557, L15. GMRT at 1.4 GHz (15′′) plus SGPS H α and CO observations of region.
 Abramowski *et al.* 2014, A&A, 562, A40. HESS and XMM-Newton observations.
 Supan *et al.* 2015, A&A, 576, A81. GMRT at 624 MHz (10′′).
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.

G15.9 + 0.2

RA: 18^h18^m52^s
Dec: –15°02′

1-GHz flux/Jy: 5.0
Spectral index: 0.63

Size/arcmin: 7×5
Type: S?

Radio: Incomplete shell, with bright concentration to the E.

X-ray: Shell, brighter to S and E.

Point sources: Central X-ray source.

References:

Clark *et al.* 1975, *AujPA*, 37, 1. Molonglo at 408 MHz (3': $S=7.7$ Jy) and Parkes 64-m at 5 GHz (4': $S=1.9$ Jy).
 Angerhofer *et al.* 1977, *A&A*, 55, 11. NRAO 140-ft at 5 GHz (6'8: $S=1.1\pm 0.2$ Jy).
 Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Caswell *et al.* 1982, *MNRAS*, 200, 1143. FIRST at 1415 MHz (44'' \times 58'').
 Dubner *et al.* 1996, *AJ*, 111, 1304. VLA at 330 MHz (61'' \times 77'': $S=11.2\pm 1.0$ Jy), 1.4 GHz (14'' \times 23'': $S=3.9\pm 0.1$ Jy) and 4.9 GHz (13'' \times 16'').
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Reynolds *et al.* 2006, *ApJ*, 652, L45. Chandra observations.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9'5: $S=2.0\pm 0.3$ Jy) including polarisation and review of flux densities.
 Klochov *et al.* 2016, *A&A*, 592, L12. Chandra observations.
 Koo *et al.* 2016, *ApJ*, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Maggi & Acero 2017, *A&A*, 597, A65. XMM-Newton observations.
 Sasaki *et al.* 2018, *MNRAS*, 479, 3033. Chandra and IR observations, plus H α non-detection.

G16.0–0.5

RA: 18^h21^m56^s
Dec: –15°14'

1-GHz flux/Jy: 2.7
Spectral index: 0.6

Size/arcmin: 15 \times 10
Type: S

Has been called G16.05–0.57.

Radio: Shell.

References:

Brogan *et al.* 2006, *ApJ*, 639, L25. VLA at 330 MHz (42'': $S=4.9\pm 0.2$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, *ApJ*, 694, L16. OH maser search.
 Beaumont *et al.* 2011, *ApJ*, 741, 14. CO of region.
 Froebrich *et al.* 2015, *MNRAS*, 454, 2586. H₂ IR observations.

G16.2–2.7

RA: 18^h29^m40^s
Dec: –16°08'

1-GHz flux/Jy: 2.5
Spectral index: 0.4

Size/arcmin: 17
Type: S

Radio: Double rim.

References:

Trushkin 1999, *A&A*, 352, L103. Review of radio observations.
 Hewitt & Yusef-Zadeh 2009, *ApJ*, 694, L16. OH maser search.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9'5: $S=1.28\pm 0.10$ Jy) including polarisation and review of flux densities.

G16.4–0.5

RA: 18^h22^m38^s
Dec: –14°55'

1-GHz flux/Jy: 4.6
Spectral index: 0.3?

Size/arcmin: 13
Type: S

Has been called G16.41–0.55.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 10.0 \pm 0.3$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 3.0 \pm 0.3$ Jy) including polarisation and review of flux densities.

G16.7+0.1

RA: 18^h20^m56^s
Dec: $-14^{\circ}20'$

1-GHz flux/Jy: 3.0
Spectral index: 0.6

Size/arcmin: 4
Type: C

Has been called G16.73+0.08.

Radio: Asymmetric shell with flat-spectrum core.

X-ray: Non-thermal core.

References:

Helfand *et al.* 1989, ApJ, 341, 151. VLA at 5 GHz ($5 : S = 0.95$ Jy) and 1.4 GHz ($15'' : S = 2.43$ Jy) and Ooty at 327 MHz ($S = 5.13$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.
 Reynoso & Mangum 2000, ApJ, 545, 874. CO observations of surroundings.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Helfand *et al.* 2003, ApJ, 592, 941. XMM-Newton observations.
 Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz ($19'' \times 25''$).
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 6 GHz.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 1.23 \pm 0.11$ Jy) including polarisation and review of flux densities.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Chang *et al.* 2018, MNRAS, 474, 2607. Chandra observations.

G17.0–0.0

RA: 18^h21^m57^s
Dec: $-14^{\circ}08'$

1-GHz flux/Jy: 0.5
Spectral index: 0.5

Size/arcmin: 5
Type: S

Has been called G17.02–0.04.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz ($42'' : S = 0.7 \pm 0.1$ Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.

G17.4–2.3

RA: 18^h30^m55^s
Dec: $-14^{\circ}52'$

1-GHz flux/Jy: 5
Spectral index: 0.5?

Size/arcmin: 24?
Type: S

Radio: Incomplete, poorly defined shell.

Optical: Filaments to SE, and diffuse emission.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.

Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2002, A&A, 385, 1042. Optical observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 2.3±0.2 Jy) including polarisation and review of flux densities.

G17.4–0.1

RA: 18^h23^m08^s
Dec: –13°46'

1-GHz flux/Jy: 0.4
Spectral index: 0.7

Size/arcmin: 6
Type: S

Has been called G17.48–0.12.

Radio: Partial shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 0.9±0.1 Jy), plus other observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G17.8–2.6

RA: 18^h32^m50^s
Dec: –14°39'

1-GHz flux/Jy: 5
Spectral index: 0.5

Size/arcmin: 24
Type: S

Radio: Well defined shell.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4'3).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 2.23±0.13 Jy) including polarisation and review of flux densities.

G18.1–0.1

RA: 18^h24^m34^s
Dec: –13°11'

1-GHz flux/Jy: 4.6
Spectral index: 0.5

Size/arcmin: 8
Type: S

Has been called G18.1–0.2 and G18.16–0.16.

Radio: Shell.

X-ray: Possibly detected.

Distance: H_I absorption suggests 6.4 kpc.

References:

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (8').
 Kassim *et al.* 1989, ApJ, 338, 152. VLA at 330 MHz (2'8×3'2) and 1.5 GHz (0'9×1'5).
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'' : S = 7.6±0.1 Jy), plus other observations.
 Hewitt & Yusef-Zadeh 2009, ApJ, 694, L16. OH maser search.
 Paron *et al.* 2013, MNRAS, 433, 1619. CO, optical and other observations of region.
 Leahy *et al.* 2014, MNRAS, 438, 1813. VGPS H_I observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

Voison *et al.* 2016, MNRAS, 458, 2813. Molecular line observations of region.
Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_I absorption observations.

G18.6–0.2

RA: 18^h25^m55^s
Dec: –12°50′

1-GHz flux/Jy: 1.4
Spectral index: 0.4

Size/arcmin: 6
Type: S

Has been called G18.62–0.28.

Radio: Partial shell.

Distance: H_I absorption suggests 4.4 kpc.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42″:S=1.9±0.1 Jy), plus other observations.
Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
Voison *et al.* 2016, MNRAS, 458, 2813. Molecular line observations of region.
Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_I absorption observations.

G18.8+0.3

Kes 67

RA: 18^h23^m58^s
Dec: –12°23′

1-GHz flux/Jy: 33
Spectral index: 0.46

Size/arcmin: 17×11
Type: S

Has been called G18.9+0.3.

Radio: Incomplete shell, in complex region near the H_{II} region W39.

Distance: Association with molecular cloud suggests 12 kpc, and H_I absorption suggests 13.8 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′:S=17±7 Jy).
Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′:S=38 Jy).
Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.
Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
Milne *et al.* 1989, PASA, 8, 187. Parkes 64-m at 8.4 GHz (3′.0:S=12.9±1.0 Jy), including polarisation.
Kassim 1992, AJ, 103, 943. VLA at 327 MHz (2′.9×3′.5:S=55 Jy).
Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (55″×75″:S=29.9±0.3 Jy).
Dubner *et al.* 1999, AJ, 118, 930. Parkes 64-m at 1.6 GHz (15″) for H_I, VLA at 1.6 GHz (12″×17″) for OH, plus CO observations.
Dubner *et al.* 2004, A&A, 426, 201. CO observations of environment.
Tian *et al.* 2007, A&A, 474, 541. VGPS at 1.4 GHz (1′) including H_I, plus CO observations of region.
Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5:S=15.3±0.9 Jy) including polarisation and review of flux densities.
Vasquez *et al.* 2012, A&A, 545, A89. CO observations of region.
Paron *et al.* 2012, A&A, 547, A60. CO and other molecular observations of region.
Paron *et al.* 2015, A&A, 580, A51. CO observations in S.
Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_I absorption observations.

G18.9–1.1**RA:** 18^h29^m50^s**Dec:** –12°58′**1-GHz flux/Jy:** 37**Spectral index:** 0.39**Size/arcmin:** 33**Type:** C?

Has been called G18.95–1.1 and G18.94–1.04.

Radio: Non-thermal, diffuse partially limb-brightened, with central ridge.

Optical: Detected.

X-ray: Partial shell.

Point sources: Compact X-ray source, with diffuse nebula.

Distance: Optical absorption suggests 1.8 kpc.

References:

Fürst *et al.* 1985, *Nature*, 314, 720. Effelsberg 100-m at 4.75 GHz (2′.4 : S = 23.8 Jy), plus other flux densities.

Odegard 1986, *AJ*, 92, 1372. TPT at 57.5 MHz (7′.2 × 8′ : S = 82 ± 15 Jy), plus review of flux densities. Barnes & Turtle 1988, *IAUCo*, 101, 347. Molonglo at 408 MHz (2′.9 × 3′.1 : S = 58 ± 9 Jy) and Parkes 64-m at 5 GHz (4′.1 × 4′.4 : S = 23 ± 6 Jy).

Patnaik *et al.* 1988, *Nature*, 332, 136. Ooty at 327 MHz (0′.6 × 1′.6).

Fürst *et al.* 1989, *A&A*, 209, 361. Effelsberg 100-m at 4.75 GHz (2′.45 : S = 23.8 Jy) and VLA at 1.5 and 4.9 GHz (14″ × 19″), and Effelsberg 100-m at 1.4 GHz (9′) for Hi.

Aschenbach *et al.* 1991, *A&A*, 246, L32. ROSAT observations.

Fürst *et al.* 1997, *A&A*, 319, 655. ROSAT observations, and Effelsberg 100-m at 10.6 GHz (1′.1).

Reich 2002, in *NSPS*, p1. Effelsberg 100-m at 10.6 GHz, including polarisation.

Harrus *et al.* 2004, *ApJ*, 603, 152. ROSAT and ASCA observations.

Tüllmann *et al.* 2010, *ApJ*, 720, 848. Chandra detection of compact source.

Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 19.6 ± 1.0 Jy) including polarisation and review of flux densities.

Stupar & Parker 2011, *MNRAS*, 414, 2282. H α observations.

Froebrich *et al.* 2015, *MNRAS*, 454, 2586. H₂ IR observations.

Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.

Shan *et al.* 2018, *ApJS*, 238, 35. Optical absorption for distance.

G19.1+0.2**RA:** 18^h24^m56^s**Dec:** –12°07′**1-GHz flux/Jy:** 10**Spectral index:** 0.5**Size/arcmin:** 27**Type:** S

Has been called G19.15+0.27.

Radio: Partial shell.

References:

Brogan *et al.* 2006, *ApJ*, 639, L25. VLA at 330 MHz (42″ : S = 17.4 ± 0.4 Jy), plus other observations.

G20.0–0.2

RA: 18^h28^m07^s
Dec: –11°35′

1-GHz flux/Jy: 10
Spectral index: 0.1

Size/arcmin: 10
Type: F

Radio: Faint, filled-centre, polarised.

X-ray: Centrally brightened.

Point sources: OH source 20.1–0.1 is nearby.

Distance: H_i absorption suggests 11.2 kpc.

References:

Becker & Helfand 1985, ApJ, 297, L25. VLA at 1.4 and 5 GHz (12'').

Odegard 1986, AJ, 92, 1372. TPT at 57.5 MHz (7'2×8': S=8.5±2 Jy), plus review of flux densities.

Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5: S=9.2±0.5 Jy) including polarisation and review of flux densities.

Petriella *et al.* 2013, A&A, 554, A73. Chandra observations, plus CO and H_i observations of region.

Acerio *et al.* 2016, ApJS, 224, 8. Fermi observations.

Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G21.0–0.4

RA: 18^h31^m12^s
Dec: –10°47′

1-GHz flux/Jy: 1.1
Spectral index: 0.6

Size/arcmin: 9×7
Type: S

Has been called G21.04–0.47.

Radio: Shell.

References:

Brogan *et al.* 2006, ApJ, 639, L25. VLA at 330 MHz (42'': S=2.3±0.2 Jy), plus other observations.

G21.5–0.9

RA: 18^h33^m33^s
Dec: –10°35′

1-GHz flux/Jy: 7
Spectral index: varies

Size/arcmin: 5
Type: C

Early observations relate to the central core only.

Radio: Filled-centre, with high frequency turnover.

X-ray: Central core, with extended, faint halo.

Point sources: Central pulsar.

Distance: H_i absorption suggests 4.4 kpc.

References:

Wilson & Weiler 1976, A&A, 53, 89. WSRT at 5 GHz (6''×35'').

Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (5''×20'') and 8 GHz (2''×7''), plus review of flux densities.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Becker & Szymkowiak 1981, ApJ, 248, L23. VLA at 5 GHz (8''), and Einstein observations.

Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT X-ray spectrum, and VLA H_i observations.

Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (26'': S=5.64±0.29 Jy).

Fürst *et al.* 1988, PASJ, 40, 347. NRO array at 22.3 GHz (4''4×7''3).

Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4'3), including polarisation.
 Salter *et al.* 1989, A&A, 225, 167. Observations at 90.7 (29''5 : $S = 3.8 \pm 0.4$ Jy) and 141.9 GHz ($S = 2.5 \pm 1.2$ Jy).
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz ($S = 3.94 \pm 0.70$ Jy), plus review of flux densities.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'3×3'4 : $S = 9$ Jy).
 Wallace *et al.* 1994, A&A, 286, 565. H α of surroundings.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Slane *et al.* 2000, ApJ, 533, L29. Chandra observations identifying X-ray halo.
 Warwick *et al.* 2001, A&A, 365, L248. XMM-Newton observations of X-ray halo.
 Bock *et al.* 2001, ApJ, 561, L203. BIMA at 94 GHz (4''6×8''6).
 Safi-Harb *et al.* 2001, ApJ, 561, 308. Chandra and other X-ray observations.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 22 GHz (8'') and 32 GHz, including polarisation.
 La Palombara & Mereghetti 2002, A&A, 383, 916. XMM-Newton upper limit on pulsations.
 Bocchino *et al.* 2005, A&A, 442, 539. XMM-Newton and Chandra observations.
 Gupta *et al.* 2005, CSci, 89, 853. Pulsar discovery.
 Camilo *et al.* 2006, ApJ, 637, 456. Pulsar discovery.
 Bietenholz & Bartel 2008, MNRAS, 386, 1411. VLA at 4.75 GHz (0''53×0''82) for expansion studies.
 Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1') including H α .
 Matheson & Safi-Harb 2010, ApJ, 724, 572. Chandra observations.
 Bietenholz *et al.* 2011, MNRAS, 412, 1221. VLA at 1.4 GHz (14''×18'').
 Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 6 GHz.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : $S = 6.5 \pm 0.4$ Jy) including polarisation and review of flux densities.
 Zajczyk *et al.* 2012, A&A, 542, A12. IR observations, including polarisation.
 Nynka *et al.* 2014, ApJ, 789, 72. NuSTAR observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 70, 100 and 143 GHz.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H α absorption observations.
 Aharonian *et al.* 2018, PASJ, 70, 38. Hitomi observations.

G21.6–0.8

RA: 18^h33^m40^s
Dec: –10°25'

1-GHz flux/Jy: 1.4
Spectral index: 0.5?

Size/arcmin: 13
Type: S

Has been called G21.64–0.84.

Radio: Faint, irregular shell.

References:

Bietenholz *et al.* 2011, MNRAS, 412, 1221. VLA at 327 MHz (85'' : $S = 2.8$ Jy) and 1.4 GHz (14''×18'').
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H $_2$ IR observations.

G21.8–0.6

Kes 69

RA: 18^h32^m45^s**1-GHz flux/Jy:** 65**Size/arcmin:** 20**Dec:** –10°08′**Spectral index:** 0.56**Type:** S**Radio:** Incomplete shell.**X-ray:** Detected.**Distance:** Association with CO indicates 5.2 kpc, and H_I absorption suggests 5.6 kpc.**References:**

- Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Wilson 1972, A&A, 19, 354. H₂CO absorption.
 Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 5 GHz (6′) and 10 GHz (3′).
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′ : S=42.3±4.6 Jy).
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.3×3′.5 : S=132 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including masers.
 Yusef-Zadeh *et al.* 2003, ApJ, 585, 319. X-ray observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Wood *et al.* 2008, AJ, 135, 2358. VLA at 4.8 GHz, including polarisation.
 Tian & Leahy 2008, MNRAS, 391, L54. VGPS at 1.4 GHz (1′) including H_I.
 Zhou *et al.* 2009, ApJ, 691, 516. CO and HCO⁺ of region.
 Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S=24.0±1.3 Jy) including polarisation and review of flux densities.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_I absorption observations.
 Sezer *et al.* 2018, MNRAS, 481, 1416. Suzaku observations.

G22.7–0.2**RA:** 18^h33^m15^s**1-GHz flux/Jy:** 33**Size/arcmin:** 26**Dec:** –09°13′**Spectral index:** 0.6**Type:** S?**Radio:** Non-thermal ring in complex region, overlapping G23.3–0.3.**X-ray:** Possible detection.**Point sources:** Variable radio source near centre, and γ -ray source near edge.**Distance:** Association with CO indicates 4.4 kpc, and H_I absorption suggests 4.7 kpc.**References:**

- Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.3×3′.4 : S=82 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Becker *et al.* 2010, AJ, 140, 157. Variable radio source detection.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Su *et al.* 2014, ApJ, 796, 122. CO observations.
 Su *et al.* 2015, ApJ, 811, 134. CO observations of region.
 Abramowski *et al.* 2015, MNRAS, 446, 1163. HESS observations.
 Mori *et al.* 2017, ApJ, 848, 80. NuSTAR, Chandra and XMM-Newton observations.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G23.3–0.3

W41

RA: 18^h34^m45^s
Dec: –08°48′

1-GHz flux/Jy: 70
Spectral index: 0.5

Size/arcmin: 27
Type: S

Radio: Distorted ring, in complex region, overlapping G22.7–0.2.

X-ray: Possible extended emission, with compact sources.

Point sources: Pulsar association suggested.

Distance: H_i and CO observations suggest 4.2 or 4.8 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.3×3′.4: S=138 Jy).
 Gaensler & Johnston 1995, MNRAS, 275, L73. Possible pulsar association.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Aharonian *et al.* 2005, Science, 307, 1938. HESS detection.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Aharonian *et al.* 2006, ApJ, 636, 777. HESS observations.
 Albert *et al.* 2006, ApJ, 643, L53. γ -ray observations.
 Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.
 Tian *et al.* 2007, ApJ, 657, L25. VGPS at 1.4 GHz (1′) including H_i, plus XMM-Newton observations.
 Leahy & Tian 2008, AJ, 135, 167. VGPS at 1.4 GHz (1′) including H_i, plus CO observations.
 Mukherjee *et al.* 2009, ApJ, 691, 1707. XMM-Newton and SWIFT observations.
 Frail *et al.* 2013, ApJ, 773, L19. OH observations.
 Castro *et al.* 2013, ApJ, 774, 36. Fermi observations.
 Abramowski *et al.* 2015, A&A, 574, A27. HESS observations.
 Su *et al.* 2015, ApJ, 811, 134. CO observations of region.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G24.7–0.6

RA: 18^h38^m43^s
Dec: –07°32′

1-GHz flux/Jy: 8
Spectral index: 0.5

Size/arcmin: 15?
Type: S?

Radio: Incomplete shell, defined in SW.

Distance: H_i absorption and CO suggests 3.8 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: S=12.3 Jy) and Parkes 64-m at 5 GHz (4′: S=3.6 Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′.8: S=2.2±0.5 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).

Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz ($44'' \times 56''$: $S = 1.9$ Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Ranasinghe & Leahy 2018, MNRAS, 477, 2243. VGPS H_I absorption, plus CO observations.

G24.7 + 0.6

RA: 18^h34^m10^s
Dec: -07°05′

1-GHz flux/Jy: 20?
Spectral index: 0.2?

Size/arcmin: 30×15
Type: C?

Radio: Filled-centre, with faint shell, and a compact H_{II} region to the S.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz (4′.3 : $S = 19 \pm 3$ Jy) and 4.75 GHz (2′.4 : $S = 17 \pm 4$ Jy) and NRO 45-m at 10.2 GHz (2′.7 : $S = 15 \pm 3$ Jy).
 Becker & Helfand 1987, ApJ, 316, 660. VLA at 1.4 GHz (12′′), and X-ray upper limit.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G25.1 – 2.3

RA: 18^h45^m10^s
Dec: -08°00′

1-GHz flux/Jy: 8
Spectral index: 0.5?

Size/arcmin: 80×30?
Type: S

Radio: Incomplete shell, extent not well defined.

References:

Gao *et al.* 2011, A&A, 532, A144. Urumqi 25-m at 5 GHz (9′.5 : $S = 3.7 \pm 0.4$ Jy), plus other observations.

G27.4 + 0.0

RA: 18^h41^m19^s
Dec: -04°56′

1-GHz flux/Jy: 6
Spectral index: 0.68

4C–04.71
Size/arcmin: 4
Type: S

Early references refer to G27.3–0.1 (Kes 73), a supposed larger remnant.

Radio: Incomplete shell.

X-ray: Diffuse emission, with central low period pulsar.

Point sources: Central AXP.

Distance: H_I absorption suggests 7.5 to 9.8 kpc or 5.8 kpc, and association with CO suggests 9 kpc.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′).
 Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′ : $S = 4.4$ Jy).
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′.8 : $S = 2.0 \pm 0.5$ Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).

Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1415 MHz ($45'' \times 60''$: $S = 3.5$ Jy). Revise $S_{408 \text{ MHz}} = 10.4$ Jy, and $S_{5 \text{ GHz}} = 1.9 \pm 0.2$ Jy.

Kriss *et al.* 1985, ApJ, 288, 703. Einstein observations, plus VLA at 1.4 and 5 GHz ($12''$).

Sanbonmatsu & Helfand 1992, AJ, 104, 2189. VLA at 1.4 GHz for H_i absorption.

Helfand *et al.* 1994, ApJ, 434, 627. ROSAT observations, particularly of central source.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations, including nearby masers.

Vasisht & Gotthelf 1997, ApJ, 486, L129. ASCA detection of pulsar.

Gotthelf & Vasisht 1997, ApJ, 486, L133. ASCA observations.

Gotthelf *et al.* 1999, ApJ, 522, L49. X-ray timing observations of pulsar.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Mereghetti *et al.* 2001, MNRAS, 321, 143. Search for optical/IR counterpart to pulsar.

Kuiper *et al.* 2004, ApJ, 613, 1173. X-ray observations of pulsar.

Tian & Leahy 2008, ApJ, 677, 292. VGPS at 1.4 GHz ($1'$) including H_i.

Mizuno *et al.* 2010, AJ, 139, 1542. Spitzer observations.

An *et al.* 2013, ApJ, 779, 163. NuSTAR and other observations.

Kumar *et al.* 2014, ApJ, 781, 41. Chandra and XMM-Newton observations.

Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).

Yeung *et al.* 2017, ApJ, 837, 69. Fermi observations.

Borkowski & Reynolds 2017, ApJ, 846, 13. Chandra expansion studies.

Liu *et al.* 2017, ApJ, 851, 37. Fermi and CO observations.

Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G27.8+0.6

RA: 18^h39^m50^s
Dec: $-04^{\circ}24'$

1-GHz flux/Jy: 30
Spectral index: varies

Size/arcmin: 50×30
Type: F

Radio: Filled-centre, with spectral turnover.

X-ray: Possible pulsar wind nebula.

References:

Reich *et al.* 1984, A&A, 133, L4. Effelsberg 100-m at 2.7 GHz ($4'.3$: $S = 23 \pm 2$ Jy), and 4.75 GHz ($2'.4$: $S = 18 \pm 2$ Jy) and NRO 45-m at 10.2 GHz (smoothed to $4'.3$: $S = 8.5 \pm 2$ Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

Misanovic *et al.* 2010, ApJ, 725, 931. XMM-Newton pulsar/wind nebula search.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'.5$: $S = 21.0 \pm 1.1$ Jy) including polarisation and review of flux densities.

Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G28.6–0.1**RA:** 18^h43^m55^s**Dec:** –03°53′**1-GHz flux/Jy:** 3?**Spectral index:** ?**Size/arcmin:** 13×9**Type:** S**Radio:** Poorly defined regions of non-thermal emission.**X-ray:** Diffuse shell, with thermal and non-thermal emission.**Point sources:** Pulsar in NE.**Distance:** H_i absorption and CO suggests 9.6 kpc.**References:**Helfand *et al.* 1989, ApJ, 341, 151. VLA at 1.4 GHz (15'') and 5 GHz (15'').Bamba *et al.* 2001, PASJ, 53, L21. ASCA observations.Ueno *et al.* 2003, ApJ, 588, 338. Chandra observations.Ebisawa *et al.* 2005, ApJ, 635, 214. Chandra observations.Zyuzin *et al.* 2018, MNRAS, 476, 2177. Pulsar observations.Ranasinghe & Leahy 2018, MNRAS, 477, 2243. VGPS H_i absorption, plus CO observations.**G28.8+1.5****RA:** 18^h39^m00^s**Dec:** –02°55′**1-GHz flux/Jy:** ?**Spectral index:** 0.4?**Size/arcmin:** 100?**Type:** S?**Radio:** Part of rim detected.**X-ray:** Diffuse, Centrally brightened.**References:**

Schwentker 1994, A&A, 286, L47. ROSAT observations.

Song *et al.* 2000, PASJ, 52, 181. ASCA observations.Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.Misanovic *et al.* 2010, ApJ, 725, 931. XMM-Newton pulsar/wind nebula search.Froeblich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.**G29.6+0.1****RA:** 18^h44^m52^s**Dec:** –02°57′**1-GHz flux/Jy:** 1.5?**Spectral index:** 0.5?**Size/arcmin:** 5**Type:** S**Radio:** Diffuse shell.**Point sources:** AXP associated.**References:**Gaensler *et al.* 1999, ApJ, 526, L37. VLA at 5 GHz (13'') and 8 GHz (8'')Vasisht *et al.* 2000, ApJ, 542, L49. X-ray observations of AXP.Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.

G29.7–0.3

Kes 75

RA: 18^h46^m25^s**1-GHz flux/Jy:** 10**Size/arcmin:** 3**Dec:** –02°59′**Spectral index:** 0.63**Type:** C

Has erroneously been called G29.6+0.1.

Radio: Shell with flatter spectrum emission from centre.

X-ray: Thermal shell and non-thermal core, and central pulsar.

Point sources: X-ray pulsar.

Distance: Association with CO implies 11 kpc, and H α absorption suggests 5.6 kpc.

References:

- Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (8′.4 : S = 5±20% Jy).
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′).
 Becker & Kundu 1976, ApJ, 204, 427. NRAO interferometer at 2.7 GHz (7″×20″) and 8 GHz (8″×25″), plus review of flux densities.
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).
 Becker *et al.* 1983, ApJ, 268, L93. VLA at 1.4 GHz (3″), plus Einstein observations.
 Becker & Helfand 1984, ApJ, 283, 154. VLA at 5 GHz (2′6), plus H α .
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (26′.5 : S = 1.02±0.07 Jy).
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.4×3′.7 : S = 27.4 Jy).
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Blanton & Helfand 1996, ApJ, 470, 961. ASCA observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Gotthelf *et al.* 2000, ApJ, 542, L37. X-ray pulsar detection.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Mereghetti *et al.* 2002, ApJ, 574, 873. BeppoSAX observations of pulsar.
 Helfand *et al.* 2003, ApJ, 582, 783. Chandra observations.
 Bock & Gaensler 2005, ApJ, 626, 343. BIMA at 88.6 GHz (10″×13″).
 Morton *et al.* 2007, ApJ, 667, 219. Spitzer and Chandra observations.
 McBride *et al.* 2008, A&A, 477, 249. INTEGRAL and Chandra observations.
 Leahy & Tian 2008, A&A, 480, L25. VGPS at 1.4 GHz (1′) including H α .
 Kumar & Safi-Harb 2008, ApJ, 678, L43. Chandra observations.
 Ng *et al.* 2008, ApJ, 686, 508. Chandra observations.
 Gavriil *et al.* 2008, Science, 319, 1802. X-ray observations of pulsar.
 Su *et al.* 2009, ApJ, 694, 376. CO observations of region, plus Chandra observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 3.6±0.6 Jy) including polarisation and review of flux densities.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H α absorption observations.
 Reynolds *et al.* 2018, ApJ, 856, 133. Chandra observations for expansion and brightness change studies.

G30.7–2.0**RA:** 18^h54^m25^s**Dec:** –02°54′**1-GHz flux/Jy:** 0.5?**Spectral index:** 0.7?**Size/arcmin:** 16**Type:** ?**Radio:** Poorly defined.**References:**Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G30.7+1.0****RA:** 18^h44^m00^s**Dec:** –01°32′**1-GHz flux/Jy:** 6**Spectral index:** 0.4**Size/arcmin:** 24×18**Type:** S?**Radio:** Non-thermal, highly polarised part shell?**Point sources:** Compact source near centre.**References:**Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2′.4 : S = 3.4±0.4 Jy), plus other flux densities.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.6×3′.9 : S = 8.6 Jy).

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 2.93±0.19 Jy) including polarisation and review of flux densities.**G31.5–0.6****RA:** 18^h51^m10^s**Dec:** –01°31′**1-GHz flux/Jy:** 2?**Spectral index:** ?**Size/arcmin:** 18?**Type:** S?

Has been called G31.55–0.65.

Radio: Distorted shell? near H_{II} region.**Optical:** Diffuse, incomplete shell.**References:**Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2′.4), plus other flux densities.Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.**G31.9+0.0****RA:** 18^h49^m25^s**Dec:** –00°55′**1-GHz flux/Jy:** 25**Spectral index:** varies**Size/arcmin:** 7×5**Type:** S

3C391

Radio: Shell, brightest in NW, with low frequency turnover.**X-ray:** Diffuse with central core.**Distance:** H_I absorption and CO association suggests 7.1 kpc.**References:**Radhakrishnan *et al.* 1972, ApJS, 24, 49. H_I absorption.

Becker & Kundu 1975, *AJ*, 80, 679. NRAO 140-ft at 10.6 GHz (3').
 Green *et al.* 1975, *A&A*, 44, 187. Effelsberg 100-m at 15 GHz (58'': $S > 1.6$).
 Becker & Kundu 1976, *ApJ*, 204, 427. NRAO interferometer at 2.7 GHz (20'' \times 23'') and 8 GHz (9'' \times 24''), plus review of flux densities.
 Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2'.6).
 Goss *et al.* 1979, *A&A*, 78, 75. FIRST at 1.4 GHz (54'' \times 66'': $S = 20 \pm 2$ Jy) and Effelsberg 100-m at 10.7 GHz (77'': $S = 7.5 \pm 0.8$ Jy).
 Wang & Seward 1984, *ApJ*, 279, 705. Einstein observations.
 Kassim 1992, *AJ*, 103, 943. VLA at 327 MHz (3'.4 \times 3'.7: $S = 44.8$ Jy).
 Reynolds & Moffett 1993, *AJ*, 105, 2226. VLA at 1.4 GHz (6''), including possible associated CO.
 Moffett & Reynolds 1994, *ApJ*, 425, 668. VLA at 330 MHz (smoothed to 30'': $S = 38.5 \pm 0.5$ Jy) 1.46 GHz (6'.7) and 4.85 GHz (6'.2 \times 6'.4), including spectral index and polarisation studies.
 Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant, including masers.
 Gorham *et al.* 1996, *ApJ*, 458, 257. Pulsar search.
 Reach & Rho 1996, *A&A*, 315, L277. ISO spectroscopy.
 Rho & Petre 1996, *ApJ*, 467, 698. ROSAT observations.
 Wilner *et al.* 1998, *AJ*, 115, 247. CO observations of surroundings.
 Reach & Rho 1998, *ApJ*, 507, L93. ISO observations.
 Reach & Rho 1999, *ApJ*, 511, 836. CO, HCO⁺ and CS observations of surroundings.
 Reach & Rho 2000, *ApJ*, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, *ApJ*, 558, 943. Erratum.
 Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
 Chen & Slane 2001, *ApJ*, 563, 202. ASCA observations.
 Reach *et al.* 2002, *ApJ*, 564, 302. Observations of shocked molecular species.
 Chen *et al.* 2004, *ApJ*, 616, 885. Chandra observations.
 Brogan *et al.* 2005, *AJ*, 130, 148. VLA at 74 MHz, 330 MHz and 1.5 GHz (70'').
 Kawasaki *et al.* 2005, *ApJ*, 631, 935. ASCA observations.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Neufeld *et al.* 2007, *ApJ*, 664, 890. Spitzer observations.
 Hewitt *et al.* 2008, *ApJ*, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Castro & Slane 2010, *ApJ*, 717, 372. Fermi observations.
 Yuan & Neufeld 2011, *ApJ*, 726, 76. Spitzer observations.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9'.5: $S = 8.9 \pm 0.6$ Jy) including polarisation and review of flux densities.
 Pihlström *et al.* 2014, *AJ*, 147, 73. VLA search for methanol masers.
 Neufeld *et al.* 2014, *ApJ*, 781, 102. Spitzer and Herschel IR spectroscopy.
 Ergin *et al.* 2014, *ApJ*, 790, 65. Fermi and Suzaku observations.
 Gusdorf *et al.* 2014, *IAUS*, 296, 178. CO observations.
 Su *et al.* 2014, *IAUS*, 296, 372. VGPS for H_I absorption.
 Lee *et al.* 2014, *MNRAS*, 443, 2650. [FeII] IR survey observations.
 Sato *et al.* 2014, *PASJ*, 66, 124. Suzaku observations.
 Froebrich *et al.* 2015, *MNRAS*, 454, 2586. H₂ IR observations.
 Kilpatrick *et al.* 2016, *ApJ*, 816, 1. CO observations, including broad lines.
 Koo *et al.* 2016, *ApJ*, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Ranasinghe & Leahy 2017, *ApJ*, 843, 119. H_I and CO observations.

G32.0–4.9

3C396.1

RA: 19^h06^m00^s**1-GHz flux/Jy:** 22?**Size/arcmin:** 60?**Dec:** –03°00'**Spectral index:** 0.5?**Type:** S?**Radio:** Possible large shell?**References:**

Milne & Hill 1969, *AJPh*, 22, 211. Parkes 64-m at 635 MHz ($31' : S = 25 \pm 30\%$ Jy), 1410 MHz ($15' : S = 19 \pm 15\%$ Jy) and 2650 MHz ($8.4' : S = 8.6 \pm 30\%$ Jy). Fluxes if size is $60'$, plus review of flux densities.
 Caswell 1970, *AJPh*, 23, 105. Revision of low frequency flux densities.
 Dickel & DeNoyer 1975, *AJ*, 80, 437. Arecibo at 111 MHz ($S = 105 \pm 30$ Jy).

G32.1–0.9

RA: $18^{\text{h}}53^{\text{m}}10^{\text{s}}$
Dec: $-01^{\circ}08'$

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 40?
Type: C?

Radio: Possible faint shell, not well defined.

X-ray: Diffuse, with clumps.

References:

Folgheraiter *et al.* 1997, *MNRAS*, 292, 365. ROSAT and ASCA observations.
 Froebrich *et al.* 2015, *MNRAS*, 454, 2586. H_2 IR observations.

G32.4+0.1

RA: $18^{\text{h}}50^{\text{m}}05^{\text{s}}$
Dec: $-00^{\circ}25'$

1-GHz flux/Jy: 0.25?
Spectral index: ?

Size/arcmin: 6
Type: S

Has been called G32.45+0.1.

Radio: Shell.

X-ray: Shell.

Distance: X-ray absorption suggests 17 kpc.

References:

Yamaguchi *et al.* 2004, *PASJ*, 56, 1059. XMM-Newton and other observations.
 Ueno *et al.* 2005, in *XRRRC*, E4.18. XMM-Newton observations.
 Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
 Kilpatrick *et al.* 2016, *ApJ*, 816, 1. CO observations, including broad lines.

G32.8–0.1

Kes 78

RA: $18^{\text{h}}51^{\text{m}}25^{\text{s}}$
Dec: $-00^{\circ}08'$

1-GHz flux/Jy: 11?
Spectral index: 0.2?

Size/arcmin: 22×15
Type: S?

Part has been called G33.1–0.1.

Radio: Incomplete, elongated shell.

Optical: Detected.

X-ray: Patchy, elongated shell.

Distance: Association with CO and H_I absorption indicate 4.8 kpc.

References:

Velusamy & Kundu 1974, *A&A*, 32, 375. NRAO 300-ft et 2.7 GHz ($5' : S = 7.2 \pm 0.5$ Jy).
 Dickel & DeNoyer 1975, *AJ*, 80, 437. Arecibo at 430 MHz ($S = 19.0 \pm 15.5$ Jy).
 Becker & Kundu 1975, *AJ*, 80, 679. NRAO 140-ft at 10.6 GHz ($3'$).
 Caswell *et al.* 1975, *AJPA*, 37, 39. Molonglo at 408 MHz ($3' : S = 12.8$ Jy) and Parkes 64-m at 5 GHz ($4' : S = 7.7$ Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'6×3'8: S=31.3 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Zhou & Chen 2011, ApJ, 743, 4. XMM-Newton observations, plus CO of region.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H $_2$ IR observations.
 Bamba *et al.* 2016, ApJ, 818, 63. Suzaku observations.
 Miceli *et al.* 2017, A&A, 599, A45. XMM-Newton observations.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H i absorption observations.

G33.2–0.6

RA: 18^h53^m50^s
Dec: –00°02'

1-GHz flux/Jy: 3.5
Spectral index: varies

Size/arcmin: 18
Type: S

Radio: Incomplete shell.

Optical: Filaments and diffuse emission.

References:

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Reich 1982, A&A, 106, 314. Effelsberg 100-m at 2.7 GHz (4'4: S=2.6±0.3 Jy) and 4.75 GHz (2'5: S=1.75±0.2 Jy).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (52''×68'' : S=2.7±0.3 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Boumis *et al.* 2009, A&A, 499, 789. Optical observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H $_2$ IR observations.

G33.6+0.1

RA: 18^h52^m48^s
Dec: +00°41'

1-GHz flux/Jy: 20
Spectral index: 0.51

Kes 79, 4C00.70, HC13

Size/arcmin: 10
Type: S

Has been called G33.7+0.0.

Radio: Shell, with bright central region, in complex region.

X-ray: Multiple shells and filaments.

Point sources: Central X-ray pulsar.

Distance: H i absorption suggests 7.8 kpc or 3.5 kpc, and CO observations suggest 5.5 kpc.

References:

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3' : S=35.5 Jy) and Parkes 64-m at 5 GHz (4' : S=7.8 Jy).
 Caswell *et al.* 1975, A&A, 45, 239. H i absorption.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S $_{430}$ MHz = 69±33 Jy.
 Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3' : S=6.8±1.5 Jy).
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6'8: S=11.4±1.1 Jy).
 Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1'), plus observations of the nearby point source.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H i absorption to nearby point source, possibly extragalactic.

Sequist & Gilmore 1982, *AJ*, 87, 378. VLA observations of nearby source, plus Einstein observations.

Green 1989, *MNRAS*, 238, 737. OH absorption.

Frail & Clifton 1989, *ApJ*, 336, 854. VLA at 1.4 GHz ($1' \times 2'9$), including H α absorption.

Velusamy *et al.* 1991, *AJ*, 102, 676. VLA at 327 MHz ($1'$), 1.5 ($7'' \times 14''$) and 5 GHz ($7''$), including spectral comparison.

Green & Dewdney 1992, *MNRAS*, 254, 686. Observations of adjacent molecular material.

Kassim 1992, *AJ*, 103, 943. VLA at 327 MHz ($3'6 \times 3'8$: $S = 34.8$ Jy).

Seward & Velusamy 1995, *ApJ*, 439, 715. ROSAT observations.

Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.

Koralesky *et al.* 1998, *AJ*, 116, 1323. VLA search for OH emission.

Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.

Tsunemi & Enoguchi 2002, *PASJ*, 54, 735. ASCA observations.

Seward *et al.* 2003, *ApJ*, 584, 414. Chandra observations.

Stanimirović 2003, *ApJ*, 592, 953. Arecibo OH absorption.

Sun *et al.* 2004, *ApJ*, 605, 742. Chandra observations.

Gotthelf *et al.* 2005, *ApJ*, 627, 390. XMM-Newton pulsar detection.

Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer possible detection.

Halpern *et al.* 2007, *ApJ*, 665, 1304. X-ray pulsar timing observations.

Giacani *et al.* 2009, *A&A*, 507, 841. VLA at 74 MHz ($36'' \times 39''$: $S = 76 \pm 10$ Jy), 324 MHz ($13''$: $S = 39 \pm 8$ Jy) and 1.5 GHz ($17'' \times 19''$: $S = 11.5 \pm 1.5$ Jy), plus review of flux densities and XMM-Newton observations.

Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz ($9'5$: $S = 9.4 \pm 0.5$ Jy) including polarisation and review of flux densities.

Auchettl *et al.* 2014, *ApJ*, 783, 32. Fermi detection.

Kilpatrick *et al.* 2016, *ApJ*, 816, 1. CO observations, including broad lines.

Zhou *et al.* 2016, *ApJ*, 831, 192. CO and other observations.

Sato *et al.* 2016, *PASJ*, 68, S8. Suzaku observations.

Ranasinghe & Leahy 2018, *AJ*, 155, 204. VGPS H α absorption observations.

Kuriki *et al.* 2018, *ApJ*, 864, 161. CO observations.

G34.7–0.4

W44, 3C392

RA: 18^h56^m00^s**1-GHz flux/Jy:** 240**Size/arcmin:** 35×27**Dec:** +01°22'**Spectral index:** 0.37**Type:** C

Has been called G34.6–0.5.

Radio: Distorted shell, brighter to the E, with pulsar and associated nebula.

Optical: Diffuse emission.

X-ray: Centrally concentrated, thermal spectrum, plus pulsar wind nebula.

Point sources: Pulsar within the boundary of the remnant.

Distance: H α absorption suggests 3.0 kpc, and optical absorption suggests 2.1 kpc.

References:

Caswell *et al.* 1975, *A&A*, 45, 239. H α absorption.

Clark *et al.* 1975, *AJPA*, 37, 75. Molonglo at 408 MHz ($3'$: $S = 299$ Jy).

Dickel & DeNoyer 1975, *AJ*, 80, 437. Arecibo $S_{430 \text{ MHz}} = 540 \pm 187$ Jy.

Dickel & Milne 1976, *AJPh*, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'4$) and 5 GHz ($4'4$).

Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz ($2'6$).

Smith *et al.* 1985, *MNRAS*, 217, 99. Einstein observations.

Wolszczan *et al.* 1991, *ApJ*, 372, L99. Pulsar detection.

Kassim 1992, *AJ*, 103, 943. VLA at 327 MHz ($3'6 \times 3'8$: $S = 469$ Jy).

- Jones *et al.* 1993, MNRAS, 265, 631. VLA at 1.4 GHz (15''), plus X-ray spectra.
- Rho *et al.* 1994, ApJ, 430, 757. Optical and ROSAT observations.
- Koo & Heiles 1995, ApJ, 442, 679. H_I of surrounding shell.
- Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ -ray emission.
- Harrus *et al.* 1996, ApJ, 464, L161. ASCA observations.
- Frail *et al.* 1996, ApJ, 464, L165. VLA at 1.5 and 8.4 GHz (7'' \times 8'' \times 9'') of pulsar nebula.
- Frail *et al.* 1996, AJ, 111, 1651. OH maser emission.
- Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
- Reach & Rho 1996, A&A, 315, L277. ISO spectroscopy.
- Harrus *et al.* 1997, ApJ, 488, 781. X-ray observations.
- Claussen *et al.* 1997, ApJ, 489, 143. VLA of associated OH masers.
- Giacani *et al.* 1997, AJ, 113, 1379. VLA at 1.4 GHz (15''), plus optical images.
- Seta *et al.* 1998, ApJ, 505, 286. CO observations of surroundings.
- Cox *et al.* 1999, ApJ, 524, 179. Revision of distance.
- Reach & Rho 2000, ApJ, 544, 843. ISO observations of interactions with surroundings.
see also: Reach & Rho 2001, ApJ, 558, 943. Erratum.
- Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
- Petre *et al.* 2002, ApJ, 579, 404. Chandra observations of pulsar and wind nebula.
- Mavromatakis *et al.* 2003, A&A, 405, 591. Optical observations.
- Shelton *et al.* 2004, ApJ, 611, 906. Chandra observations of part.
- Reach *et al.* 2005, ApJ, 618, 297. Molecular line and near-IR observations.
- Hoffman *et al.* 2005, ApJ, 627, 803. Observations of OH masers.
- Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
- Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
- Neufeld *et al.* 2007, ApJ, 664, 890. Spitzer observations.
- Castelletti *et al.* 2007, A&A, 471, 537. VLA at 74 MHz (36'' \times 39'' : $S = 634 \pm 70$ Jy) and 324 MHz (13'' : $S = 411 \pm 50$ Jy).
- Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
- Abdo *et al.* 2010, Science, 327, 1103. Fermi observations.
- Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'.5 : $S = 118 \pm 6$ Jy) including polarisation and review of flux densities.
- Giuliani *et al.* 2011, ApJ, 742, L30. γ -ray observations.
- Uchiyama *et al.* 2012, ApJ, 749, L35. Fermi observations.
- Uchida *et al.* 2012, PASJ, 64, 141. Suzaku observations.
- Ackermann *et al.* 2013, Science, 339, 807. Fermi observations.
- Yoshiike *et al.* 2013, ApJ, 768, 179. CO and H_I observations of region.
- Sashida *et al.* 2013, ApJ, 774, 10. HCO⁺ and CO observations of region.
- Park *et al.* 2013, ApJ, 777, 14. Arecibo H_I observations of region.
- Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
- Anderl *et al.* 2014, A&A, 569, A81. CO observations of regions in NE.
- Su *et al.* 2014, IAUS, 296, 372. VGPS for H_I absorption.
- Froeblich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
- Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30, 44 and 70 GHz.
- Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
- McEwen *et al.* 2016, ApJ, 826, 189. NH₃ and CH₃OH observations.
- Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
- Génova-Santos *et al.* 2017, MNRAS, 464, 4107. Radio observations at 10 to 20 GHz.
- Egron *et al.* 2017, MNRAS, 470, 1329. SRT at 1.5 GHz (11' : $S = 214 \pm 6$ Jy) and 7 GHz (2'.7 : $S = 94 \pm 4$ Jy).
- Yamada *et al.* 2017, ApJ, 834, L3. CO and HCO⁺ observations.
- Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.
- Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_I absorption observations.

G35.6–0.4**RA:** 18^h57^m55^s**Dec:** +02°13′**1-GHz flux/Jy:** 9
Spectral index: 0.5**Size/arcmin:** 15×11
Type: S?

Re-identified as SNR in 2009.

Radio: Diffuse, with some limb brightening.**Distance:** H_I absorption suggests 3.8 kpc.**References:**

Green 2009, MNRAS, 399, 177. Identification in the radio as a SNR.
 Paron & Giacani 2010, A&A, 509, L4. CO and IR observations of region.
 Zhu *et al.* 2013, ApJ, 775, 95. H_I and other observations.
 Paredes *et al.* 2014, A&A, 561, A56. GMRT at 610 MHz (4′′.8×12′′.2).
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_I absorption observations.

G36.6–0.7**RA:** 19^h00^m35^s**Dec:** +02°56′**1-GHz flux/Jy:** 1.0
Spectral index: 0.7?**Size/arcmin:** 25?
Type: S?**Radio:** polarised arc, possibly part of a larger shell?**References:**

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2′.4), plus other flux densities.
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.4×3′.7 : S=6.7 Jy).
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S=0.39±0.04 Jy) including polarisation and review of flux densities.

G36.6+2.6**RA:** 18^h48^m49^s**Dec:** +04°26′**1-GHz flux/Jy:** 0.7?
Spectral index: 0.5?**Size/arcmin:** 17×13?
Type: S**Radio:** Poorly resolved shell.**References:**

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).

G38.7–1.3**RA:** 19^h06^m40^s**Dec:** +04°28′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 32×19?**Type:** S

G38.7–1.4 refers to the E portion.

Radio: Incomplete shell.**Optical:** Arc of filaments, brighter to E.**X-ray:** Detected in E.**References:**Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT of E, and radio survey observations.Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.Huang *et al.* 2014, ApJ, 785, 118. XMM-Newton and Chandra observations.Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.**G39.2–0.3****RA:** 19^h04^m08^s**Dec:** +05°28′**1-GHz flux/Jy:** 18**Spectral index:** 0.34

3C396, HC24, NRAO 593

Size/arcmin: 8×6**Type:** C**Radio:** Shell, brighter to W, with faint ‘tail’ to E.**X-ray:** Diffuse, brighter to W, with central core.**Point sources:** Central X-ray source.**Distance:** H_I absorption suggests 8.5 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S₄₃₀ MHz = 54±38 Jy.

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′:S=4.1±1.0 Jy).

Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (48″×65″).

Becker & Helfand 1987, AJ, 94, 1629. VLA at 1.4 GHz (12″:S=14 Jy) and 5 GHz, plus Einstein observations.

Patnaik *et al.* 1990, A&A, 232, 467. VLA at 1.5 GHz (25″) and 1.4 GHz (7″5×7″8) and 5 GHz (25″) including polarisation, plus Ooty at 327 MHz (31″×100″), including review of flux densities.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′5×3′6:S=42.5 Jy).

Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz for spectral index studies.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

Harrus & Slane 1999, ApJ, 516, 811. ASCA observations.

Aharonian *et al.* 2001, A&A, 375, 1008. HESS limit.Olbert *et al.* 2003, ApJ, 592, L45. Chandra observations.Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.Scaife *et al.* 2007, MNRAS, 377, L69. 33 GHz observations.Lee *et al.* 2009, ApJ, 691, 1042. IR observations.Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5:S=8.8±0.5 Jy) including polarisation and review of flux densities.Su *et al.* 2011, ApJ, 727, 43. Chandra and CO observations of region.

Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Cruciani *et al.* 2016, MNRAS, 459, 4224. Parkes 64 m at 8.4, 13.5, 18.6 and 21.5 GHz, plus review of flux densities.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G39.7–2.0

W50, SS433

RA: 19^h12^m20^s**1-GHz flux/Jy:** 85?**Size/arcmin:** 120×60**Dec:** +04°55′**Spectral index:** 0.7?**Type:** ?

Eastern part has been called G40.0–3.1. Is this a SNR?

Radio: Elongated shell, containing SS433, adjacent to the H_{II} region S74.

Optical: Faint filaments at the edge of the radio emission.

X-ray: Emission from SS433 and two lobes.

Point sources: SS433 is the compact source in the centre of the W50.

Distance: H_i absorption and CO observations indicate 4.9 kpc.

References:

van den Bergh 1980, ApJ, 236, L23. Optical in H α and [S_{II}].
 Zealey *et al.* 1980, MNRAS, 192, 731. Optical spectra.
 van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H_i absorption to nearby point source (not SS433).
 Seaquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
 Watson *et al.* 1983, ApJ, 273, 688. X-ray observations.
 Downes *et al.* 1986, MNRAS, 218, 393. Effelsberg 100-m at 4.75 GHz (2′.4 : S = 34±4 Jy), plus previous 1.7 and 2.7 GHz data.
 Romney *et al.* 1987, ApJ, 321, 822. VLBI of SS433, including distance.
 Elston & Baum 1987, AJ, 94, 1633. Mosaic with VLA at 1.4 GHz (30′′) of fine structure only.
 Kawai *et al.* 1989, PASJ, 41, 491. X-ray observations of SS433.
 Band 1989, ApJ, 336, 937. Einstein and EXOSAT observations.
 Yamauchi *et al.* 1994, PASJ, 46, L109. X-ray spectral observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Brinkmann *et al.* 1996, A&A, 312, 306. ROSAT observations.
 Kotani *et al.* 1996, PASJ, 48, 619. X-ray line emission from SS433.
 Safi-Harb & Ögelman 1997, ApJ, 483, 868. ROSAT observations.
 Dubney *et al.* 1998, AJ, 116, 1842. VLA at 328 MHz (60′′×64′′ : S = 160±20 Jy), and 1.4 GHz (54′′×56′′), plus NRAO 140-ft at 1.4 GHz (21′) for H_i observations.
 Safi-Harb & Petre 1999, ApJ, 512, 784. X-ray observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. HESS limit.
 Brinkmann *et al.* 2007, A&A, 463, 611. XMM-Newton observations of E lobe.
 Boumis *et al.* 2007, MNRAS, 381, 308. Optical observations.
 Lockman *et al.* 2007, MNRAS, 381, 881. H_i observations.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 37±4 Jy), including polarisation and review of flux densities.
 Farnes *et al.* 2017, MNRAS, 467, 4777. ATCA at 1.4 to 3.1 GHz, including polarisation and H α observations.
 Broderick *et al.* 2018, MNRAS, 475, 5360. LOFAR at 115 to 189 MHz, including 140 MHz (55′′×78′′).
 Su *et al.* 2018, ApJ, 863, 103. CO and H_i observations.

G40.5–0.5

RA: 19^h07^m10^s
Dec: +06°31′

1-GHz flux/Jy: 11
Spectral index: 0.4

Size/arcmin: 22
Type: S

Radio: Shell, brightest to the NE.

Point sources: Central pulsar.

References:

- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Downes *et al.* 1980, A&A, 92, 47. Effelsberg 100-m at 1.7 GHz (7′.6 : S = 9.3 ± 1.3 Jy), and 2.7 GHz (4′.4 : S = 7.2 ± 0.5 Jy), plus review of flux densities.
 Aharonian *et al.* 2001, A&A, 375, 1008. HESS limit.
 Yang *et al.* 2006, ChJAA, 6, 210. CO observations of surroundings.
 Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 6.4 ± 0.3 Jy) including polarisation and review of flux densities.
 Aliu *et al.* 2014, ApJ, 787, 166. γ -ray observations of region.
 Lyne *et al.* 2017, ApJ, 834, 137. Pulsar detection.

G41.1–0.3

3C397

RA: 19^h07^m34^s
Dec: +07°08′

1-GHz flux/Jy: 25
Spectral index: 0.50

Size/arcmin: 4.5 × 2.5
Type: S

Radio: 3C397 is two sources: the E is the SNR, the W is a H_{II} region.

X-ray: Brighter to the E and W, with central component.

Distance: H_I absorption suggest 8.5 kpc.

References:

- Kundu *et al.* 1974, AJ, 79, 132. NRAO 140-ft at 10 GHz (3′) and 5 GHz (6′).
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′ : S = 21.3 ± 1.2 Jy for both components).
 Caswell *et al.* 1975, A&A, 45, 239. H_I absorption.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S_{430 MHz} = 82 ± 51 Jy, also Algonquin 46-m at 10.6 GHz (3′ : S = 12 ± 2 Jy), and Haystack 36-m at 15.5 GHz (2′.3 : S = 8.5 ± 3.0 Jy).
 Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′ : S = 29.8 Jy) and Parkes 64-m at 5 GHz (4′ : S = 8.7 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Caswell *et al.* 1982, MNRAS, 200, 1143. FIRST at 1.4 GHz (52″ × 58″).
 Becker *et al.* 1985, ApJ, 296, 461. VLA at 1.4 and 5 GHz (8″), plus Einstein observations.
 Morsi & Reich 1987, A&AS, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30″ : S = 4.10 ± 0.19 Jy).
 Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3′.5 × 3′.6 : S = 46.3 Jy).
 Anderson & Rudnick 1993, ApJ, 408, 514. VLA at 1.45 and 4.89 GHz, for spectral index studies.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Chen *et al.* 1999, ApJ, 520, 737. ASCA and ROSAT observations.
 Dyer & Reynolds 1999, ApJ, 526, 365. VLA at 1.5 GHz (6′.6 × 6′.9) and 4.8 GHz (5′.6 × 6′.4), including polarisation and comparison with ROSAT image.
 Safi-Harb *et al.* 2000, ApJ, 545, 922. ROSAT, ASCA and other X-ray observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Aharonian *et al.* 2001, A&A, 375, 1008. HESS limit.
 Safi-Harb *et al.* 2005, ApJ, 618, 321. Chandra observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Jiang *et al.* 2010, ApJ, 712, 1147. CO observations of region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S=18.5±1.1 Jy) including polarisation and review of flux densities.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
 Yamaguchi *et al.* 2015, ApJ, 801, L31. Suzaku observations.
 Clark *et al.* 2015, ApJ, 809, L2. Fermi observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Leahy & Ranasinghe 2016, ApJ, 817, 74. VGPS for H_i absorption.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G41.5+0.4

RA: 19^h05^m50^s
Dec: +07°46'

1-GHz flux/Jy: 1?
Spectral index: ?

Size/arcmin: 10
Type: S?

Radio: Partial clumpy shell, brighter to NE.

References:

Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 332 MHz (20'':S=1.8±0.4 Jy).
 Alves *et al.* 2012, MNRAS, 422, 2429. Radio observations.

G42.0-0.1

RA: 19^h08^m10^s
Dec: +08°00'

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 8
Type: S?

Radio: Irregular shell.

References:

Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 332 MHz (20'':S=1.8±0.4 Jy).
 Alves *et al.* 2012, MNRAS, 422, 2429. Radio observations.

G42.8+0.6

RA: 19^h07^m20^s
Dec: +09°05'

1-GHz flux/Jy: 3?
Spectral index: 0.5?

Size/arcmin: 24
Type: S

Has been called G42.8+0.65.

Radio: Faint shell.

Point sources: Near soft gamma repeater, and young pulsar.

References:

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2'4":S=1.5±0.2 Jy), plus other flux densities.
 Vasisht *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3'2×3'4).
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.
 Lorimer & Xilouris 2000, ApJ, 545, 385. Pulsar detection.
 Aharonian *et al.* 2001, A&A, 375, 1008. HESS limit.
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50''), and other observations of the region.

G43.3–0.2

W49B

RA: 19^h11^m08^s
Dec: +09°06′**1-GHz flux/Jy:** 38
Spectral index: 0.46**Size/arcmin:** 4×3
Type: S**Radio:** Shell, brightest to the SE and W, near the H_{II} region W49A.**X-ray:** Centrally brightened, elongated E--W.**Point sources:** Compact X-ray source.**Distance:** H_I absorption suggests 11.3 kpc.**References:**

- Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 2.7 GHz (4′).
Downes & Wilson 1974, *A&A*, 34, 133. Effelsberg 100-m at 10.7 GHz (1′3).
Green *et al.* 1975, *A&A*, 44, 187. Effelsberg 100-m at 15.0 GHz (58″: $S=9.0\pm 0.7$ Jy).
Lockhart & Goss 1978, *A&A*, 67, 355. H_I absorption.
Altenhoff *et al.* 1979, *A&AS*, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6).
Pye *et al.* 1984, *MNRAS*, 207, 649. Einstein observations, and VLA at 1.4 and 5 GHz (both 12″).
Smith *et al.* 1985, *ApJ*, 296, 469. EXOSAT spectrum.
Morsi & Reich 1987, *A&AS*, 71, 189. Effelsberg 100-m at 32 GHz (smoothed to 30″: $S=6.90\pm 0.38$ Jy).
- Moffett & Reynolds 1994, *ApJ*, 437, 705. VLA at 330 MHz (6″7×7″7: $S=64.4$ Jy), 1.48 GHz (4″8×5″2: $S=31.8$ Jy) and 4.85 GHz (4″0×4″1), including polarisation.
Fujimoto *et al.* 1995, *PASJ*, 47, L31. ASCA observations.
Gorham *et al.* 1996, *ApJ*, 458, 257. Pulsar search.
Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
Hwang *et al.* 2000, *ApJ*, 532, 970. ROSAT image and ASCA spectroscopy.
Sugizaki *et al.* 2001, *ApJS*, 134, 77. ASCA survey observations.
Aharonian *et al.* 2001, *A&A*, 375, 1008. HESS limit.
Brogan & Troland 2001, *ApJ*, 550, 799. VLA at 1.4 GHz (24″×27″ and 5″) for H_I Zeeman splitting.
Lacey *et al.* 2001, *ApJ*, 559, 954. VLA at 74 MHz (23″×26″: $S=55.6$ Jy) and 326 MHz (6″2×6″6: $S=56.0$ Jy).
Kaplan *et al.* 2002, *ApJ*, 566, 378. VLA at 333 MHz (50″), and other observations of the region.
Kawasaki *et al.* 2005, *ApJ*, 631, 935. ASCA observations.
Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
Miceli *et al.* 2006, *A&A*, 453, 567. XMM-Newton observations.
Keohane *et al.* 2007, *ApJ*, 654, 938. IR and Chandra observations.
Ozawa *et al.* 2009, *ApJ*, 706, L71. Suzaku observations.
Abdo *et al.* 2010, *ApJ*, 722, 1303. Fermi observations.
Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz (9′5: $S=19.1\pm 1.0$ Jy) including polarisation and review of flux densities.
Rodes-Roca 2013, *A&A*, 555, A115. IR of compact X-ray source.
Yang *et al.* 2013, *ApJ*, 766, 44. Suzaku spectroscopy.
Lopez *et al.* 2013, *ApJ*, 777, 145. Chandra spectroscopy.
Zhu *et al.* 2014, *ApJ*, 793, 95. Spitzer and other observations.
Froeblich *et al.* 2015, *MNRAS*, 454, 2586. H₂ IR observations.
Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
Kilpatrick *et al.* 2016, *ApJ*, 816, 1. CO observations, including broad lines.
Koo *et al.* 2016, *ApJ*, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
Abdalla *et al.* 2018, *A&A*, 612, A1. HESS observations.
Ranasinghe & Leahy 2018, *AJ*, 155, 204. VGPS H_I absorption observations.
Abdalla *et al.* 2018, *A&A*, 612, A5. HESS observations.
Zhou & Vink 2018, *A&A*, 615, A150. Chandra observations.
Tanaka *et al.* 2018, *ApJ*, 866, L26. NuSTAR observations.

Yamaguchi *et al.* 2018, ApJ, 868, L35. NuSTAR observations.

G43.9+1.6

RA: 19^h05^m50^s
Dec: +10°30′

1-GHz flux/Jy: 9.0
Spectral index: 0.5

Size/arcmin: 60?
Type: S?

Radio: Large, poorly defined faint shell.

Point sources: Soft gamma repeater nearby.

References:

Reich *et al.* 1988, IAUCo, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).
 Vasisht *et al.* 1994, ApJ, 431, L35. VLA at 327 MHz (3′.2×3′.4).
 Hurley *et al.* 1996, ApJ, 463, L13. Observations of soft gamma repeater field.
 Kaplan *et al.* 2002, ApJ, 566, 378. VLA at 333 MHz (50′′), and other observations of the region.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 4.55±0.24 Jy) including polarisation and review of flux densities.

G45.7–0.4

RA: 19^h16^m25^s
Dec: +11°09′

1-GHz flux/Jy: 4.2?
Spectral index: 0.4?

Size/arcmin: 22
Type: S

Radio: Shell, brightest to the SE, poorly defined to NW.

References:

Fürst *et al.* 1987, A&AS, 69, 403. Effelsberg 100-m at 4.75 GHz (2′.4 : S = 2.6±0.3 Jy), plus other flux densities.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G46.8–0.3

(HC30)

RA: 19^h18^m10^s
Dec: +12°09′

1-GHz flux/Jy: 17
Spectral index: 0.54

Size/arcmin: 15
Type: S

Has been called G46.6–0.2.

Radio: Shell, two bright arcs to NNW and SSE.

Distance: H_i absorption suggests 5.7 to 11.4 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′ : S = 9.8±0.9 Jy), and 37-m at 1.7 GHz (S = 14.5±5.5 Jy).
 Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′ : S = 20.3 Jy) and Parkes 64-m at 5 GHz (4′ : S = 7.1 Jy).
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo S₄₃₀ MHz = 46±21 Jy.
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′.8 : S = 5.9±0.6 Jy).
 Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′.6).
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (53′′×58′′ : S = 13.3±0.1 Jy).

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S = 7.02±0.18 Jy) including polarisation and review of flux densities.

Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.

G49.2–0.7

(W51)

RA: 19^h23^m50^s**1-GHz flux/Jy:** 160?**Size/arcmin:** 30**Dec:** +14°06'**Spectral index:** 0.3?**Type:** S?

Has erroneously been called G49.1–0.1.

Radio: In complex region, parameters uncertain.

Optical: Some diffuse emission possibly associated.

X-ray: Elongated east--west.

Distance: Association with CO gives 6 kpc, optical absorption suggests 5.7 kpc, and H_i absorption suggests 5.4 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Parkes 64-m at 5 GHz (4').

Sato 1973, PASJ, 25, 135. H_i absorption.

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5'), S = 51.5±3.2 Jy, for the non-thermal component, but probably confused.

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).

Seward 1990, ApJS, 73, 781. Einstein observations.

Copetti & Schmidt 1991, MNRAS, 250, 127. 151 MHz observations.

Kassim 1992, AJ, 103, 943. VLA at 327 MHz (3'1×3'5).

Subrahmanyan & Goss 1995, MNRAS, 275, 755. VLA at 330 MHz (1'1).

Koo *et al.* 1995, ApJ, 447, 211. ROSAT observations.

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Green *et al.* 1997, AJ, 114, 2058. OH masers.

Koo & Moon 1997, ApJ, 475, 194. Arecibo (3') and VLA (40''×42'') at 1.4 GHz for H_i.

Koo & Moon 1997, ApJ, 485, 263. NRAO 12-m CO and HCO⁺ observations.

Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.

Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

Koo *et al.* 2002, AJ, 123, 1629. ASCA observations.

Koo *et al.* 2005, ApJ, 633, 946. Chandra observations.

Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_i.

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.

Abdo *et al.* 2009, ApJ, 706, L1. Fermi observations.

Koo *et al.* 2010, AJ, 140, 262. H_i Zeeman splitting observations.

Ceccarelli *et al.* 2011, ApJ, 740, L4. Molecular line observations of region.

Aleksić *et al.* 2012, A&A, 541, A13. γ -ray observations.

Hanabata *et al.* 2013, PASJ, 65, 42. Suzaku observations.

Tian & Leahy 2013, ApJ, 769, L17. H_i observations of region.

Brogan *et al.* 2013, ApJ, 771, 91. VLA at 74 MHz (84''×92'') and 320 MHz (33''×35''), plus OH, molecular line and other observations.

Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_i in region.

Sasaki *et al.* 2014, A&A, 563, A9. XMM-Newton observations.

Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.

Dumes *et al.* 2014, ApJ, 786, L24. Molecular line observations.

Jogler & Funk 2016, ApJ, 816, 100. Fermi observations.

McEwen *et al.* 2016, ApJ, 826, 189. NH₃ and CH₃OH observations.

Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H α absorption observations.

G53.4+0.0

RA: 19^h29^m57^s
Dec: +18°10′

1-GHz flux/Jy: 1.5
Spectral index: 0.6?

Size/arcmin: 10?
Type: S

Has been called G53.41+0.03.

Radio: Asymmetric shell.

X-ray: Detected.

References:

Anderson *et al.* 2017, A&A, 605, A58. VLA at 1 to 2 GHz.

Driessen *et al.* 2018, ApJ, 860, 133. LOFAR at 140 MHz, plus other radio observations, and XMM-Newton observations.

G53.6–2.2

RA: 19^h38^m50^s
Dec: +17°14′

1-GHz flux/Jy: 8
Spectral index: 0.50

3C400.2, NRAO 611
Size/arcmin: 33×28
Type: S

Has been called G53.7–2.2.

Radio: Ring of emission, with extension to NW.

Optical: Filaments and diffuse emission.

X-ray: Centrally brightened, offset to NW.

Distance: Association with H α gives 2.8 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′: $S=5.3\pm 0.6$ Jy).

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′: $S=4.8\pm 0.3$ Jy).

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′: $S=11.7$ Jy).

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}}=20\pm 10$ Jy, $S_{318\text{ MHz}}=20\pm 3.6$ Jy.

Goss *et al.* 1975, A&A, 43, 459. WSRT at 610 MHz (1′×3′: $S=13.2\pm 1.6$ Jy).

Sabbadin & d’Odorico 1976, A&A, 49, 119. Optical spectra.

van den Bergh 1978, ApJS, 38, 119. Optical observations.

Rosado 1983, RMxAA, 8, 59. Optical spectra.

Blair & Long 1988, PASP, 100, 461. Optical imaging and spectroscopy.

see also: Blair & Long 1988, PASP, 100, 651. Erratum.

Long *et al.* 1991, ApJ, 373, 567. Einstein and optical observations.

Winkler *et al.* 1993, ApJ, 405, 608. Optical imaging.

Dubner *et al.* 1994, AJ, 108, 207. VLA at 327 MHz (59′′) and 1.49 GHz (52′′), plus X-rays.

Saken *et al.* 1995, ApJ, 443, 231. ROSAT observations.

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Giacani *et al.* 1998, A&AS, 133, 61. DRAO at 1.4 GHz for H α studies, including distance.

Yoshita *et al.* 2001, PASJ, 53, 93. ASCA observations, and spectral comparison with ROSAT.

Ambrocio-Cruz *et al.* 2006, RMxAA, 42, 241. Optical imaging and spectroscopy.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′:5′: $S=4.00\pm 0.22$ Jy) including polarisation and review of flux densities.

Broersen & Vink 2015, MNRAS, 446, 3885. Chandra observations.

Ergin *et al.* 2017, ApJ, 842, 22. Suzaku and Fermi observations.

G54.1+0.3

RA: 19^h30^m31^s
Dec: +18°52'

1-GHz flux/Jy: 0.5
Spectral index: 0.1

Size/arcmin: 12?
Type: C?

Radio: Filled-centre core, with possible faint diffuse emission.

X-ray: Centrally concentrated, with more extended diffuse emission.

Point sources: Central pulsar.

Distance: H_i absorption suggests 4.9, association with CO suggests 8.2 kpc, and optical absorption suggests 6.3 kpc.

References:

- Green 1985, MNRAS, 216, 691. Radio at 2.7 GHz (7''×20'').
 Reich *et al.* 1985, A&A, 151, L10. Effelsberg 100-m at 4.75 GHz (2'.4: S=0.37±0.04 Jy)
 Velusamy & Becker 1988, AJ, 95, 1162. VLA at 1.4 (14'' : S = 0.48±0.03 Jy), 1.6 (14'' : S = 0.42±0.03 Jy) and 5 GHz (5'' : S = 0.33±0.02 Jy), Ooty at 327 MHz (S = 0.50±0.08 Jy), plus review of flux densities.
 Seward 1989, AJ, 97, 481. Einstein observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Lu *et al.* 2001, A&A, 370, 570. ROSAT and ASCA observations.
 Lu *et al.* 2002, ApJ, 568, L49. Chandra observations.
 Camilo *et al.* 2002, ApJ, 574, L71. Pulsar detection.
 Kaplan & Moon 2006, ApJ, 644, 1056. IR upper limit for pulsar.
 Leahy *et al.* 2008, AJ, 136, 1477. VGPS at 1.4 GHz (1') including H_i.
 Koo *et al.* 2008, ApJ, 673, L147. Akari observations of surroundings.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Bocchino *et al.* 2010, A&A, 520, A71. XMM-Newton and Suzaku observations.
 Lang *et al.* 2010, ApJ, 709, 1125. VLA at 1.4 GHz (6''.6×6''.8), 4.7 GHz (3''.2×3''.3), and 8.2 GHz (3''.0×3''.2) and Spitzer observations.
 Acciari *et al.* 2010, ApJ, 719, L69. γ -ray observations.
 Lee *et al.* 2012, JKAS, 45, 117. CO observations of region.
 Krivonos *et al.* 2017, MNRAS, 470, 512. INTEGRAL observations.
 Temim *et al.* 2017, ApJ, 836, 129. Spitzer, Herschel and Akari observations.
 Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.
 Ranasinghe & Leahy 2018, AJ, 155, 204. VGPS H_i absorption observations.
 Rho *et al.* 2018, MNRAS, 479, 5101. Spitzer, Herschel and other observations.
 Driessen *et al.* 2018, ApJ, 860, 133. LOFAR observations at 144 MHz, plus other observations.

G54.4–0.3

(HC40)

RA: 19^h33^m20^s
Dec: +18°56'

1-GHz flux/Jy: 28
Spectral index: 0.5

Size/arcmin: 40
Type: S

Has been called G54.5–0.3.

Radio: Shell, in complex region.

Optical: Faint filaments.

Point sources: Pulsar outside NW rim.

Distance: H_i and CO observations suggest 6.6 kpc.

References:

- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5' : S = 34.4±5.0 Jy).

Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2'6).
 Caswell 1985, AJ, 90, 1224. DRAO at 1.4 GHz (1'3×2'6: $S=18\pm 4$ Jy).
 Velusamy *et al.* 1986, JApA, 7, 105. WSRT at 609 MHz (50''×191'' smoothed to 100''×200'').
 Junkes *et al.* 1992, A&AS, 96, 1. Surrounding CO.
 Junkes *et al.* 1992, A&A, 261, 289. Nearby IRAS sources.
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_I.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Park *et al.* 2013, ApJ, 777, 14. Arecibo of H_I in region.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Karpova *et al.* 2017, MNRAS, 466, 1757. X-ray observations of pulsar.
 Ranasinghe & Leahy 2017, ApJ, 843, 119. H_I and CO observations.

G55.0+0.3

RA: 19^h32^m00^s
Dec: +19°50'

1-GHz flux/Jy: 0.5?
Spectral index: 0.5?

Size/arcmin: 20×15?
Type: S

Has been called G55.2+0.5.

Radio: Faint, partial shell.

Point sources: Old pulsar nearby.

Distance: Association with H_I features implies 14 kpc.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1'0×2'5), and northern sky survey at 4.9 GHz.
 Matthews *et al.* 1998, ApJ, 493, 312. WSRT at 327 MHz (1'0×2'9: $S=0.98\pm 0.15$ Jy), DRAO at
 1.4 GHz (1'0×2'9: $S=0.25\pm 0.12$ Jy), plus H_I observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G55.7+3.4

RA: 19^h21^m20^s
Dec: +21°44'

1-GHz flux/Jy: 1?
Spectral index: 0.3?

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Point sources: Old pulsar within the boundary of the remnant.

References:

Goss *et al.* 1977, A&A, 61, 93. WSRT observations at 610 MHz (57''×156'': $S=1.9\pm 0.2$ Jy) and
 1415 MHz (27''×72'': $S=1.0\pm 0.1$ Jy).
 Bhatnagar *et al.* 2011, ApJ, 739, L20. VLA at 1.3 to 1.9 GHz (30'').
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5: $S=0.52\pm 0.03$ Jy) including polarisation
 and review of flux densities.

G57.2 + 0.8

(4C21.53)

RA: 19^h34^m59^s**Dec:** +21°57'**1-GHz flux/Jy:** 1.8**Spectral index:** 0.35**Size/arcmin:** 12?**Type:** S?**Radio:** Extended non-thermal arc.**Point sources:** Central magnetar.**Distance:** H_I observations suggest 12.5 kpc.**References:**

Sieber & Seiradakis 1984, A&A, 130, 257. Effelsberg 100-m at 1.4 GHz (8'8 : 1.34±0.1), 2.7 GHz (4'3 : 0.86±0.1), plus other surveys of the area.

Caswell *et al.* 1985, AJ, 90, 488. DRAO at 1.4 GHz (1'×3').

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S=0.74±0.04 Jy) including polarisation and review of flux densities.

Surnis *et al.* 2016, ApJ, 826, 184. GMRT at 610 MHz and VLA at 1.4 GHz.

Israel *et al.* 2016, MNRAS, 457, 3448. Chandra, XMM-Newton and Swift observations of magnetar.

Kothes *et al.* 2018, ApJ, 852, 54. DRAO at 408 MHz (2'8×8'5) and 1.4 GHz (0'82×2'5), including H_I and polarisation, plus other radio observations.

G59.5 + 0.1**RA:** 19^h42^m33^s**Dec:** +23°35'**1-GHz flux/Jy:** 3?**Spectral index:** ?**Size/arcmin:** 15**Type:** S

Has been called G59.6+0.1.

Radio: Incomplete shell.**Optical:** Diffuse shell.**References:**

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1'0×2'5 : S=5.1±0.2 Jy), and northern sky survey at 4.9 GHz.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Xu & Wang 2012, A&A, 543, A24. CO observations of SE.

G63.7 + 1.1**RA:** 19^h47^m52^s**Dec:** +27°45'**1-GHz flux/Jy:** 1.8**Spectral index:** 0.24**Size/arcmin:** 8**Type:** F**Radio:** Centrally brightened, with core.**X-ray:** Diffuse emission.**References:**

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1'0×2'2), and northern sky survey at 4.9 GHz.

Wallace *et al.* 1997, AJ, 114, 2068. WSRT at 1.4 GHz (14''×26'' : S=1.63 Jy), DRAO at 1.4 GHz (smoothed to 2'), plus review of flux densities and other observations.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S=1.12±0.06 Jy) including polarisation and review of flux densities.

Matheson *et al.* 2016, ApJ, 825, 134. XMM-Newton and Chandra observations.

G64.5 + 0.9

RA: 19^h50^m25^s
Dec: +28°16'

1-GHz flux/Jy: 0.15?
Spectral index: 0.5

Size/arcmin: 8
Type: S?

Radio: Shell with central source.

Optical: Filaments in N and W.

References:

Hurley-Walker *et al.* 2009, MNRAS, 398, 249. Radio identification.

Neustadt *et al.* 2017, MNRAS, 469, 516. Optical observations.

G65.1 + 0.6

RA: 19^h54^m40^s
Dec: +28°35'

1-GHz flux/Jy: 5.5
Spectral index: 0.61

Size/arcmin: 90×50
Type: S

Radio: Large, faint shell.

Point sources: Old pulsar nearby.

Distance: Possible association with H_I suggests 9 kpc.

References:

Landecker *et al.* 1990, A&A, 232, 207. DRAO at 408 MHz (3'5×7'0":S=9.5±0.1 Jy), and 1.4 GHz (1'0×2'0":S=5.4±1.0 Jy).

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Tian & Leahy 2006, A&A, 455, 1053. CGPS at 408 MHz (2'8×5'9":S=8.6±0.8 Jy) and 1.4 GHz (0'8×1'7":S=4.9±0.5 Jy) including H_I.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3':S=9.1±1.0 Jy) and 1420 MHz (~1':S=3.9±0.5 Jy), including review of flux densities.

Aleksić *et al.* 2010, ApJ, 725, 1629. γ -ray observations.

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9'5":S=3.2±0.3 Jy), including polarisation and review of flux densities.

Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.

G65.3 + 5.7

RA: 19^h33^m00^s
Dec: +31°10'

1-GHz flux/Jy: 42
Spectral index: 0.6

Size/arcmin: 310×240
Type: S?

Has been called G65.2+5.7.

Radio: Large, faint ring, near S91 and S94.

Optical: Filamentary ring.

X-ray: Diffuse, centrally brightened.

Distance: Optical proper motions and velocities indicates 0.8 kpc.

References:

Gull *et al.* 1977, ApJ, 215, L69. Optical plates.

Reich *et al.* 1979, A&A, 72, 270. Effelsberg 100-m observations at 1.42 GHz (smoothed to $11' : S = 42.4 \pm 1.6$ Jy), estimate $S_{408 \text{ MHz}} = 91 \pm 5$ Jy from previous sky survey.
 Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
 Rosado 1981, ApJ, 250, 222. Optical interferometry.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [O III] imagery.
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Mavromatakis *et al.* 2002, A&A, 388, 355. Optical observations.
 Boumis *et al.* 2004, A&A, 424, 583. Optical observations.
 Shelton *et al.* 2004, ApJ, 615, 275. ROSAT observations.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Xiao *et al.* 2009, A&A, 503, 827. Effelsberg 100-m at 2.7 GHz ($4.4' : S = 22 \pm 3$ Jy), and Urumqi 21-m at 4.8 GHz ($9.5' : S = 16.8 \pm 1.8$ Jy) including polarisation and review of flux densities.
 Gosachinskii 2010, AstL, 36, 260. H α observations.
 Kim *et al.* 2010, ApJ, 722, 388. Far UV observations.

G65.7 + 1.2

DA 495

RA: $19^{\text{h}}52^{\text{m}}10^{\text{s}}$
Dec: $+29^{\circ}26'$

1-GHz flux/Jy: 5.1
Spectral index: varies

Size/arcmin: 22
Type: F

Has mistakenly been called G55.7+1.2.

Radio: Centrally brightened with thick shell?

X-ray: Centrally brightened.

Point sources: Compact X-ray source near centre.

Distance: H α polarisation observations suggest 1.5 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 2.8 \pm 0.4$ Jy), and 37-m at 1.7 GHz ($S = 4.4 \pm 0.5$ Jy), plus review of flux densities.
see also: Willis 1973, A&A, 27, 483. Erratum.
 Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430 \text{ MHz}} = 8.7 \pm 4.9$ Jy, $S_{318 \text{ MHz}} = 9.7 \pm 2.2$ Jy.
 Landecker & Caswell 1983, AJ, 88, 1810. DRAO at 1.4 GHz ($0.9 \times 1.5' : S = 4.4 \pm 0.2$ Jy).
 Velusamy *et al.* 1989, JApA, 10, 161. Ooty at 327 MHz ($36'' \times 64''$), WSRT at 610 MHz ($62''$) and VLA at 1.4 GHz ($36''$ and $12''$), including IRAS imaging.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Kothes *et al.* 2004, ApJ, 607, 855. H α polarisation absorption.
 Arzoumanian *et al.* 2004, ApJ, 610, L101. ROSAT and ASCA observations of compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 6.5 \pm 0.6$ Jy) and 1420 MHz ($\sim 1' : S = 4.0 \pm 0.2$ Jy), including polarisation and review of flux densities.
 Arzoumanian *et al.* 2008, ApJ, 687, 505. Chandra observations.
 Kothes *et al.* 2008, ApJ, 687, 516. CGPS at 408 MHz ($2.9 \times 6.0' : S = 6.5 \pm 0.5$ Jy) and 1.4 GHz ($0.82 \times 1.75' : S = 4.0 \pm 0.2$ Jy), Effelsberg 100-m at 4.85 GHz ($2.45' : S = 1.6 \pm 0.1$ Jy) and 10.55 GHz (smoothed to $2.45' : S = 1.1 \pm 0.1$ Jy), plus review of flux densities.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9.5' : S = 1.95 \pm 0.10$ Jy) including polarisation and review of flux densities.
 Karpova *et al.* 2015, MNRAS, 453, 2241. Chandra and XMM-Newton observations.

G66.0–0.0

RA: 19^h57^m50^s
Dec: +29°03′

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 31×25?
Type: S

Radio: Some emission in N.

Optical: Incomplete shell.

Distance: Optical absorption suggests 2.3 kpc.

References:

Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G67.6+0.9

RA: 19^h57^m45^s
Dec: +30°53′

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 50×45?
Type: S

Radio: Arc in S.

Optical: Filamentary shell.

Distance: Optical absorption suggests 2.0 kpc.

References:

Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.

Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G67.7+1.8

RA: 19^h54^m32^s
Dec: +31°29′

1-GHz flux/Jy: 1.0
Spectral index: 0.61

Size/arcmin: 15×12
Type: S

Radio: Double arc shell.

Optical: Filaments in N.

X-ray: Detected.

Point sources: Compact X-ray source.

Distance: Optical absorption suggests 1.5--5.7 kpc.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1′0×1′9 : S = 1.9±0.1 Jy), and northern sky survey at 4.9 GHz (S = 0.42±0.05 Jy).

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Mavromatakis *et al.* 2001, A&A, 370, 265. Optical observations.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: S = 1.1±0.1 Jy) and 1420 MHz ($\sim 1'$: S = 0.68±0.04 Jy), including polarisation and review of flux densities.

Gök *et al.* 2008, Ap&SS, 318, 207. Optical observations.

Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Hui & Becker 2009, A&A, 494, 1005. Chandra observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5 : S = 0.30±0.03 Jy) including polarisation and review of flux densities.

Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G67.8+0.5**RA:** 20^h00^m00^s**Dec:** +30°51′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 7×5**Type:** ?**Radio:** Poorly resolved arc.**Optical:** Diffuse shell, brighter to W.**References:**Sabin *et al.* 2013, MNRAS, 431, 279. H α and radio survey observations.**G68.6–1.2****RA:** 20^h08^m40^s**Dec:** +30°37′**1-GHz flux/Jy:** 1.1**Spectral index:** 0.2**Size/arcmin:** 23**Type:** ?**Radio:** Faint, poorly defined source.**References:**Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz (4′.3), including polarisation.Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′.3).Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz ($\sim 1'$: $S=0.57\pm 0.08$ Jy), including review of flux densitiesSun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′.5: $S=0.80\pm 0.04$ Jy), including polarisation and review of flux densities.**G69.0+2.7****RA:** 19^h53^m20^s**Dec:** +32°55′**1-GHz flux/Jy:** 120?**Spectral index:** varies**Size/arcmin:** 80?**Type:** ?

CTB 80

An association with a SN in AD1408 has been suggested. Has been called G68.8+2.8.

Radio: Compact core, flat spectrum plateau, and steeper spectrum extensions, with spectral break?**Optical:** Expanding nebulosity near centre, with filaments to the SW and far NE.**X-ray:** Diffuse emission with compact source.**Point sources:** Pulsar at western edge of core.**Distance:** H α observations suggest 1.5 kpc, and optical absorption suggests 4.6 kpc.**References:**Angerhofer *et al.* 1981, A&A, 94, 313. WSRT at 610 MHz (56″×103″) 1.4 GHz (24″×44″) and 5 GHz (7″×13″), plus optical.Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.Sofue *et al.* 1983, PASJ, 35, 437. NRO 45-m at 10.2 GHz (2′.7).

Velusamy & Kundu 1983, JApA, 4, 253. VLA of compact sources.

Blair *et al.* 1984, ApJ, 282, 161. Optical images and spectra.

Wang & Seward 1984, ApJ, 285, 607. Einstein observations.

- Strom *et al.* 1984, A&A, 139, 43. Radio observations of flat spectrum component, VLA 5 GHz ($1''.7$) and 1.4 GHz.
- Mantovani *et al.* 1985, A&A, 145, 50. Bologna at 408 MHz ($2'.6 \times 4'.9 : S = 67.5 \pm 10.5$ Jy), Effelsberg 100-m at 1.41 ($9' : S = 62 \pm 9$ Jy), 1.72 ($7'.6 : S = 66 \pm 5$ Jy), 2.7 ($4'.5 : S = 52 \pm 4$ Jy) and 4.75 GHz ($2'.4 : S = 44 \pm 3.3$ Jy), plus review of flux densities.
- Kulkarni *et al.* 1988, Nature, 331, 50. Pulsar detection.
- Angelini *et al.* 1988, ApJ, 330, L43. EXOSAT spectra.
- Fesen *et al.* 1988, Nature, 334, 229. IRAS of surrounding shell.
- Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz ($4'.3$), including polarisation.
- Whitehead *et al.* 1989, MNRAS, 237, 1109. Optical of core.
- Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz of core, plus review of flux densities.
- Hester & Kulkarni 1989, ApJ, 340, 362. Optical imaging and spectroscopy.
- Koo *et al.* 1990, ApJ, 364, 178. Large, expanding H α shell.
- Greidanus & Strom 1990, A&A, 240, 376. Optical kinematics of core.
- Koo *et al.* 1993, ApJ, 417, 196. VLA at 1.4 GHz of surrounding H α ($1'$), plus IRAS.
- Safi-Harbi *et al.* 1995, ApJ, 439, 722. ROSAT observations.
- Srinivasan 1997, ApJ, 489, 170. γ -ray observations of pulsar.
- Mavromatakis *et al.* 2001, A&A, 371, 300. Optical observations.
- Butler *et al.* 2002, A&A, 395, 845. HST detection of pulsar.
- Migliazzo *et al.* 2002, ApJ, 567, L141. Pulsar proper motion study.
- Castelletti *et al.* 2003, AJ, 126, 2114. GMRT at 240 and 618 MHz ($17'' \times 26''$ and $6'' \times 10''$) and VLA at 324 and 1380 MHz ($63'' \times 73''$ and $78'' \times 93''$).
- Moon *et al.* 2004, ApJ, 610, L33. Chandra and HST observations of core.
- Li *et al.* 2005, ApJ, 628, 931. Chandra observations of pulsar and surroundings.
- Golden *et al.* 2005, ApJ, 635, L153. High resolution radio observations of pulsar and surroundings.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 72 \pm 7$ Jy) and 1420 MHz ($\sim 1' : S = 56 \pm 5$ Jy), including polarisation and review of flux densities.
- Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H α .
- Albert *et al.* 2007, ApJ, 669, 1143. γ -ray observations.
- Zeiger *et al.* 2008, ApJ, 674, 271. Proper motion of pulsar.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'.5 : S = 36 \pm 4$ Jy), including polarisation and review of flux densities.
- Leahy & Ranasinghe 2012, MNRAS, 423, 718. CGPS at 1.4 GHz, including H α , plus ROSAT observations.
- Park *et al.* 2013, ApJ, 777, 14. Arecibo of H α in region.
- Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30 and 44 GHz.
- Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G69.7 + 1.0

RA: 20^h02^m40^s
Dec: +32°43'

1-GHz flux/Jy: 2.0
Spectral index: 0.7

Size/arcmin: 16×14
Type: S

Radio: Poorly resolved source.

X-ray: Detected.

References:

- Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
- Junkes *et al.* 1988, LNP, 316, 134. Effelsberg 100-m at 2.7 GHz ($4'.3$), including polarisation.
- Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz ($4'.3$).
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Yoshita *et al.* 2000, PASJ, 52, 867. ROSAT and ASCA observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.2 \pm 0.4$ Jy) and 1420 MHz ($\sim 1' : S = 1.5 \pm 0.1$ Jy), including review of flux densities.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S=0.78±0.07 Jy) including polarisation and review of flux densities.

G70.0–21.5

RA: 21^h24^m00^s
Dec: +19°23'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 330×240
Type: S

Radio: Not detected.

Optical: Large, faint shell of filaments.

X-ray: Partially detected.

References:

Boumis *et al.* 2002, A&A, 396, 225. Optical and ROSAT observations.

Fesen *et al.* 2015, ApJ, 812, 37. H α and other optical/UV line and ROSAT observations.

G73.9+0.9

RA: 20^h14^m15^s
Dec: +36°12'

1-GHz flux/Jy: 9
Spectral index: 0.23

Size/arcmin: 27
Type: S?

Radio: Diffuse, centrally brightened to SW.

Optical: Faint shell.

References:

Reich *et al.* 1986, A&A, 155, 185. Effelsberg 100-m at 4.75 GHz (2'4":S=6.7±0.5 Jy), plus other flux densities.

Chastenay & Pineault 1988, IAUCom, 101, 297. DRAO at 408 MHz (3'5×5'9) and 1.4 GHz (1'0×1'7).

Pineault & Chastenay 1990, MNRAS, 246, 169. DRAO at 408 MHz (3'4×5'8":S=12.7±1.2 Jy) and 1.4 GHz (1'0×1'7":S=7.4±1.0 Jy).

Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.

Pineault *et al.* 1996, AJ, 112, 201. DRAO at 1.4 GHz (smoothed to 2') for H α .

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Mavromatakis 2003, A&A, 398, 153. Optical observations.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (\sim 3':S=10.0±1.7 Jy) and 1420 MHz (\sim 1':S=7.6±0.6 Jy), including polarisation and review of flux densities.

Sitnik 2010, ARep, 54, 317. H α and CO observations of region.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S=6.2±0.3 Jy) including polarisation and review of flux densities.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

Acerro *et al.* 2016, ApJS, 224, 8. Fermi observations.

Zdziarski *et al.* 2016, MNRAS, 455, 1451. Fermi observations.

G74.0–8.5

Cygnus Loop

RA: 20^h51^m00^s**1-GHz flux/Jy:** 210**Size/arcmin:** 230×160**Dec:** +30°40′**Spectral index:** varies**Type:** S

Has been suggested that this is two overlapping remnants.

Radio: Shell, brightest to the NE, with fainter breakout region to S, with spectral variations.

Optical: Large filamentary loop, brightest to the NE, not well defined to the S or W.

X-ray: Shell in soft X-rays.

Point sources: Several compact radio sources within the boundary of the remnant, including CL4, plus X-ray sources in S.

Distance: Stellar interactions gives 0.74 kpc.

References:

- Green 1990, AJ, 100, 1927. DRAO at 408 MHz (3′3×6′7) for spectral index study, plus X-ray and optical.
- Graham *et al.* 1991, AJ, 101, 175. Shocked molecular H outside rim in NE.
- Shull & Clarke 1991, PASP, 103, 811. Optical spectroscopy of nearside filaments.
- Fesen *et al.* 1992, AJ, 104, 719. H α imagery.
- Long *et al.* 1992, ApJ, 400, 214. Optical of Balmer dominated filament.
- Arendt *et al.* 1992, ApJ, 400, 562. IRAS observations.
- Hester *et al.* 1994, ApJ, 420, 721. H α , [OIII] and other optical observations of Balmer dominated filaments in NE.
- Gorham *et al.* 1996, ApJ, 458, 257. Pulsar search.
- Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
- Leahy *et al.* 1997, AJ, 114, 2081. DRAO at 1.4 GHz (1′×2′), including polarisation.
- Leahy & Röger 1998, ApJ, 505, 784. DRAO at 1.4 GHz (1′0×1′9) and 408 MHz (3′4×6′9), for spectral index studies in comparison with other radio observations.
- Levenson *et al.* 1998, ApJS, 118, 541. Optical images.
- Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density (S=1400±400 Jy).
- Bohigas *et al.* 1999, ApJ, 518, 324. Optical spectroscopy of surroundings.
- Levenson *et al.* 1999, ApJ, 526, 874. ROSAT images.
- Blair *et al.* 1999, AJ, 118, 942. HST observations, for distance.
- Danforth *et al.* 2000, AJ, 119, 2319. UV, optical and X-ray comparison of selected regions.
- Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
- Miyata *et al.* 2001, ApJ, 550, 1023. ASCA observations of compact X-ray sources.
- Danforth *et al.* 2001, AJ, 122, 938. Far-UV spectroscopy, H α and other optical observations of NE region.
- Levenson & Graham 2001, ApJ, 559, 948. HST of SE region.
- Uyaniker *et al.* 2002, A&A, 389, L61. Effelsberg 100-m at 2.7 GHz (4′3) including polarisation, and comparison with ROSAT data.
- Leahy 2002, AJ, 123, 2689. DRAO at 1.4 GHz (2′×4′) for H α .
- Blair *et al.* 2002, ApJS, 140, 367. UV spectroscopy.
- Levenson *et al.* 2002, ApJ, 576, 798. Chandra observations of W edge.
- Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′5: S=184±18 Jy).
- Leahy 2004, MNRAS, 351, 385. Chandra observations of SW.
- Uyaniker *et al.* 2004, A&A, 426, 909. Effelsberg 100-m at 2.7 GHz (4′3), with comparison with other data for spectral index studies.
- Blair *et al.* 2005, AJ, 129, 2268. HST of outer filaments.
- Leahy 2005, AJ, 130, 165. DRAO at 1.4 GHz of SE.
- Levenson & Graham 2005, ApJ, 622, 366. Chandra observations of knot in SE.

- Sun *et al.* 2006, A&A, 447, 937. Urumqi 25-m at 4.8 GHz (9'5": $S=90\pm 9$ Jy), with comparisons with other data for spectral index studies.
- Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
- Seon *et al.* 2006, ApJ, 644, L175. Far UV observations.
- Sankrit *et al.* 2007, AJ, 133, 1383. UV observations of part.
- Tsunemi *et al.* 2007, ApJ, 671, 1717. XMM-Newton observations of NE to SW.
- Nemes *et al.* 2008, ApJ, 675, 1293. XMM-Newton observations of NE.
- Katsuda *et al.* 2008, ApJ, 680, 1198. Chandra observations of NE.
- Uchida *et al.* 2008, ApJ, 688, 1102. XMM-Newton observations.
- Uchida *et al.* 2009, PASJ, 61, 503. Suzaku observations of N.
- Kimura *et al.* 2009, PASJ, 61, S137. Suzaku observations from NE to SW.
- Tsunemi *et al.* 2009, PASJ, 61, S147. Suzaku observations of SE.
- Kosugi *et al.* 2010, PASJ, 62, 1035. Suzaku observations of SE.
- Sankrit *et al.* 2010, ApJ, 712, 1092. Spitzer observations.
- McEntaffer & Brantseg 2011, ApJ, 730, 99. Chandra observations of E.
- Katagiri *et al.* 2011, ApJ, 741, 44. Fermi observations.
- Katsuda *et al.* 2012, ApJ, 754, L7. Suzaku and XMM-Newton observations.
- Leahy & Hassan 2013, ApJ, 764, 55. Suzaku and XMM-Newton observations of SW.
- Oakley *et al.* 2013, ApJ, 766, 51. X-ray spectroscopy.
- Kim *et al.* 2014, ApJ, 784, 12. Far UV observations.
- Sankrit *et al.* 2014, ApJ, 787, 3. Spitzer spectroscopy in SE.
- Medina *et al.* 2014, ApJ, 791, 30. Optical spectroscopy in NE.
- Raymond *et al.* 2015, ApJ, 805, 152. HST observations in NE.
see also: Raymond *et al.* 2015, ApJ, 814, 165. Erratum.
- Roberts & Wang 2015, MNRAS, 449, 1340. Suzaku observations.
- Arnaud *et al.* 2016, A&A, 586, A134. Planck flux density 30 GHz.
- Katsuda *et al.* 2016, ApJ, 819, L32. H α spectroscopy of NE.
- Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
- Anero *et al.* 2016, ApJS, 224, 8. Fermi observations.
- Boubert *et al.* 2017, A&A, 606, A14. Gaia search for runaway progenitor companion.
- Fesen *et al.* 2018, MNRAS, 481, 1786. Optical observations for distance.

G74.9+1.2

CTB 87

RA: 20^h16^m02^s
Dec: +37°12'

1-GHz flux/Jy: 9
Spectral index: varies

Size/arcmin: 8×6
Type: F

Radio: Filled-centre, with high polarisation and high frequency turnover.

X-ray: Centrally brightened.

Point sources: Compact X-ray source in SE.

Distance: Optical extinction gives 6.1 kpc.

References:

- Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo $S_{430\text{ MHz}}=12.2\pm 9.7$ Jy, $S_{318\text{ MHz}}=17.7\pm 5.0$ Jy.
- Weiler & Shaver 1978, A&A, 70, 389. WSRT at 610 MHz (57''×94'' : $S=9.1\pm 1.2$ Jy), 1.4 (24''×40'' : $S=8.7\pm 1.2$ Jy) and 5 GHz (24''×40'' : $S=5.6\pm 1.3$ Jy).
- Geldzahler *et al.* 1980, A&A, 84, 237. Effelsberg 100-m at 2.7 GHz (4'4" : $S=7.6\pm 0.5$ Jy).
- Wilson 1980, ApJ, 241, L19. Einstein observations.
- van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H α absorption of nearby compact source.
- Seauquist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.
- Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (smoothed to 40'' : $S=1.47\pm 0.19$ Jy).

Green & Gull 1989, MNRAS, 237, 555. VLA at 1.4 GHz (1'2×1'4) including H α .

Salter *et al.* 1989, *Apj*, 338, 171. NRAO 12-m at 84.2 GHz, plus review of flux densities.
 Pineault & Chastenay 1990, *MNRAS*, 246, 169. DRAO at 408 MHz ($3'.4 \times 5'.8 : S = 11.6 \pm 0.4$ Jy) and 1.4 GHz ($1'.0 \times 1'.7 : S = 7.2 \pm 0.3$ Jy).
 Asaoka & Koyama 1990, *PASJ*, 42, 625. Ginga X-ray spectrum.
 Wendker *et al.* 1991, *A&A*, 241, 551. DRAO at 408 MHz ($3'.5 \times 5'.2 : S = 13.3 \pm 0.8$ Jy) and Effelsberg 100-m at 4.8 GHz ($S = 7.5 \pm 0.7$ Jy).
 Wallace *et al.* 1994, *A&A*, 286, 565. H α of surroundings.
 Cho *et al.* 1994, *Aj*, 108, 634. CO of adjacent molecular clouds.
 Gorham *et al.* 1996, *Apj*, 458, 257. Pulsar search.
 Biggs & Lyne 1996, *MNRAS*, 282, 691. Pulsar search.
 Wallace *et al.* 1997, *A&A*, 317, 212. DRAO at 408 MHz ($3'.4 \times 5'.5$) and 1.4 GHz ($1'.0 \times 1'.6$) including H α (smoothed to $2'$).
 Lorimer *et al.* 1998, *A&A*, 331, 1002. Pulsar search.
 Kothes *et al.* 2003, *Apj*, 588, 852. CGPS at 1.4 GHz ($1' \times 1'.6$) including H α , plus CO observations.
 Kothes *et al.* 2006, *A&A*, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 11.9 \pm 0.9$ Jy) and 1420 MHz ($\sim 1' : S = 7.1 \pm 1.1$ Jy), including polarisation and review of flux densities.
 Hurley-Walker *et al.* 2009, *MNRAS*, 396, 365. Radio observations at 14 to 18 GHz.
 Sitnik 2010, *ARep*, 54, 317. H α and CO observations of region.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 6.4 \pm 0.4$ Jy) including polarisation and review of flux densities.
 Matheson *et al.* 2013, *Apj*, 774, 33. Chandra observations.
 Bassani *et al.* 2014, *A&A*, 561, A108. X-ray and γ -ray observations.
 Aliu *et al.* 2014, *Apj*, 788, 78. γ -ray detection.
 Saha 2016, *MNRAS*, 460, 3563. Fermi observations.
 Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
 Liu *et al.* 2018, *Apj*, 859, 173. CO observations.
 Abeysekara *et al.* 2018, *Apj*, 861, 134. γ -ray observations.

G76.9 + 1.0

RA: 20^h22^m20^s
Dec: +38°43'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 9
Type: C

Radio: Bipolar shell.

Point sources: Central pulsar.

References:

Taylor *et al.* 1992, *Aj*, 103, 931. WSRT at 327 MHz ($1'.0 \times 1'.6$), and northern sky survey at 4.9 GHz.
 Landecker *et al.* 1993, *A&A*, 276, 522. VLA at 1.49 GHz ($14''$), 4.86 GHz ($13'' \times 16''$) and 8.55 GHz ($11'' \times 12''$), including polarisation and review of flux densities.
 Landecker *et al.* 1997, *A&AS*, 123, 199. Miyun at 232-MHz ($3'.8 \times 5'.4$).
 Lorimer *et al.* 1998, *A&A*, 331, 1002. Pulsar search.
 Kothes *et al.* 2006, *A&A*, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 2.3 \pm 0.2$ Jy) and 1420 MHz ($\sim 1' : S = 1.35 \pm 0.07$ Jy), including polarisation and review of flux densities.
 Hurley-Walker *et al.* 2009, *MNRAS*, 396, 365. Radio observations at 14 to 18 GHz.
 Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 0.79 \pm 0.07$ Jy) including polarisation and review of flux densities.
 Marthi *et al.* 2011, *MNRAS*, 416, 2560. GMRT at 618 MHz ($51'' \times 54''$), 1160 MHz ($2''.2 \times 3''.4$), and Chandra observations of central source.
 Arzoumanian *et al.* 2011, *Apj*, 739, 39. Pulsar detection.
 Jeong *et al.* 2012, *Ap&SS*, 342, 389. CO observations of region.

G78.2+2.1**RA:** 20^h20^m50^s**Dec:** +40°26′**1-GHz flux/Jy:** 320**Spectral index:** 0.51DR4, γ Cygni SNR**Size/arcmin:** 60**Type:** S

Has been called G78.1+1.8.

Radio: In complex region (early catalogues refer to other proposed remnants in this region).

Optical: Faint filaments, spectra indicate a SNR superposed on a H II region.

X-ray: Weak emission from the SE of the remnant.

Point sources: X-ray pulsar at edge of remnant, with nebula.

Distance: Associations with other objects suggests 1.7 to 2.6 kpc.

References:

- Higgs *et al.* 1977, *AJ*, 82, 718. DRAO at 1.4 GHz ($2' \times 3' : S = 270 \pm 40$ Jy) plus some 10 GHz (4') survey data, reveals true extent of remnant.
- d'Odorico & Sabbadin 1977, *A&AS*, 28, 439. Optical spectra.
- van den Bergh 1978, *ApJS*, 38, 119. Optical observations.
- Landecker *et al.* 1980, *A&AS*, 39, 133. DRAO H α observations ($2' \times 3' .1$).
- Higgs *et al.* 1983, *AJ*, 88, 97. CO of surroundings.
- Bohigas *et al.* 1983, *RMxAA*, 8, 155. Optical spectra, find thermal only.
- Braun & Strom 1986, *A&AS*, 63, 345. WSRT H α observations.
- Fukui & Tatematsu 1988, *IAUCo*, 101, 261. CO observations of the vicinity ($2' .7$).
- Green 1989, *MNRAS*, 238, 737. OH observations.
- Pineault & Chastenay 1990, *MNRAS*, 246, 169. DRAO at 408 MHz ($3' .4 \times 5' .8 : S = 480 \pm 60$ Jy) and 1.4 GHz ($1' .0 \times 1' .7 : S = 270 \pm 40$ Jy).
- Wendker *et al.* 1991, *A&A*, 241, 551. DRAO at 408 MHz ($3' .5 \times 5' .2 : S = 540 \pm 40$ Jy) and Effelsberg 100-m at 4.8 GHz ($S = 150 \pm 15$ Jy).
- Esposito *et al.* 1996, *ApJ*, 461, 820. Associated γ -ray emission.
- Brazier *et al.* 1996, *MNRAS*, 281, 1033. γ -ray and X-ray point source.
- Biggs & Lyne 1996, *MNRAS*, 282, 691. Pulsar search.
- Zhang *et al.* 1997, *A&A*, 324, 641. Multi-frequency radio comparison.
- Lorimer *et al.* 1998, *A&A*, 331, 1002. Pulsar search.
- Roberts *et al.* 2001, *ApJS*, 133, 451. ASCA observations.
- Uchiyama *et al.* 2002, *ApJ*, 571, 866. ASCA observations.
- Mavromatakis 2003, *A&A*, 408, 237. Optical observations.
- Bykov 2004, *A&A*, 427, L21. Hard X-ray observations.
- Becker *et al.* 2004, *ApJ*, 615, 897. Chandra and other observations of compact sources.
- Weisskopf *et al.* 2006, *ApJ*, 652, 387. Chandra and other observations of compact sources.
- Kothes *et al.* 2006, *A&A*, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 500 \pm 35$ Jy) and 1420 MHz ($\sim 1' : S = 226 \pm 19$ Jy), including review of flux densities.
- Kang & Koo 2007, *ApJS*, 173, 85. SGPS of high velocity H α .
- Casandjian & Grenier 2008, *A&A*, 489, 849. γ -ray observations.
- Ladouceur & Pineault 2008, *A&A*, 490, 197. CGPS at 408 MHz ($2' .9 \times 4' .5$) and 1.4 GHz ($0' .8 \times 1' .5$).
- Gao *et al.* 2011, *A&A*, 529, A159. Urumqi 25-m at 5 GHz ($9' .5 : S = 170 \pm 18$ Jy), including polarisation and review of flux densities.
- Leahy *et al.* 2013, *MNRAS*, 436, 968. ROSAT and Chandra observations, and CGPS for H α .
- Aliu *et al.* 2013, *ApJ*, 770, 93. γ -ray observations.
- Lin *et al.* 2013, *ApJ*, 770, L9. Pulsar detection.
- Hui *et al.* 2015, *ApJ*, 799, 76. XMM-Newton and Chandra observations of pulsar, including proper motion.
- Fraija & Araya 2016, *ApJ*, 826, 31. Fermi observations.
- Acerio *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
- Abeysekara *et al.* 2018, *ApJ*, 861, 134. γ -ray observations.

G82.2 + 5.3

W63

RA: 20^h19^m00^s
Dec: +45°30′**1-GHz flux/Jy:** 120?
Spectral index: 0.5?**Size/arcmin:** 95×65
Type: S

Has been called G82.5+5.3.

Radio: Shell in the Cygnus X complex.

Optical: In complex region, but spectra indicate SNR filaments.

X-ray: Detected.

Distance: Optical absorption suggests 3.2 kpc.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′: $S=59.0\pm 3.5$ Jy).
 Sabbadin 1976, A&A, 51, 159. Optical spectra.
 Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz (6′8). Incomplete mapping.
 Rosado & González 1981, RMxAA, 5, 93. Optical spectra.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′5: $S=82.5\pm 5.5$ Jy).
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1′) including polarisation, of part.
 Mavromatakis *et al.* 2004, A&A, 415, 1051. ROSAT, ASCA and optical observations.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3′$: $S=144\pm 12$ Jy) and 1420 MHz ($\sim 1′$:
 $S=93\pm 5$ Jy), including review of flux densities.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′5: $S=49\pm 5$ Jy), including polarisation
 and review of flux densities.
 Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G83.0–0.3**RA:** 20^h46^m55^s
Dec: +42°52′**1-GHz flux/Jy:** 1
Spectral index: 0.4**Size/arcmin:** 9×7
Type: S

Radio: Incomplete shell.

References:

Taylor *et al.* 1992, AJ, 103, 931. WSRT at 327 MHz (1′0×1′5), and northern sky survey at 4.9 GHz.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3′$: $S=1.2\pm 0.3$ Jy) and 1420 MHz ($\sim 1′$:
 $S=0.8\pm 0.1$ Jy), including polarisation and review of flux densities.

G84.2–0.8**RA:** 20^h53^m20^s
Dec: +43°27′**1-GHz flux/Jy:** 11
Spectral index: 0.5**Size/arcmin:** 20×16
Type: S

Radio: Elongated shell, with a filament aligned with the major axis.

X-ray: Detected.

Distance: H α absorption suggests 6 kpc.

References:

Matthews *et al.* 1977, A&A, 55, 1. WSRT at 610 MHz ($56'' \times 81'' : S = 12.4 \pm 1.5$ Jy) and Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 6.8 \pm 1.3$ Jy).
 Matthews & Shaver 1980, A&A, 87, 255. WSRT at 1415 MHz ($23'' \times 32''$), and Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 5.6 \pm 0.5$ Jy).
 Feldt & Green 1993, A&A, 274, 421. DRAO at 1.4 GHz ($1' \times 1'.5$), including H α , plus CO observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz ($1'$) including polarisation.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 14.5 \pm 0.5$ Jy) and 1420 MHz ($\sim 1' : S = 7.2 \pm 0.8$ Jy), including review of flux densities.
 Leahy & Green 2012, ApJ, 760, 25. CGPS, including H α , plus Chandra observations.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G85.4 + 0.7

RA: 20^h50^m40^s
Dec: +45°22'

1-GHz flux/Jy: ?
Spectral index: 0.2

Size/arcmin: 24?
Type: S

Radio: Faint, incomplete shell, within larger thermal shell.

X-ray: Centrally brightened.

Distance: H α observations suggest 3.5 kpc, and optical absorption suggests 4.4 kpc.

References:

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($2'.8 \times 4'.4 : S < 0.45$ Jy) and 1.4 GHz ($0'.8 \times 1'.1$), plus H α and X-ray data.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 2.9 \pm 0.5$ Jy) and 1420 MHz ($\sim 1' : S = 2.3 \pm 0.2$ Jy), including review of flux densities.
 Jackson *et al.* 2008, ApJ, 674, 936. XMM-Newton and H α observations.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
 Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.

G85.9 – 0.6

RA: 20^h58^m40^s
Dec: +44°53'

1-GHz flux/Jy: ?
Spectral index: 0.2

Size/arcmin: 24
Type: S

Radio: Faint, incomplete shell.

Optical: Diffuse shell.

X-ray: Centrally brightened.

Distance: H α observations suggest 4.8 kpc.

References:

Kothes *et al.* 2001, A&A, 376, 641. CGPS at 408 MHz ($2'.8 \times 4'.4 : S < 0.9$ Jy) and 1.4 GHz ($0'.8 \times 1'.1$), plus H α , X-ray and optical data.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 3.0 \pm 1.3$ Jy) and 1420 MHz ($\sim 1' : S = 2.2 \pm 0.8$ Jy), including review of flux densities.
 Jackson *et al.* 2008, ApJ, 674, 936. XMM-Newton and H α observations.
 Gök *et al.* 2009, Ap&SS, 324, 17. Optical observations.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G89.0+4.7

HB21

RA: 20^h45^m00^s**1-GHz flux/Jy:** 220**Size/arcmin:** 120×90**Dec:** +50°35′**Spectral index:** 0.38**Type:** S

Radio: Distorted shell (4C50.52, an extragalactic double, is within the boundary of the remnant).

Optical: Filaments and patches.

X-ray: Centrally brightened.

Distance: Various associations suggest 0.8 kpc, and optical absorption suggests 1.9 kpc.

References:

- Hirabayashi & Takahashi 1972, PASJ, 24, 231. 30-m dish at 4.2 GHz (11′:S=160±40 Jy).
 Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′:S=148±16 Jy), plus optical filaments.
 Hill 1974, MNRAS, 169, 59. Half-Mile Telescope at 1.4 GHz (3′×3′9).
 Haslam *et al.* 1975, A&A, 39, 453. Effelsberg 100-m at 2.7 GHz (4′4)
 Fukui & Tatematsu 1988, IAUCom, 101, 261. CO observations of the vicinity (2′7).
 Tatematsu *et al.* 1990, A&A, 237, 189. DRAO at 408 MHz (3′5×4′5) and 1.4 GHz (1′0×1′3), including H_I, plus CO observations of adjacent molecular cloud.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Leahy & Aschenbach 1996, A&A, 315, 260. ROSAT observations.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koo *et al.* 2001, ApJ, 552, 175. NRAO 12-m and other CO observations (27″ and 45″) of eastern part.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′5:S=228±5 Jy).
 Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1′) including polarisation.
 Byun *et al.* 2006, ApJ, 637, 283. CO observations of surroundings.
 Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
 Leahy 2006, ApJ, 647, 1125. CGPS at 408 MHz (2′8×3′7) and 1.4 GHz (0′8×1′1).
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′:S=259±19 Jy) and 1420 MHz (~1′:S=183±9 Jy), including polarisation and review of flux densities.
 Kang & Koo 2007, ApJS, 173, 85. SGPS of high velocity H_I.
 Mavromatakis *et al.* 2007, A&A, 461, 991. Optical observations.
 Shinn *et al.* 2009, ApJ, 693, 1883. IR observations.
 Pannuti *et al.* 2010, AJ, 140, 1787. ASCA and observations.
 Shinn *et al.* 2010, AdSpR, 45, 445. IR observations in S.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′5:S=107±11 Jy), including polarisation and review of flux densities.
 Reichardt *et al.* 2012, A&A, 546, A21. Fermi detection.
 Shinn *et al.* 2012, ApJ, 759, 34. Akari observations of H₂.
 Pivato *et al.* 2013, ApJ, 779, 179. Fermi observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30 and 44 GHz.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Boubert *et al.* 2017, A&A, 606, A14. Gaia search for runaway progenitor companion.
 Shan *et al.* 2018, ApJS, 238, 35. Optical absorption for distance.
 Suzuki *et al.* 2018, PASJ, 70, 75. Suzaku observations.

G93.3+6.9

RA: 20^h52^m25^s
Dec: +55°21′

1-GHz flux/Jy: 9
Spectral index: 0.45

DA 530, 4C(T)55.38.1

Size/arcmin: 27×20
Type: C?

Has been called G93.2+6.7.

Radio: Shell, with two bright limbs, highly polarised.

X-ray: Compact central source.

Distance: H_I observations suggest 2.2 kpc.

References:

Roger & Costain 1976, A&A, 51, 151. DRAO at 1.42 GHz (2′×2′.4: S=6.9 Jy).

Haslam *et al.* 1980, A&A, 92, 57. Effelsberg 100-m at 1.72 GHz (7′.6: S=6.47±0.52 Jy) and 2.7 GHz (4′.4: S=5.64±0.64 Jy), plus review of flux densities.

Lalitha *et al.* 1984, A&A, 131, 196. Effelsberg 100-m at 4.75 GHz (smoothed to 3′: S=4.01±0.57 Jy).

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Landecker *et al.* 1999, ApJ, 527, 866. DRAO at 408 MHz (3′.5×4′.3) and 1.4 GHz (1′.0×1′.2), including polarisation and H_I.

Foster & Routledge 2003, ApJ, 598, 1005. H_I for distance.

Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′: S=10.5±0.7 Jy) and 1420 MHz, including review of flux densities.

Jiang *et al.* 2007, ApJ, 670, 1142. Chandra observations.

Bocchino *et al.* 2008, AdSpR, 41, 407. XMM-Newton observations.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G93.7–0.2

RA: 21^h29^m20^s
Dec: +50°50′

1-GHz flux/Jy: 65
Spectral index: 0.65

CTB 104A, DA 551

Size/arcmin: 80
Type: S

Has been called G93.6–0.2 and G93.7–0.3.

Radio: Distorted, faint shell.

Distance: Association with H_I features suggests 1.5 kpc.

References:

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′: S=18.4±1.0 Jy).

Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz (7′.6: S=53.5±5.0 Jy), plus review of flux densities.

Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to 2′: S=58±6 Jy).

Mantovani *et al.* 1991, A&A, 247, 545. Effelsberg 100-m at 4.75 GHz (smoothed to 3′: S=33.5±4.0 Jy), including polarisation, plus review of flux densities.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.

Mantovani *et al.* 1991, A&A, 116, 1323. VLA search for OH emission.

Uyaniker *et al.* 2002, ApJ, 565, 1022. CGPS 1.4 GHz (49″×54″), including H_I, and 408 MHz (2′.8×3′.7).

Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1′) including polarisation.

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′: S=67±6 Jy) and 1420 MHz (~1′: S=35±4 Jy), including polarisation and review of flux densities.

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5: S=25.0±2.5 Jy), including polarisation and review of flux densities.

G94.0+1.0

3C434.1

RA: 21^h24^m50^s
Dec: +51°53′**1-GHz flux/Jy:** 13
Spectral index: 0.45**Size/arcmin:** 30×25
Type: S**Radio:** Incomplete shell, containing H_I shell.**Distance:** Association with stellar wind bubble implies 5.2 kpc.**References:**

- Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′:S=6.1±0.8 Jy), and 37-m at 1.7 GHz (S=11±3 Jy).
- Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5′:S=5.8±0.4 Jy). Also NRAO 140-ft at 5 GHz (6′).
- Mantovani *et al.* 1982, A&A, 105, 176. Effelsberg 100-m at 1.7 GHz (7′6″:S=12.0±1.3 Jy), plus review of flux densities.
- Goss *et al.* 1984, A&A, 138, 469. WSRT at 610 MHz (smoothed to 100″:S=16±1.7 Jy) and Effelsberg 100-m at 4.75 GHz (2′4″:S=7.2±0.5 Jy).
- Landecker *et al.* 1985, AJ, 90, 1082. DRAO at 1.4 GHz (smoothed to 2′:S=16±3 Jy).
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Uyaniker *et al.* 2003, ApJ, 585, 785. CGPS at 1.4 GHz (1′) including polarisation.
- Foster *et al.* 2004, A&A, 417, 79. DRAO at 1.4 GHz, including H_I.
- Foster 2005, A&A, 441, 1043. CGPS at 408 MHz (2′8″×3′6″) and 1.4 GHz (0′8″×1′0″) for spectral index studies, plus other observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′:S=20±2 Jy) and 1420 MHz (~1′:S=11.3±1.0 Jy), including review of flux densities.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5″:S=6.2±0.4 Jy) including polarisation and review of flux densities.
- Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
- Jeong *et al.* 2013, ApJ, 770, 105. CO observations of region.

G96.0+2.0**RA:** 21^h30^m30^s
Dec: +53°59′**1-GHz flux/Jy:** 0.35
Spectral index: 0.6**Size/arcmin:** 26
Type: S**Radio:** Faint, arc in S, poorly defined in N.**Distance:** Association for H_I indicates 4 kpc.**References:**

- Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz (2′8″×3′5″) and 1.4 GHz (50″×61″) including H_I.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′:S=0.42±0.06 Jy) and 1420 MHz (~1′:S=0.24±0.02 Jy), including review of flux densities.
- Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9′5″:S=0.14±0.02 Jy) including polarisation and review of flux densities.

G106.3 + 2.7**RA:** 22^h27^m30^s**Dec:** +60°50′**1-GHz flux/Jy:** 6**Spectral index:** 0.6**Size/arcmin:** 60×24**Type:** C?

Incorporates the pulsar wind nebula G106.6+2.9 (the ‘Boomerang’).

Radio: Faint extended source, which brighter ‘head’ to NE.

X-ray: Pulsar and wind nebula.

Point sources: Pulsar.

References:

- Pineault & Joncas 2000, AJ, 120, 3218. DRAO at 408 MHz (3′.5×3′.9: $S=10.5\pm 0.3$ Jy) and 1.4 GHz (1′.0×1′.2: $S=4.9\pm 0.6$ Jy), plus H_i.
- Halpern *et al.* 2001, ApJ, 547, 323. X-ray and radio observations of the ‘head’.
- Halpern *et al.* 2001, ApJ, 552, L125. Pulsar detection.
- Kothes *et al.* 2001, ApJ, 560, 236. CGPS at 1.4 GHz, including H_i, plus CO and other observations.
- Ng & Romani 2004, ApJ, 601, 479. Chandra detection of pulsar wind nebula.
- Kothes *et al.* 2004, ApJ, 607, 855. H_i polarisation absorption.
- Kothes *et al.* 2006, ApJ, 638, 225. Effelsberg 100-m at 4.85 (2′.4), 8.35 (1′.4), 10.5 (1′.2) and 32 GHz (0′.45) of pulsar wind nebula, including polarisation.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: $S=8.6\pm 1.0$ Jy) and 1420 MHz ($\sim 1'$: $S=4.8\pm 0.5$ Jy), including polarisation and review of flux densities.
- Abdo *et al.* 2007, ApJ, 664, L91. γ -ray observations.
- Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
- Acciari *et al.* 2009, ApJ, 703, L6. γ -ray observations.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5: $S=2.0\pm 0.3$ Jy), including polarisation and review of flux densities.

G108.2 – 0.6**RA:** 22^h53^m40^s**Dec:** +58°50′**1-GHz flux/Jy:** 8**Spectral index:** 0.5**Size/arcmin:** 70×54**Type:** S

Radio: Faint shell.

Distance: Possible associated H_i structures suggest 3.2 kpc.

References:

- Tian *et al.* 2007, A&A, 465, 907. DRAO at 408 MHz (2′.8×3′.3: $S=11.5\pm 1.2$ Jy) and 1.4 GHz (1′.0×1′.2: $S=6.6\pm 0.7$ Jy) including H_i.

G109.1–1.0

CTB 109

RA: 23^h01^m35^s**1-GHz flux/Jy:** 20**Size/arcmin:** 28**Dec:** +58°53′**Spectral index:** 0.45**Type:** S

Radio: Semicircular shell, with the Molecular cloud S152 is to the immediate W.

Optical: Faint optical filaments.

X-ray: Semicircular shell, with pulsar at W edge.

Point sources: Long period X-ray pulsar (magnetar).

Distance: Various observations imply 3.2 kpc.

References:

- Hughes *et al.* 1981, *Apj*, 246, L127. WSRT at 610 MHz ($1' : S = 40 \pm 5$ Jy) shows bad CLEAN artefacts.
- Blair & Kirshner 1981, *Nature*, 291, 132. Optical spectra.
- Downes 1983, *MNRAS*, 203, 695. Effelsberg 100-m at 2.7 GHz ($4'.4 : S = 13.0 \pm 1.5$ Jy).
- Sofue *et al.* 1983, *PASJ*, 35, 447. NRO 45-m at 10.2 GHz ($2'.7$).
- Hughes *et al.* 1984, *Apj*, 283, 147. WSRT at 610 MHz ($20'' : S = 26 \pm 3$ Jy) 1.4 GHz ($10''$), DRAO at 1.4 GHz ($1' \times 1'.2 : S = 16.8 \pm 2$ Jy) and Algonquin 46-m at 6.5 GHz ($4'.5 : S = 6.7 \pm 1$ Jy).
- Braun & Strom 1986, *A&AS*, 63, 345. WSRT H α observations.
- Tatematsu *et al.* 1987, *PASJ*, 39, 755. NRO 45-m at 10 GHz ($2'.7$), plus polarisation.
- Tatematsu *et al.* 1987, *A&A*, 184, 279. CO observations of the surroundings ($2'.7$).
- Hanson *et al.* 1988, *A&A*, 195, 114. EXOSAT of pulsar.
- see also:* Hanson *et al.* 1988, *A&A*, 207, 204. Erratum.
- Morini *et al.* 1988, *Apj*, 333, 777. EXOSAT observations.
- Koyama *et al.* 1989, *PASJ*, 41, 461. X-ray observations of pulsar.
- Coe *et al.* 1989, *MNRAS*, 238, 649. IRAS observations of surroundings.
- Green 1989, *MNRAS*, 238, 737. OH observations.
- Tatematsu *et al.* 1990, *Apj*, 351, 157. CO of surroundings, plus X-ray observations.
- Davies & Coe 1991, *MNRAS*, 249, 313. Optical and IR observations near pulsar.
- Fesen & Hurford 1995, *Aj*, 110, 747. Optical observations.
- Hurford & Fesen 1995, *MNRAS*, 277, 549. ROSAT imaging.
- Rho & Petre 1997, *Apj*, 484, 828. ROSAT observations.
- Parmar *et al.* 1998, *A&A*, 330, 175. X-ray observations.
- Lorimer *et al.* 1998, *A&A*, 331, 1002. Pulsar search.
- Koralesky *et al.* 1998, *Aj*, 116, 1323. VLA search for OH emission.
- Patel *et al.* 2001, *Apj*, 563, L45. Chandra observations of pulsar.
- Kothes *et al.* 2002, *Apj*, 576, 169. CGPS at 1.4 GHz ($59'' \times 68''$), including H α , plus CO observations.
- Sasaki *et al.* 2004, *Apj*, 617, 322. XMM-Newton observations.
- Sasaki *et al.* 2006, *Apj*, 642, L149. CO observations of surroundings, plus Chandra observations.
- Kothes *et al.* 2006, *A&A*, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 26 \pm 3$ Jy) and 1420 MHz ($\sim 1' : S = 17.4 \pm 1.2$ Jy), including polarisation and review of flux densities.
- Tian *et al.* 2010, *MNRAS*, 404, L1. CGPS for H α absorption.
- Sun *et al.* 2011, *A&A*, 536, A83. Urumqi 25-m at 5 GHz ($9'.5 : S = 9.8 \pm 0.5$ Jy) including polarisation and review of flux densities.
- Kothes & Foster 2012, *Apj*, 746, L4. H α and CO observations of region.
- Castro *et al.* 2012, *Apj*, 756, 88. Fermi observations.
- Sasaki *et al.* 2013, *A&A*, 552, A45. Chandra observations of NE.
- Tendulakar *et al.* 2013, *Apj*, 772, 31. Pulsar proper motion study.
- Vogel *et al.* 2014, *Apj*, 789, 75. NuSTAR observations of pulsar.
- Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
- Nakano *et al.* 2017, *PASJ*, 69, 40. Suzaku observations.
- Sánchez-Cruces *et al.* 2018, *MNRAS*, 473, 1705. Optical observations.

G111.7–2.1

Cassiopeia A, 3C461

RA: 23^h23^m26^s**Dec:** +58°48′**1-GHz flux/Jy:** 2300**Spectral index:** 0.77**Size/arcmin:** 5**Type:** S

Presumably the remnant of a late 17th century SN.

Radio: Bright shell with compact knots and extended plateau of emission; shows secular decline.

Optical: Fast knots and quasi-stationary flocculi, with many filaments at large radii, and NE ‘jet’.

X-ray: Incomplete shell, with hard spectral component.

Point sources: Central compact X-ray source.

Distance: Optical expansion gives 3.3 kpc.

References:

- Anderson & Rudnick 1995, *Apj*, 441, 307. VLA at 1.4 GHz and 4.8 GHz, for proper motion studies.
- Kassim *et al.* 1995, *Apj*, 455, L59. VLA at 74 (25′) and 332 MHz (6′), including spectral index comparisons.
- Anderson & Rudnick 1996, *Apj*, 456, 234. VLA for spectral index comparisons.
- Schwarz *et al.* 1997, *A&AS*, 123, 43. WSRT at 1.4 GHz (30′) for H_i absorption.
- O’Sullivan & Green 1999, *MNRAS*, 303, 575. Flux density changes at 13.5, 15.5 and 16.5 GHz.
- Agüeros & Green 1999, *MNRAS*, 305, 957. 151 MHz observations for bulk expansion studies.
- Gotthelf *et al.* 2001, *Apj*, 552, L39. Chandra observations, showing outer shock.
- Hwang *et al.* 2001, *Apj*, 560, L175. Chandra observations of Doppler shifted lines.
- Fesen *et al.* 2001, *AJ*, 122, 2644. HST observations.
- Reynoso & Goss 2002, *Apj*, 575, 871. VLA at 5 GHz (6′′.0×6′′.4) for H₂CO absorption studies.
- Krause *et al.* 2005, *Science*, 308, 1604. Spitzer light echoes from surroundings.
- Ennis *et al.* 2006, *Apj*, 652, 376. Spitzer observations.
- Kang & Koo 2007, *ApJS*, 173, 85. SGPS of high velocity H_i.
- Rho *et al.* 2008, *Apj*, 673, 271. Spitzer observations.
- Rest *et al.* 2008, *Apj*, 681, L81. SN light echo.
- Krause *et al.* 2008, *Science*, 320, 1195. SN light echo spectrum.
- Helmboldt & Kassim 2009, *AJ*, 138, 838. Low radio frequency temporal variations.
- Patnaude & Fesen 2009, *Apj*, 697, 535. X-ray proper motions.
- Barlow *et al.* 2010, *A&A*, 518, L138. Herschel observations.
- Sibthorpe *et al.* 2010, *Apj*, 719, 1553. Akari and sub-mm observations of region.
- DeLaney *et al.* 2010, *Apj*, 725, 2038. Spitzer and Chandra observations for 3-D structure.
- Patnaude *et al.* 2011, *Apj*, 729, L28. Chandra observations of fading.
- Rest *et al.* 2011, *Apj*, 732, 3. SN light echo observations.
- Fesen *et al.* 2011, *Apj*, 736, 109. HST variability studies.
- Besel & Krause 2012, *A&A*, 541, L3. IR light echoes.
- Rho *et al.* 2012, *Apj*, 747, L6. Akari CO observations.
- Vogt *et al.* 2012, *Apj*, 750, 155. Spitzer light echoes.
- Asgekar *et al.* 2013, *A&A*, 551, L11. LOFAR of carbon recombination lines.
- Yang *et al.* 2013, *Apj*, 766, 44. Suzaku spectroscopy.
- Rutherford *et al.* 2013, *Apj*, 769, 64. Chandra spectroscopy.
- Milisavljevic & Fesen 2013, *Apj*, 772, 134. Optical spectroscopy for 3-D structure.
- Koo *et al.* 2013, *Science*, 342, 1346. IR spectroscopy.
- DeLaney *et al.* 2014, *Apj*, 785, 7. VLA plus Pie Town at 74 MHz (9′′) and comparison with higher frequencies.
- Arendt *et al.* 2014, *Apj*, 786, 55. Spitzer and Herschel IR observations.
- Patnaude & Fesen 2014, *Apj*, 789, 138. Multi-epoch optical and X-ray observations.
- Lee *et al.* 2014, *Apj*, 789, 7. Spitzer and CO observations.
- Vinyaikin 2014, *ARep*, 58, 626. Time evolution of radio emission.

Alarie *et al.* 2014, MNRAS, 441, 2996. Optical imaging/spectroscopy, including distance from expansion.
 Grefenstette *et al.* 2014, Nature, 506, 339. NuSTAR ^{44}Ti observations.
 Grefenstette *et al.* 2015, ApJ, 802, 15. NuSTAR observations.
 Lee *et al.* 2015, ApJ, 808, 98. WISE, Spitzer and other IR observations.
 Milisavljevic & Fesen 2015, Science, 347, 526. Near-IR observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 7 frequencies between 30 and 353 GHz.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Fesen & Milisavljevic 2016, ApJ, 818, 17. HST [SIII] and [SII] observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Wang & Li 2016, ApJ, 825, 102. INTEGRAL observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Oonk *et al.* 2017, MNRAS, 465, 1066. LOFAR observations, including H and C lines, 33 to 78 MHz, including 69 MHz ($9''8 \times 11''2$).
 De Looze *et al.* 2017, MNRAS, 465, 3309. Spitzer and Herschel observations.
 Trotter *et al.* 2017, MNRAS, 469, 1299. Time evolution of radio emission.
 Grefenstette *et al.* 2017, ApJ, 834, 19. NuSTAR observations.
 Sato *et al.* 2017, ApJ, 836, 225. Multi-epoch Chandra observations.
 Lee *et al.* 2017, ApJ, 837, 118. IR observations.
 Arias *et al.* 2018, A&A, 612, A110. LOFAR at 30 to 77 MHz ($7'' \times 17''$).
 Salas *et al.* 2018, MNRAS, 475, 2496. LOFAR C recombination lines as 43, 54, 148 and 340 MHz ($70''$).
 Sato *et al.* 2018, ApJ, 853, 46. Chandra and NuSTAR observations.
 Zhou *et al.* 2018, ApJ, 865, 6. CO observations.
 Raymond *et al.* 2018, ApJ, 866, 128. IR observations.
 Koo *et al.* 2018, ApJ, 866, 139. IR observations.

G113.0+0.2

RA: $23^{\text{h}}26^{\text{m}}50^{\text{s}}$
Dec: $+61^{\circ}26'$

1-GHz flux/Jy: 4
Spectral index: 0.5?

Size/arcmin: $40 \times 17?$
Type: ?

Radio: Elongated, extent not well defined.

Point sources: Contains old pulsar.

Distance: Association for H_I indicates 3.1 kpc.

References:

Kothes *et al.* 2005, A&A, 444, 871. CGPS at 408 MHz ($2'8 \times 3'1$) and 1.4 GHz ($49'' \times 55''$) including H_I.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$) and 1420 MHz ($\sim 1'$).
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5 : S = 1.9 \pm 0.5$ Jy) including polarisation and review of flux densities.

G114.3 + 0.3**RA:** 23^h37^m00^s**Dec:** +61°55′**1-GHz flux/Jy:** 5.5**Spectral index:** 0.5**Size/arcmin:** 90×55**Type:** S**Radio:** Shell, with H_{II} region S165 within the boundary of the remnant.**Optical:** Faint emission in centre and to S.**Point sources:** Pulsar near centre of remnant.**Distance:** Association with H_I and other features implies 0.7 kpc.**References:**

- Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4′.4 : S = 3.6 Jy) and S_{1.4 GHz} = 4.4 Jy from 1.4 GHz survey data, plus H_I from Maryland--Green Bank survey.
- Kulkarni *et al.* 1993, Nature, 362, 135. Pulsar association.
- Fürst *et al.* 1993, A&A, 276, 470. Pulsar association.
- Becker *et al.* 1996, A&A, 306, 464. ROSAT of pulsar.
- Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
- Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
- Mavromatakis *et al.* 2002, A&A, 383, 1011. Optical observations.
- Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz (49″×55″), including H_I (1′.0×1′.1).
- Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz (3′.4×3′.9 : S = 12.0±6.0 Jy) and 1.4 GHz (1′.0×1′.1 : S = 9.8±0.8 Jy).
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 1420 MHz (~1′ : S = 5.4±0.8 Jy), including review of flux densities.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 6.9±0.7 Jy), including polarisation and review of flux densities.

G116.5 + 1.1**RA:** 23^h53^m40^s**Dec:** +63°15′**1-GHz flux/Jy:** 10**Spectral index:** 0.5**Size/arcmin:** 80×60**Type:** S**Radio:** Distinct shell, with high polarisation.**Optical:** Detected.**Distance:** Association with H_I features implies 1.6 kpc.**References:**

- Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz (4′.4 : S = 4.7±0.4 Jy) and S_{1.4 GHz} = 8.0±0.8 Jy from 1.4 GHz survey data, plus H_I from Maryland--Green Bank survey.
- Fesen *et al.* 1997, AJ, 113, 767. Optical observations.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz (49″×55″), including H_I (1′.0×1′.1).
- Mavromatakis *et al.* 2005, A&A, 435, 141. Optical observations.
- Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz (3′.4×3′.8 : S = 15.0±1.5 Jy) and 1.4 GHz (1′.0×1′.1 : S = 10.6±0. Jy).
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′ : S = 12.5±1.6 Jy) and 1420 MHz (~1′ : S = 10.3±0.70 Jy), including polarisation and review of flux densities.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 5.7±0.6 Jy), including polarisation and review of flux densities.

G116.9+0.2

CTB 1

RA: 23^h59^m10^s**Dec:** +62°26′**1-GHz flux/Jy:** 8**Spectral index:** 0.57**Size/arcmin:** 34**Type:** S

Has been called G117.3+0.1 and G116.9+0.1.

Radio: Incomplete shell.

Optical: Filaments on sky survey.

X-ray: Centrally brightened, with NE 'breakout'.

Point sources: Pulsar outside rim to E.

Distance: Association with H_I features implies 1.6 kpc.

References:

Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz ($5' : S = 3.9 \pm 1.0$ Jy), and 37-m at 1.7 GHz ($S = 5.5 \pm 2.0$ Jy), plus review of flux densities.

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz ($5' : S = 4.2 \pm 0.2$ Jy).

Angerhofer *et al.* 1977, A&A, 55, 11. NRAO 140-ft at 5 GHz ($6'8 : S = 3.0 \pm 0.3$ Jy).

Dickel & Willis 1980, A&A, 85, 55. WSRT at 610 MHz ($56'' \times 64''$) and 1.4 GHz ($22'' \times 25''$).

Reich & Braunsfurth 1981, A&A, 99, 17. Effelsberg 100-m at 2.7 GHz ($4'4 : S = 4.8 \pm 0.4$ Jy) and $S_{1.4 \text{ GHz}} = 7.8 \pm 0.8$ Jy from 1.4 GHz survey data, plus H_I from Maryland--Green Bank survey.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Landecker *et al.* 1982, AJ, 87, 1379. DRAO at 1.42 GHz and H_I ($2' \times 2'3 : S = 8.3 \pm 0.5$ Jy), plus review of flux densities.

see also: Landecker *et al.* 1983, AJ, 88, 877. Erratum.

Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.

Hailey & Craig 1994, ApJ, 434, 635. Optical spectroscopy.

Hailey & Craig 1995, ApJ, 455, L151. ROSAT of nearby pulsar.

Fesen *et al.* 1997, AJ, 113, 767. Optical observations.

Craig *et al.* 1997, ApJ, 488, 307. ROSAT observations.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 10.6 GHz ($69''$), including polarisation.

Yar-Uyaniker *et al.* 2004, ApJ, 616, 247. CGPS at 1.4 GHz ($49'' \times 55''$), including H_I ($1'0 \times 1'1$).

Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.

Tian & Leahy 2006, ChJAA, 6, 543. CGPS at 408 MHz ($3'4 \times 3'8 : S = 15.0 \pm 1.5$ Jy) and 1.4 GHz ($1'0 \times 1'1 : S = 8.1 \pm 0.4$ Jy).

Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 10.5 \pm 0.8$ Jy) and 1420 MHz ($\sim 1' : S = 7.0 \pm 0.8$ Jy), including review of flux densities.

Pannuti *et al.* 2010, AJ, 140, 1787. ASCA and Chandra observations.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz ($9'5 : S = 3.6 \pm 0.4$ Jy) including polarisation and review of flux densities.

Clark *et al.* 2017, ApJ, 834, 106. Fermi pulsar detection.

Wu *et al.* 2018, ApJ, 854, 99. Radio detection of pulsar.

Zyuzin *et al.* 2018, MNRAS, 476, 2177. X-ray observations of pulsar.

Katsuragawa *et al.* 2018, PASJ, 70, 110. Suzaku observations.

G119.5 + 10.2

CTA 1

RA: 00^h06^m40^s**1-GHz flux/Jy:** 36**Size/arcmin:** 90?**Dec:** +72°45'**Spectral index:** 0.6**Type:** S

Has been called G119.5+10.3.

Radio: Incomplete shell, with 'breakout' to NW.

Optical: Faint diffuse nebulosities.

X-ray: Centrally brightened.

Point sources: Central pulsar.

Distance: Associated H_I shell indicates 1.4 kpc.

References:

- Sieber *et al.* 1979, A&A, 74, 361. Effelsberg 100-m at 2.7 GHz (4'4": $S=23.6\pm 2.8$ Jy).
 Sieber *et al.* 1981, A&A, 103, 393. Effelsberg 100-m at 2.7 GHz and 1.7 GHz (7'6": $S=31.6\pm 2.5$ Jy), and 151 MHz (4' : $S=62.6\pm 6$ Jy).
 Fesen *et al.* 1981, ApJ, 247, 148. Optical, including spectra.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Simonetti 1992, ApJ, 386, 170. VLA observations of background sources for rotation measure studies.
 Pineault *et al.* 1993, AJ, 105, 1060. DRAO at 1.4 GHz (1' : $S=34\pm 4$ Jy), plus H_I and IRAS.
 Seward *et al.* 1995, ApJ, 453, 284. ROSAT observations.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Slane *et al.* 1997, ApJ, 485, 221. ASCA and ROSAT observations.
 Pineault *et al.* 1997, A&A, 324, 1152. DRAO at 408 MHz (3'5") and 1.4 GHz (1'0).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Brazier *et al.* 1998, MNRAS, 295, 819. Studies of central, compact γ -ray source.
 Mavromatakis *et al.* 2000, A&A, 353, 371. Optical imagery.
 Roberts *et al.* 2001, ApJS, 133, 451. ASCA observations.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 1.4 GHz (9'4).
 Slane *et al.* 2004, ApJ, 601, 1045. XMM-Newton observations of central source.
 Halpern *et al.* 2004, ApJ, 612, 398. Chandra observations of central nebula, plus optical and radio limits for compact source.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Abdo *et al.* 2008, Science, 322, 1218. Fermi detection of pulsar.
 Lin *et al.* 2010, ApJ, 725, L1. XMM-Newton observations of pulsar.
 Caraveo *et al.* 2010, ApJ, 725, L6. XMM-Newton observations of pulsar.
 Sun *et al.* 2011, A&A, 535, A64. Urumqi 25-m at 4.8 GHz (9'5": $S=11.6\pm 1.2$ Jy) and Effelsberg 100-m at 2.6 GHz (4'4": $S=20.3\pm 2.0$ Jy) including polarisation.
 Lin *et al.* 2012, MNRAS, 426, 2283. Suzaku observations.
 Mignani *et al.* 2013, MNRAS, 430, 1354. Optical limits for pulsar.
 Aliu *et al.* 2013, ApJ, 764, 38. γ -ray observations.
 Li *et al.* 2016, ApJ, 831, 19. Fermi observations of pulsar.
 Ackermann *et al.* 2018, ApJS, 237, 32. Fermi observations.

G120.1 + 1.4

Tycho, 3C10, SN1572

RA: 00^h25^m18^s**1-GHz flux/Jy:** 50**Size/arcmin:** 8**Dec:** +64°09′**Spectral index:** 0.58**Type:** S

This is the remnant of the Tycho's SN of AD1572.

Radio: Shell, brightest to the NE.

Optical: Faint filaments/knots to the NNW, NE and E.

X-ray: Shell, brighter to the NE.

Point sources: Faint radio source near centre of the remnant, thought to be extragalactic.

Distance: H α observations suggest 2.3--3 kpc, optical proper motion and shock velocity gives 2.4 kpc.

References:

- Duin & Strom 1975, A&A, 39, 33. WSRT at 610 MHz (57'' \times 64'') and 5 GHz (7'' \times 8'').
- Klein *et al.* 1979, A&A, 76, 120. Effelsberg 100-m at 10.7 GHz (1'2 : S = 13.1 \pm 0.8 Jy), plus review of flux densities.
- Strom *et al.* 1982, MNRAS, 200, 473. WSRT at 1415 MHz (27'' \times 31'') from 1971 and 1979, for expansion.
- Dickel *et al.* 1982, ApJ, 257, 145. Comparison of radio, X-ray and optical observations.
- Reid *et al.* 1982, ApJ, 261, 485. Einstein observations.
- Seward *et al.* 1983, ApJ, 266, 287. Einstein observations.
- Tan & Gull 1985, MNRAS, 216, 949. Radio at 2.7 GHz (4'') from 1980 and 1983, and One-Mile Telescope at 1.4 GHz (23'') from 1965 and 1980, for expansion.
- Albinson *et al.* 1986, MNRAS, 219, 427. H α observations.
- Green & Gull 1987, MNRAS, 224, 1055. VLA H α absorption observations towards central radio source.
- Wood *et al.* 1992, AJ, 103, 1338. VLA at 5 GHz (1''5) polarisation studies.
- Vancura *et al.* 1995, ApJ, 441, 680. X-ray spectra and ROSAT image.
- Predehl & Schmitt 1995, A&A, 293, 889. ROSAT of dust scattered halo.
- Schwarz *et al.* 1995, A&A, 299, 193. WSRT and Effelsberg 100-m at 1.4 GHz (50''), and VLA at 1.4 GHz (13''), for neutral hydrogen studies.
- Hwang & Gotthelf 1997, ApJ, 475, 665. ASCA observations.
- Reynoso *et al.* 1997, ApJ, 491, 816. VLA at 1.4 GHz (1''4 \times 1''5) from 1984 and 1994 for expansion studies.
- Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
- Reynoso *et al.* 1999, AJ, 117, 1827. VLA at 1.4 GHz (13''), for H α studies.
- Katz-Stone *et al.* 2000, ApJ, 529, 453. VLA at 333 MHz and 1.4 GHz (7''7 \times 9''5), for spectral index studies.
- Ghavamian *et al.* 2000, ApJ, 535, 266. Optical observations of shock precursors.
see also: Ghavamian *et al.* 2017, ApJ, 843, 77. Erratum.
- Hughes 2000, ApJ, 545, L53. ROSAT X-ray expansion.
- Decourchelle *et al.* 2001, A&A, 365, L218. XMM-Newton observations.
- Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
- Douvion *et al.* 2001, A&A, 373, 281. ISO observations.
- Lee *et al.* 2004, ApJ, 605, L113. Observations of molecular clouds in vicinity.
- Warren *et al.* 2005, ApJ, 634, 376. Chandra observations.
- Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (\sim 3' : S = 86 \pm 5 Jy) and 1420 MHz (\sim 1' : S = 40.5 \pm 1.5 Jy), including polarisation and review of flux densities.
- Cassam-Chenai *et al.* 2007, ApJ, 665, 315. Chandra observations.
- Rest *et al.* 2008, ApJ, 681, L81. SN light echo.
- Krause *et al.* 2008, Nature, 456, 617. SN light echo spectrum.
- Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Raymond *et al.* 2010, ApJ, 712, 901. H α spectroscopy.
 Lee *et al.* 2010, ApJ, 715, L146. H α observations.
 Hayato *et al.* 2010, ApJ, 725, 894. Suzaku observations of expansion.
 Ishihara *et al.* 2010, A&A, 521, L61. Akari observations.
 Tian & Leahy 2011, ApJ, 729, L15. H α and CO observations.
 Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 20.0 \pm 2.0 Jy) including polarisation and review of flux densities.
 Giordano *et al.* 2012, ApJ, 744, L2. Fermi observations.
 Gomez *et al.* 2012, MNRAS, 420, 3557. Herschel IR dust observations.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.
 Williams *et al.* 2013, ApJ, 770, 129. Spitzer observations.
 Yamaguchi *et al.* 2014, ApJ, 780, 136. Suzaku observations.
 Wang & Li 2014, ApJ, 789, 123. INTEGRAL observations.
 Troja *et al.* 2014, ApJ, 797, L6. Swift observations.
 Miceli *et al.* 2015, ApJ, 805, 120. XMM-Newton observations.
 Lu *et al.* 2015, ApJ, 805, 142. Chandra observations.
 Katsuda *et al.* 2015, ApJ, 808, 49. Suzaku observations.
 Tran *et al.* 2015, ApJ, 812, 101. Chandra observations.
 Lopez *et al.* 2015, ApJ, 814, 132. NuSTAR observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 5 frequencies between 30 and 143 GHz.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Williams *et al.* 2016, ApJ, 823, L32. Multi-epoch X-ray and radio observations for expansion.
 Zhou *et al.* 2016, ApJ, 826, 34. CO observations.
 Chen *et al.* 2017, A&A, 604, A113. CO observations.
 Yamaguchi *et al.* 2017, ApJ, 834, 124. Suzaku observations of E rim.
 Archambault *et al.* 2017, ApJ, 836, 23. γ -ray observations.
 Sato & Hughes 2017, ApJ, 840, 112. Chandra and Suzaku observations.
 Williams *et al.* 2017, ApJ, 842, 28. Multi-epoch Chandra observations for expansion studies.
 Knežević *et al.* 2017, ApJ, 846, 167. H α observations of NE.
 Kerzendorf *et al.* 2018, MNRAS, 479, 5696. HST search for progenitor companion.
 Vinyaikin *et al.* 2018, ARep, 62, 130. Time evolution of radio emission.

G126.2 + 1.6

RA: 01^h22^m00^s

Dec: +64°15'

1-GHz flux/Jy: 6

Spectral index: 0.5

Size/arcmin: 70

Type: S?

Radio: Poorly defined shell.

Optical: Filaments, mostly in W.

References:

Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz (9' : S = 6.8 \pm 0.7 Jy) and 2.7 GHz (4'4 : S = 3.9 \pm 0.4 Jy).
 Blair *et al.* 1980, ApJ, 242, 592. Optical detection and spectra.
 Rosado 1982, RMxAA, 5, 127. Optical spectra.
 Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz (4'4) and 4.8 GHz (2'6).
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz (3'5 \times 3'9 : S = 12 \pm 2.5 Jy) and part at 1.4 GHz (1'0 \times 1'1), plus review of flux densities.
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5 : S = 6.1 \pm 1.6 Jy).
 Boumis *et al.* 2005, A&A, 443, 175. Optical observations.

Tian & Leahy 2006, A&A, 447, 205. CGPS at 408 MHz ($3'.4 \times 3'.8 : S = 9.7 \pm 3.9$ Jy) and 1.4 GHz ($1'.0 \times 1'.1 : S = 6.7 \pm 2.1$ Jy), plus other observations for spectral index studies.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 5.7 \pm 0.7$ Jy) and 1420 MHz ($\sim 1' : S = 6.4 \pm 1.1$ Jy), including review of flux densities.
 Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz ($9'.5 : S = 2.6 \pm 0.6$ Jy), including polarisation.
see also: Sun *et al.* 2007, A&A, 469, 1003. Erratum.

G127.1+0.5

R5

RA: 01^h28^m20^s**1-GHz flux/Jy:** 12**Size/arcmin:** 45**Dec:** +63°10'**Spectral index:** 0.45**Type:** S

Has been called G127.3+0.7.

Radio: Distinct shell, with bright central source.

Optical: Detected.

Point sources: Flat radio spectrum (extragalactic) source at centre of remnant.

Distance: 1.2--1.3 kpc if associated with NGC 559.

References:

Caswell 1977, MNRAS, 181, 789. Half-Mile Telescope at 1.42 GHz ($3'.5 \times 3'.9$), plus other observations of central source.
 Pauls 1977, A&A, 59, L13. Effelsberg 100-m at 1.4 GHz ($9' : S = 8 \pm 1$ Jy).
 Salter *et al.* 1978, A&A, 66, 77. Effelsberg 100-m at 2.7 GHz ($4'.4$), plus 5 and 8.7 GHz of the central source.
 Reich *et al.* 1979, A&A, 78, L13. Effelsberg 100-m at 1.4 GHz ($9' : S = 10.8 \pm 1.3$ Jy).
 Pauls *et al.* 1982, A&A, 112, 120. WSRT at 610 MHz ($56'' \times 62''$) and H_i absorption to the point source with the VLA.
 Geldzahler & Shaffer 1982, ApJ, 260, L69. Observations of central source.
 Fürst *et al.* 1984, A&A, 133, 11. Effelsberg 100-m at 2.7 GHz ($4'.4$) and 4.8 GHz ($2'.6$).
 Goss & van Gorkom 1984, JApA, 5, 425. WSRT H_i absorption of central source.
 Joncas *et al.* 1989, A&A, 219, 303. DRAO at 408 MHz ($3'.5 \times 3'.9 : S = 17.9 \pm 2.0$ Jy) and 1.4 GHz ($1'.0 \times 1'.13 : S = 10.1 \pm 0.8$ Jy), plus review of flux densities.
 Xilouris *et al.* 1993, A&A, 270, 393. Optical imaging.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'.5 : S = 14.6 \pm 0.8$ Jy).
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact sources.
 Leahy & Tian 2006, A&A, 451, 251. CGPS at 408 MHz ($3'.4 \times 3'.8 : S = 17.1 \pm 1.7$ Jy) and 1.4 GHz ($1'.0 \times 1'.2 : S = 10.0 \pm 0.8$ Jy).
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3' : S = 15.9 \pm 1.0$ Jy) and 1420 MHz ($\sim 1' : S = 9.7 \pm 0.6$ Jy), including polarisation and review of flux densities.
 Sun *et al.* 2007, A&A, 463, 993. Urumqi 25-m at 5 GHz ($9'.5 : S = 6.3 \pm 0.7$ Jy), including polarisation.
see also: Sun *et al.* 2007, A&A, 469, 1003. Erratum.
 Zhou *et al.* 2014, ApJ, 791, 109. CO observations.

G130.7 + 3.1

3C58, SN1181

RA: 02^h05^m41^s**1-GHz flux/Jy:** 33**Size/arcmin:** 9×5**Dec:** +64°49′**Spectral index:** 0.07**Type:** F

This is the remnant of the SN of AD1181.

Radio: Filled-centre, highly polarised, with high frequency turnover.

Optical: Faint filaments.

X-ray: Centrally brightened, with faint jet.

Point sources: Central pulsar.

Distance: H α absorption indicates 2 kpc.

References:

- Green *et al.* 1975, A&A, 44, 187. Effelsberg 100-m at 15.0 GHz (58'' : S = 26.7±0.5 Jy).
 Wilson & Weiler 1976, A&A, 49, 357. WSRT at 610 MHz (58''×64''), 1.4 GHz (24''×27'') and 5 GHz (7''×8'').
 van den Bergh 1978, ApJ, 220, L9. Optical observations.
 Becker *et al.* 1982, ApJ, 255, 557. X-ray observations.
 Green & Gull 1982, Nature, 299, 606. H α absorption distance.
 Fesen 1983, ApJ, 270, L53. Optical spectra.
 Reynolds & Aller 1985, AJ, 90, 2312. VLA at 1.4 GHz (2''), for limits of shell.
 Davelaar *et al.* 1986, ApJ, 300, L59. EXOSAT spectrum.
 Green 1986, MNRAS, 218, 533. 151 MHz observations (1'2×1'3 : S = 36±4 Jy), plus 2.7 GHz (4''), plus Einstein observations for limit on shell.
 Green 1987, MNRAS, 225, 11P. Flux density increase at 408 MHz.
 Morsi & Reich 1987, A&AS, 69, 533. Effelsberg 100-m at 32 GHz (26'5 : S = 24.2±1.4 Jy).
 Reynolds & Aller 1988, ApJ, 327, 845. VLA at 1.4 (2''4) and 4.9 GHz (2''5).
 Salter *et al.* 1989, ApJ, 338, 171. NRAO 12-m at 84.2 GHz (90'' : S = 15.0±2.0 Jy), plus review of flux densities.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Green & Scheuer 1992, MNRAS, 258, 833. IRAS upper limits.
 Roberts *et al.* 1993, A&A, 274, 427. H α absorption.
 Wallace *et al.* 1994, A&A, 286, 565. H α of surroundings.
 Helfand *et al.* 1995, ApJ, 453, 741. ROSAT observations.
 Torii *et al.* 2000, PASJ, 52, 875. ASCA observations.
 Bietenholz *et al.* 2001, ApJ, 560, 772. VLA at 74 MHz (26'' : S = 33.6 Jy) and 327 MHz (8''2 : S = 33.9 Jy), for spectral index and expansion studies.
 Bocchino *et al.* 2001, A&A, 369, 1078. XMM-Newton observations.
 Murray *et al.* 2002, ApJ, 568, 226. Chandra pulsar detection.
 Camilo *et al.* 2002, ApJ, 571, L41. Pulsar detection in radio.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 32 GHz (26'') for polarised intensity.
 Slane *et al.* 2002, ApJ, 571, L45. Chandra observations.
 Slane *et al.* 2004, ApJ, 616, 403. Deep Chandra imaging.
 Bietenholz 2006, ApJ, 645, 1180. VLA at 1.4 GHz (1'36) for expansion studies.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3' : S = 32.2±2.0 Jy) and 1420 MHz (~1' : S = 31.9±1.0 Jy), including polarisation and review of flux densities.
 Gotthelf *et al.* 2007, ApJ, 654, 267. XMM-Newton observations.
 Slane *et al.* 2008, ApJ, 676, L33. Spitzer and other IR observations.
 Fesen *et al.* 2008, ApJS, 174, 379. Optical observations for proper motion studies.
 Shearer & Neustroev 2008, MNRAS, 390, 235. Optical observations of pulsar nebula.
 Shibano *et al.* 2008, A&A, 486, 273. Optical observations of pulsar nebula.
 Abdo *et al.* 2009, ApJ, 699, L102. Fermi observations of pulsar.
 Livingstone *et al.* 2009, ApJ, 706, 1163. Pulsar observations.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5":S=31.7±3.0 Jy) including polarisation and review of flux densities.
 Bietenholz *et al.* 2013, MNRAS, 431, 2590. Proper motion study of pulsar.
 Kothes *et al.* 2013, A&A, 560, A18. CGPS H α observations for distance.
 Aleksić *et al.* 2014, A&A, 567, L8. γ -ray detection.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 7 frequencies between 30 and 353 GHz.

G132.7 + 1.3

HB3

RA: 02^h17^m40^s**1-GHz flux/Jy:** 45**Size/arcmin:** 80**Dec:** +62°45'**Spectral index:** 0.6**Type:** S

Has been called G132.4+2.2.

Radio: Faint shell, adjacent to W3/4/5 complex.

Optical: Complete, filamentary shell, shock excited spectra.

X-ray: Partial shell.

Point sources: Pulsar nearby.

Distance: Interaction with surroundings gives 2 kpc.

References:

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5':S=33.8±7.0 Jy).
 d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.
 Read 1981, MNRAS, 194, 863. Radio at 151 MHz (4'4) and 1.4 GHz (2') showing H α shell.
 Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.
 Fesen & Gull 1983, PASP, 95, 196. Optical image.
 Leahy *et al.* 1985, ApJ, 294, 183. Einstein observations.
 Landecker *et al.* 1987, AJ, 94, 111. DRAO at 408 MHz (3'5×4':S=75±15 Jy), plus review of flux densities.
 Routledge *et al.* 1991, A&A, 247, 529. DRAO at 1.4 GHz (1'0×1'1) for H α , plus CO observations.
 Fesen *et al.* 1995, AJ, 110, 2876. Optical imaging and spectroscopy, DRAO at 408 MHz (3'5×4' and 1.4 GHz (1'0×1'1)).
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14'5":S=51.5±3.5 Jy).
 Tian & Leahy 2005, A&A, 436, 187. CGPS at 408 MHz (3'4×3'8") and 1.4 GHz (1'0×1'1), for spectral index studies.
see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.
 Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3':S=61±9 Jy) and 1420 MHz (~1':S=29.4±2.7 Jy), including review of flux densities.
 Lazendic & Slane 2006, ApJ, 647, 350. X-ray observations.
 Green 2007, BASI, 35, 77. Review of radio spectrum.
 Shi *et al.* 2008, A&A, 487, 601. Urumqi 25-m at 4.8 GHz (9'5), plus other survey observations for spectral studies.
 Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
 Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Katagiri *et al.* 2016, ApJ, 818, 114. Fermi observations.
 Zhou *et al.* 2016, ApJ, 833, 4. CO observations of region.

G150.3 + 4.5**RA:** 04^h27^m00^s**Dec:** +55°28′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 180×150**Type:** S**Radio:** Faint radio shell.**References:**

Gao & Han 2014, A&A, 567, A59. Radio surveys at 1.4, 2.7 and 5 GHz (9′.4, 4′.3, 9′.5).

Ackermann *et al.* 2017, ApJ, 843, 139. Fermi observations.Ackermann *et al.* 2018, ApJS, 237, 32. Fermi observations.**G152.4–2.1****RA:** 04^h07^m50^s**Dec:** +49°11′**1-GHz flux/Jy:** 3.5?**Spectral index:** 0.7?**Size/arcmin:** 100×95**Type:** S**Radio:** Bilateral shell.**References:**Foster *et al.* 2013, A&A, 549, A107. Effelsberg 100-m at 2.7 GHz, including polarisation, plus various radio survey observations.**G156.2 + 5.7****RA:** 04^h58^m40^s**Dec:** +51°50′**1-GHz flux/Jy:** 5**Spectral index:** 0.5**Size/arcmin:** 110**Type:** S**Radio:** Faint shell, brighter in E and W.**Optical:** Filamentary ring and smaller patchy ring.**X-ray:** Faint shell.**Distance:** Optical/X-ray observations imply >1.7 kpc.**References:**Pfeffermann *et al.* 1991, A&A, 246, L28. ROSAT detection.Reich *et al.* 1992, A&A, 256, 214. Effelsberg 100-m at 1.4 (9′ : S = 4.2±1.0 Jy) and 2.7 GHz (4′.3 : S = 3.0±1.0 Jy), plus H α and IRAS.Yamauchi *et al.* 1993, PASJ, 45, 795. Hard X-ray observations.Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.Yamauchi *et al.* 1999, PASJ, 51, 13. ASCA observations of some regions.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz, including polarisation.

Pannuti & Allen 2004, AdSpR, 33, 434. ASCA and RXTE observations.

Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz ($\sim 3'$: S = 8.1±1.3 Jy), including review of flux densities.Xu *et al.* 2007, A&A, 470, 969. Urumqi 25-m at 4.8 GHz (9′.5 : S = 2.5±0.5 Jy), including polarisation.

Gerardy & Fesen 2007, MNRAS, 376, 929. Optical observations.

Katsuda *et al.* 2009, PASJ, 61, S155. Suzaku observations.Uchida *et al.* 2012, PASJ, 64, 61. Suzaku observations.Katsuda *et al.* 2016, ApJ, 826, 108. Multi-epoch H α observations for expansion.

G159.6 + 7.3**RA:** 05^h20^m00^s**Dec:** +50°00′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 240×180?**Type:** S**Radio:** Not detected.**Optical:** Large, faint shell.**X-ray:** Possible emission.**References:**Fesen & Milisavljevic 2010, AJ, 140, 1163. H α and ROSAT observations**G160.9 + 2.6**

HB9

RA: 05^h01^m00^s**Dec:** +46°40′**1-GHz flux/Jy:** 110**Spectral index:** 0.64**Size/arcmin:** 140×120**Type:** S

Has been called G160.5+2.8 and G160.4+2.8.

Radio: Large, filamentary shell.**Optical:** Incomplete shell.**X-ray:** Centrally brightened.**Point sources:** Pulsar within boundary of the remnant, plus several nearby compact radio sources.**Distance:** Various observations suggests less than 4 kpc.**References:**

d'Odorico & Sabbadin 1977, A&AS, 28, 439. Optical spectra.

Damashek *et al.* 1978, ApJ, 225, L31. Pulsar.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Dwarakanath *et al.* 1982, JApA, 3, 207. Radio observations at 34.5 MHz (26′×40′: S=750±150 Jy), plus review of flux densities.van Gorkom *et al.* 1982, MNRAS, 198, 757. WSRT H α absorption to nearby point source.

Sequist & Gilmore 1982, AJ, 87, 378. VLA observations of nearby source.

Leahy 1987, ApJ, 322, 917. Einstein observations.

Leahy & Roger 1991, AJ, 101, 1033. DRAO at 408 MHz (3′.5×4′.8) and 1.4 GHz (1′.0×1′.4), including H α and discussion of distance.

Yamauchi & Koyama 1993, PASJ, 45, 545. Hard X-ray observations.

Leahy & Aschenbach 1995, A&A, 293, 853. ROSAT observations.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Leahy *et al.* 1998, A&A, 339, 601. 232 MHz (3′.8×5′.2), 151 MHz (4′.2×5′.8) and Effelsberg 100-m at 4.7 GHz (2′.5) for spectral index studies.Roger *et al.* 1999, A&AS, 137, 7. 22 MHz flux density (S=1130±340 Jy).Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′.5: S=91±3 Jy).

Fürst & Reich 2004, in MIM, p141. Effelsberg 100-m at 1.4 and 2.7 GHz (9′.3), including polarisation.

Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′: S=102±7 Jy) and 1420 MHz (~1′: S=54.0±2.9 Jy), including polarisation and review of flux densities.

Leahy & Tian 2007, A&A, 461, 1013. CGPS at 408 MHz (2′.8×3′.9: S=117.8±5.3 Jy) and 1.4 GHz (58″×80″: S=65.9±3.4 Jy).

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5: S=34±3 Jy), including polarisation and review of flux densities.Gosachinskii 2013, AstL, 39, 179. H α observations of region.

Araya *et al.* 2014, MNRAS, 444, 860. Fermi detection.

G166.0+4.3

RA: 05^h26^m30^s
Dec: +42°56′

1-GHz flux/Jy: 7
Spectral index: 0.37

VRO 42.05.01
Size/arcmin: 55×35
Type: S

Radio: Two arcs of strikingly different radii.

Optical: Nearly complete ring.

X-ray: Predominantly in SW.

Distance: Hi indicates 4.5 kpc.

References:

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
Willis 1973, A&A, 26, 237. NRAO 300-ft at 2.7 GHz (5′:S=5.2±1.0 Jy).
Lozinskaya 1979, AujPh, 32, 113. H α interferometry.
Landecker *et al.* 1982, ApJ, 261, L41. DRAO at 1.4 GHz (1′0×1′4), plus review of flux densities.
Fesen *et al.* 1983, ApJS, 51, 337. Deep [OIII] imagery.
Pineault *et al.* 1985, A&A, 151, 52. VLA at 1.4 GHz (16″×20″) of part of remnant, and optical observations.
Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
Braun & Strom 1986, A&AS, 63, 345. WSRT Hi Observations.
Pineault *et al.* 1987, ApJ, 315, 580. DRAO and VLA combined at 1.4 GHz (20″).
Landecker *et al.* 1989, MNRAS, 237, 277. DRAO at 1.4 GHz (1′0×1′4), including Hi.
Burrows & Guo 1994, ApJ, 421, L19. ROSAT images and spectra.
Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
Guo & Burrows 1997, ApJ, 480, L51. ASCA observations.
Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
Leahy & Tian 2005, A&A, 440, 929. CGPS at 408 MHz (3′4×5′0) and 1.4 GHz (1′0×1′4), for spectral index studies.
see also: Tian & Leahy 2006, A&A, 451, 991. Erratum.
Kothes *et al.* 2006, A&A, 457, 1081. CGPS at 408 MHz (~3′:S=8.1±0.9 Jy) and 1420 MHz (~1′:S=5.1±0.4 Jy), including polarisation and review of flux densities.
Bocchino *et al.* 2009, A&A, 498, 139. XMM-Newton observations.
Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′5:S=3.3±0.3 Jy), including polarisation and review of flux densities.
Araya 2013, MNRAS, 434, 2202. Fermi observations.
Matsumura *et al.* 2017, PASJ, 69, 30. Suzaku observations.

G178.2–4.2

RA: 05^h25^m05^s
Dec: +28°11′

1-GHz flux/Jy: 2
Spectral index: 0.5

Size/arcmin: 72×62
Type: S

Radio: Faint shell, brighter in NE.

References:

- Gao *et al.* 2011, A&A, 532, A144. Urumqi 25-m at 5 GHz (9′5:S=1.0±0.1 Jy), plus other observations.

G179.0+2.6**RA:** 05^h53^m40^s**Dec:** +31°05′**1-GHz flux/Jy:** 7**Spectral index:** 0.4**Size/arcmin:** 70**Type:** S?**Radio:** Thick shell, with background extragalactic sources near centre.**Optical:** Nearly complete shell.**Point sources:** Pulsar near centre.**References:**

- Fürst & Reich 1986, A&A, 154, 303. Effelsberg 100-m at 1.4 (9′.4), 2.7 (4′.3) and 4.75 GHz (2′.4).
 Fürst *et al.* 1989, A&A, 223, 66. Observations of central, extragalactic source.
 Lorimer *et al.* 1998, A&A, 331, 1002. Pulsar search.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
 Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 3.2±0.3 Jy), including polarisation and review of flux densities.
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
 Pletsch *et al.* 2013, ApJ, 779, L11. Pulsar detection.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 How *et al.* 2018, MNRAS, 478, 1987. Optical observations.

G180.0–1.7

S147

RA: 05^h39^m00^s**Dec:** +27°50′**1-GHz flux/Jy:** 65**Spectral index:** varies**Size/arcmin:** 180**Type:** S**Radio:** Large faint shell, with spectral break.**Optical:** Wispy ring.**X-ray:** Possible detection.**Point sources:** Pulsar within boundary, with faint wind nebula.**Distance:** Various observations suggest about 1.2 kpc.**References:**

- van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.
 Sofue *et al.* 1980, PASJ, 32, 1. Effelsberg 100-m at 5 GHz (2′.6) of parts.
 Kundu *et al.* 1980, A&A, 92, 225. Effelsberg 100-m at 2.7 GHz (5′.5 : S = 34.9±4 Jy) and 1.6 GHz (10′ : S = 60.2±6 Jy).
 Angerhofer & Kundu 1981, AJ, 86, 1003. Arecibo at 430 MHz (9′ : S = 97±20 Jy).
 Fürst *et al.* 1982, A&A, 115, 428. Observations of compact radio sources near the remnant.
 Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.
 Fürst & Reich 1986, A&A, 163, 185. Effelsberg 100-m at 1.4, 2.7 and 4.7/5.0 GHz (9′.4, 4′.3 and 2′.4/2′.6).
 Sauvageot *et al.* 1990, A&A, 227, 183. EXOSAT possible detection.
 Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.
 Anderson *et al.* 1996, ApJ, 468, L55. Pulsar detection.
 Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz.
 Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′.5 : S = 77±10 Jy).
 Romani & Ng 2003, ApJ, 585, L41. Chandra of pulsar.
 Kramer *et al.* 2003, ApJ, 593, L31. Pulsar observations.
 Sallmen & Welsh 2004, A&A, 426, 555. Optical absorption towards background stars.
 Drew *et al.* 2005, MNRAS, 362, 753. H α imaging.
 Ng *et al.* 2007, ApJ, 654, 487. Pulsar and wind nebula observations.

Xiao *et al.* 2008, A&A, 482, 783. Urumqi 25-m at 4.8 GHz (9'5 : S = 15.4±3.0 Jy) and Effelsberg 100-m at 2.6 GHz (4'4 : S = 34.6±4.0 Jy).
 Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.
 Katsuta *et al.* 2012, ApJ, 752, 135. Fermi observations.
 Dinçel *et al.* 2015, MNRAS, 448, 3196. Identification of OB runaway star near centre.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Boubert *et al.* 2017, A&A, 606, A14. Gaia search for runaway progenitor companion.
 Chen *et al.* 2017, MNRAS, 472, 3924. Optical and IR observations, including distance.
 Ren *et al.* 2018, RAA, 18, 111. Optical spectroscopy.

G181.1 + 9.5

RA: 06^h26^m40^s
Dec: +32°30'

1-GHz flux/Jy: ?
Spectral index: 0.45?

Size/arcmin: 74
Type: S

Radio: Faint shell.

X-ray: Detected.

Distance: H α observations suggest 0.5--2.5 kpc.

References:

Kothes *et al.* 2017, A&A, 597, A116. DRAO at 1.4 GHz (50''×90'') including H α and Effelsberg 100-m at 4850 MHz (2'45) including polarisation, plus ROSAT survey observations.

G182.4 + 4.3

RA: 06^h08^m10^s
Dec: +29°00'

1-GHz flux/Jy: 0.5
Spectral index: 0.4

Size/arcmin: 50
Type: S

Radio: Incomplete shell.

Optical: Brighter in S and NW.

X-ray: Diffuse emission.

References:

Kothes *et al.* 1998, A&A, 331, 661. Effelsberg 100-m at 1.4, 2.7, 4.9 and 10.5 GHz (9'4 : S = 0.36±0.08 Jy, 4'4 : S = 0.25±0.04 Jy, 2'5 : S = 0.20±0.02 Jy and 1'2 : S = 0.15±0.03 Jy), plus X-ray upper limit.

Reich 2002, in NSPS, p1. Effelsberg 100-m at 2.7 GHz and 4.9 GHz (3').

Sun *et al.* 2011, A&A, 536, A83. Urumqi 25-m at 5 GHz (9'5 : S = 0.26±0.5 Jy) including polarisation and review of flux densities.

Sezer *et al.* 2012, MNRAS, 427, 1168. Optical and XMM-Newton observations.

Jeong *et al.* 2012, Ap&SS, 342, 389. CO observations of region.

G184.6–5.8

Crab Nebula, 3C144, SN1054

RA: 05^h34^m31^s**1-GHz flux/Jy:** 900**Size/arcmin:** 7×5**Dec:** +22°01′**Spectral index:** 0.30**Type:** F

This is the remnant of the SN of AD1054.

Radio: Filled-centre, central pulsar, with faint ‘jet’ (or tube) extending from the N edge; shows secular decline.

Optical: Strongly polarised filaments, diffuse synchrotron emission, with ‘jet’ faintly visible.

X-ray: Central ‘torus’ around the pulsar.

Point sources: Pulsar powering the remnant.

Distance: Proper motions and radial velocities give 2 kpc.

References:

- Velusamy 1984, *Nature*, 308, 251. VLA at 1.4 GHz, radio detection of ‘jet’.
- Velusamy 1985, *MNRAS*, 212, 359. VLA at 1.4 GHz (15’’).
- Aller & Reynolds 1985, *ApJ*, 293, L73. Radio flux density decrease.
- Fesen & Gull 1986, *ApJ*, 306, 259. Deep [OIII] imagery of ‘jet’.
- Marcelin *et al.* 1990, *A&A*, 228, 471. Optical of ‘jet’.
- Bietenholz & Kronberg 1990, *ApJ*, 357, L13. VLA at 1.4 GHz (1’’8).
- Hester *et al.* 1990, *ApJ*, 357, 539. Optical and IR images.
- Hickson & van den Bergh 1990, *ApJ*, 365, 224. Optical polarisation.
- Bietenholz & Kronberg 1992, *ApJ*, 393, 206. VLA at 1.5 and 5 GHz (1’’8) and 1.5 and 14 GHz (6’’5) for spectral studies.
- Fesen & Staker 1993, *MNRAS*, 263, 69. [OIII] imaging of ‘jet’, and proper motion studies.
- Kassim *et al.* 1993, *AJ*, 106, 2218. VLA at 74 MHz (20’’).
- Véron-Cetty & Woltjer 1993, *A&A*, 270, 370. Continuum and [OIII] photometry.
- Wallace *et al.* 1994, *A&A*, 286, 565. H α of surroundings.
- Frail *et al.* 1995, *ApJ*, 454, L129. VLA at 333 MHz (20’’) for limits on shell.
- Bietenholz *et al.* 1997, *ApJ*, 490, 291. Comparison of VLA observations at 74 MHz, 327 MHz, 1.5 GHz and 5 GHz for spectral index studies.
- Blair *et al.* 1997, *ApJS*, 109, 473. HST imaging.
- Fesen *et al.* 1997, *AJ*, 113, 354. Limits on H α halo.
- Nugent 1998, *PASP*, 110, 831. Optical expansion.
- Sankrit *et al.* 1998, *ApJ*, 504, 344. HST images.
- Wallace *et al.* 1999, *ApJS*, 124, 181. DRAO at 1.4 GHz (1’0×2’8), plus Effelsberg 100-m, for H α studies.
- Greiveldinger & Aschenbach 1999, *ApJ*, 510, 305. X-ray variability of torus.
- Weisskopf *et al.* 2000, *ApJ*, 536, L81. Chandra observations.
- Sollerman *et al.* 2000, *ApJ*, 537, 861. HST observations.
- Aharonian *et al.* 2000, *ApJ*, 539, 317. HESS observations.
- Willingale *et al.* 2001, *A&A*, 365, L212. XMM-Newton observations.
- Douvion *et al.* 2001, *A&A*, 373, 281. ISO observations.
- Bietenholz *et al.* 2001, *ApJ*, 560, 254. Multi-epoch VLA observations, showing variations near the pulsar.
- Bandiera *et al.* 2002, *A&A*, 386, 1044. 1.3 mm observations.
- Hester *et al.* 2002, *ApJ*, 577, L49. HST and Chandra multi-epoch observations.
- Atkins *et al.* 2003, *ApJ*, 595, 803. γ -ray observations.
- Green *et al.* 2004, *MNRAS*, 355, 1315. Sub-mm and ISO observations.
- Mori *et al.* 2004, *ApJ*, 609, 186. Chandra observations.
- Čadež *et al.* 2004, *ApJ*, 609, 797. Optical observations.
- Bietenholz *et al.* 2004, *ApJ*, 615, 794. VLA at 5 GHz (1’’4) and HST multi-epoch observation for proper motion studies.

Melatos *et al.* 2005, ApJ, 633, 931. Multi-epoch near-IR observations central region.
 Seward *et al.* 2006, ApJ, 636, 873. Chandra observations of scattering halo.
 Temim *et al.* 2006, AJ, 132, 1610. Spitzer observations.
see also: Temim *et al.* 2009, AJ, 137, 5155. Erratum.
 Aharonian *et al.* 2006, A&A, 457, 899. HESS observations
 Seward *et al.* 2006, ApJ, 652, 1277. Chandra observations.
 MacAlpine *et al.* 2007, AJ, 133, 81. Optical spectroscopy.
 Kaplan *et al.* 2008, ApJ, 677, 1201. HST proper motion of pulsar.
 Rudie *et al.* 2008, MNRAS, 384, 1200. [OIII] observations of 'jet', for proper motion.
 Hurley-Walker *et al.* 2009, MNRAS, 396, 365. Radio observations at 14 to 18 GHz.
 Tziamtzis *et al.* 2009, A&A, 497, 167. Limits on H α halo.
 Carlebois *et al.* 2010, AJ, 139, 2083. Optical imaging spectroscopy.
 Amount *et al.* 2010, A&A, 514, A70. IRAM 30-m at 150 GHz (16''7 : S = 244 \pm 24 Jy) and GBT at 90 GHz (9''3).
 Satterfield *et al.* 2012, AJ, 144, 27. Optical spectroscopy.
 Temim *et al.* 2012, ApJ, 753, 72. Spitzer observations of dust.
 Gomez *et al.* 2012, ApJ, 760, 96. Herschel IR and sub-mm observations of dust.
 Loh *et al.* 2012, MNRAS, 421, 789. IR observations of H $_2$.
 Lundqvist & Tziamtzis 2012, MNRAS, 423, 1571. Optical limits on outer shell.
 Loll *et al.* 2013, ApJ, 765, 152. HST optical line and continuum observations.
 Barlow *et al.* 2013, Science, 342, 1343. Herschel far-IR spectroscopy.
 Black & Fesen 2015, MNRAS, 447, 2540. [OIII] observations of 'jet'.
 Owen & Barlow 2015, ApJ, 801, 141. Herschel and Spitzer observations.
 Bietenholz & Nugent 2015, MNRAS, 454, 2416. Multi-epoch VLA and optical observations for expansion study.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 9 frequencies between 30 and 857 GHz.
 Trotter *et al.* 2017, MNRAS, 469, 1299. Time evolution of radio emission.
 Dubner *et al.* 2017, ApJ, 840, 82. VLA, HST and Chandra observations, plus ALMA of central region, and UV observations.
 Ritacco *et al.* 2018, A&A, 616, A35. Observations 150 GHz (18''), including polarisation.

G189.1+3.0

IC443, 3C157

RA: 06^h17^m00^s**1-GHz flux/Jy:** 165**Size/arcmin:** 45**Dec:** +22°34'**Spectral index:** 0.36**Type:** C**Radio:** Limb-brightened to NE, with faint extension to the E.**Optical:** Brightest to the NE, with faint filaments outside the NE boundary.**X-ray:** Shell, brightest to the NE, plus compact source with nebula.**Point sources:** X-ray source and nebula in S.**Distance:** Optical observations imply 1.9 kpc.**References:**

Fesen 1984, ApJ, 281, 658. Optical of filament to far NE.

Erickson & Mahoney 1985, ApJ, 290, 596. TPT at 4 frequencies between 31 and 74 MHz (11'x13' to 4'7x5'4), plus review of flux densities.

Braun & Strom 1986, A&A, 164, 193. WSRT at 327 MHz (72''x185'') and 1.4 GHz (17''x43''), plus H α and IRAS.

Green 1986, MNRAS, 221, 473. 151 MHz observations (1'2x3'1) and Half-Mile Telescope at 1.4 GHz (2'1x5'4).

Mufson *et al.* 1986, AJ, 92, 1349. Radio, IR, optical, UV and X-ray comparison, including VLA at 1.6 GHz (3''3x3''8 and 40'').Petre *et al.* 1988, ApJ, 335, 215. Einstein and other X-ray observations.

- Dickel *et al.* 1989, AJ, 98, 1363. VLA at 1.4 GHz ($1''.1 \times 1''.2$) of NE.
- Wood *et al.* 1991, AJ, 102, 224. VLA at 5 GHz ($3''.6 \times 3''.8$) of northeast, including polarisation.
- Asaoka & Aschenbach 1994, A&A, 284, 573. X-ray, including possible overlapping remnant.
- Tauber *et al.* 1994, ApJ, 421, 570. Observations of shocked molecular species.
- Claussen *et al.* 1999, ApJ, 522, 349. High resolution observations of OH masers.
- Rho *et al.* 2001, ApJ, 547, 885. ISO and 2Mass IR spectroscopy and imaging.
- Olbert *et al.* 2001, ApJ, 554, L205. Chandra of compact X-ray source and surrounding nebula.
- Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz ($14'.5 : S = 160 \pm 5$ Jy).
- Welsh & Sallmen 2003, A&A, 408, 545. Optical absorption studies.
- Leahy 2004, AJ, 127, 2277. DRAO at 408 MHz ($3'.3 \times 8'.6$) and 1.4 GHz ($1'.0 \times 2'.6$), for spectral index studies.
- see also:* Leahy 2004, AJ, 128, 1478. Addendum.
- Bykov *et al.* 2005, ApJ, 624, L41. Chandra observations of compact source.
- Gaensler *et al.* 2006, ApJ, 648, 1037. Chandra of X-ray source and nebula.
- Hewitt *et al.* 2006, ApJ, 652, 1288. GBT at 1.6 and 1.7 GHz ($7''.2$) for OH, and VLA at 330 MHz ($64'' \times 74''$).
- Rosado *et al.* 2007, AJ, 133, 89. Observations of shocked H₂.
- Albert *et al.* 2007, ApJ, 664, L87. γ -ray observations.
- Troja *et al.* 2008, A&A, 485, 777. XMM-Newton observations.
- Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.
- Bykov *et al.* 2008, ApJ, 676, 1050. XMM-Newton, Chandra and Spitzer observations.
- Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
- Lee *et al.* 2008, AJ, 135, 796. VLA at 1.4 GHz ($39'' \times 42''$) including Hi.
- Bocchino *et al.* 2008, AdSpR, 41, 396. INTEGRAL observations.
- Bocchino *et al.* 2009, A&A, 498, 139. XMM-Newton observations.
- Acciari *et al.* 2009, ApJ, 698, L133. γ -ray observations.
- Yamaguchi *et al.* 2009, ApJ, 705, L6. Suzaku observations.
- Koo *et al.* 2010, AJ, 140, 262. Hi Zeeman splitting observations.
- Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz ($9'.5 : S = 85 \pm 9$ Jy), including polarisation and review of flux densities.
- Castelletti *et al.* 2011, A&A, 534, A21. VLA at 74 MHz ($35'' : S = 470 \pm 51$ Jy) and 330 MHz ($17'' : S = 248 \pm 15$ Jy), plus review of flux densities.
- Yuan & Neufeld 2011, ApJ, 726, 76. Spitzer observations.
- Xu *et al.* 2011, ApJ, 727, 81. CO observations of region.
- Shinn *et al.* 2011, ApJ, 732, 124. Akari observations of H₂.
- Lee *et al.* 2012, ApJ, 749, 34. CO and HCO⁺ of region.
- Taylor *et al.* 2012, ApJ, 750, L15. Optical absorption of background stars.
- Yuan *et al.* 2012, ApJ, 753, 126. Spitzer spectroscopy.
- Ackermann *et al.* 2013, Science, 339, 807. Fermi observations.
- Hezareth *et al.* 2013, A&A, 558, A45. CO observations, including polarisation, of region.
- Kokusho *et al.* 2013, ApJ, 768, L8. IR observations of [FeII], plus Akari and Spitzer observations.
- Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
- Ohnishi *et al.* 2014, ApJ, 784, 74. Suzaku observations.
- Yamaguchi *et al.* 2014, ApJ, 785, L27. Suzaku observations.
- Su *et al.* 2014, ApJ, 788, 122. CO observations.
- Gusdorf *et al.* 2014, IAUS, 296, 178. CO observations.
- Mitra *et al.* 2014, IAUS, 296, 376. Combined GMRT and CLFST observations at 150 MHz ($19'' \times 24''$).
- Swartz *et al.* 2015, ApJ, 808, 84. Chandra observations of compact source and nebula.
- Kokusho *et al.* 2015, P&SS, 116, 92. IR [FeII] observations.
- Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 9 frequencies between 30 and 857 GHz.
- Kilpatrick *et al.* 2016, ApJ, 816, 1. CO observations, including broad lines.
- Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
- McEwen *et al.* 2016, ApJ, 826, 189. NH₃ and CH₃OH observations.

Aceró *et al.* 2016, ApJS, 224, 8. Fermi observations.

Egron *et al.* 2017, MNRAS, 470, 1329. SRT at 1.5 GHz ($11' : S = 134 \pm 4$ Jy) and 7 GHz ($2'7 : S = 67 \pm 3$ Jy).

Ambrocio-Cruz *et al.* 2017, MNRAS, 472, 51. Optical observations of NW, for distance.

Madsen *et al.* 2017, ApJ, 841, 56. NuSTAR observations.

Greco *et al.* 2018, A&A, 615, A157. XMM-Newton observations.

Zhang *et al.* 2018, ApJ, 859, 141. NuSTAR, XMM-Newton and Chandra observations.

G190.9–2.2

RA: 06^h01^m55^s
Dec: +18°24'

1-GHz flux/Jy: 1.3?
Spectral index: 0.7?

Size/arcmin: 70×60
Type: S

Radio: Incomplete shell.

References:

Foster *et al.* 2013, A&A, 549, A107. Various radio survey observations.

G205.5+0.5

RA: 06^h39^m00^s
Dec: +06°30'

1-GHz flux/Jy: 140
Spectral index: 0.4

Monoceros Nebula
Size/arcmin: 220
Type: S

Radio: In complex region, parts may be H_{II} regions.

Optical: Large ring, near Rosette nebula.

X-ray: Possibly detected.

Distance: Association with molecular cloud suggests 1.6 to 2.0 kpc.

References:

Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (9').

Velusamy & Kundu 1974, A&A, 32, 375. NRAO 300-ft at 2.7 GHz (5'), part only.

Dickel & DeNoyer 1975, AJ, 80, 437. Arecibo at 111 MHz ($1' : S = 462 \pm 180$ Jy) and $S_{610 \text{ MHz}} = 245$ Jy.

Davies *et al.* 1978, A&AS, 31, 271. Deep optical plates.

Lozinskaya 1981, SvAL, 7, 17. Mean optical velocity.

Graham *et al.* 1982, A&A, 109, 145. Effelsberg 100-m at 2.7 GHz ($4'4 : S = 97.6 \pm 12.5$ Jy), plus review of flux densities.

Fesen *et al.* 1985, ApJ, 292, 29. Optical spectra.

Leahy *et al.* 1986, MNRAS, 220, 501. Einstein observations.

Odegard 1986, ApJ, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24', 19' and 16').

Esposito *et al.* 1996, ApJ, 461, 820. Possible associated γ -ray emission.

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Oliver *et al.* 1996, A&A, 315, 578. CO observations of some of surroundings.

Jaffe *et al.* 1997, ApJ, 484, L129. γ -ray detection.

Aharonian *et al.* 2004, A&A, 417, 973. HESS limit.

Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.

Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.

Xiao & Zhu 2012, A&A, 545, A86. Review of radio, H_I and H α observations.

Dirks & Meyer 2016, ApJ, 819, 45. Time variation of optical line absorption.

Katagiri *et al.* 2016, ApJ, 831, 106. Fermi observations.

Aceró *et al.* 2016, ApJS, 224, 8. Fermi observations.

Boubert *et al.* 2017, A&A, 606, A14. Gaia search for runaway progenitor companion.

Su *et al.* 2017, ApJ, 836, 211. CO observations.

Zhao *et al.* 2018, ApJ, 855, 12. IR observations.

G206.9+2.3

PKS 0646+06

RA: 06^h48^m40^s
Dec: +06°26′**1-GHz flux/Jy:** 6
Spectral index: 0.5**Size/arcmin:** 60×40
Type: S?**Radio:** Diffuse source near the Monoceros Nebula.**Optical:** Filaments detected.**X-ray:** Possibly detected.**References:**

Davies & Meaburn 1978, A&A, 69, 443. Optical observations.

Nousek *et al.* 1981, Apj, 248, 152. HEAO-1 X-ray limit.Graham *et al.* 1982, A&A, 109, 145. Effelsberg 100-m at 2.7 GHz (4′.4 : S = 4.1±0.6 Jy), plus review of flux densities.

Rosado 1982, RMxAA, 5, 127. Optical observations.

Fesen *et al.* 1985, Apj, 292, 29. Optical spectra.

Leahy 1986, A&A, 156, 191. Einstein observations.

Odegard 1986, Apj, 301, 813. TPT at 20.6, 25.6 and 30.9 MHz (24′, 19′ and 16′).

Biggs & Lyne 1996, MNRAS, 282, 691. Pulsar search.

Gao *et al.* 2011, A&A, 529, A159. Urumqi 25-m at 5 GHz (9′.5 : S = 2.9±0.3 Jy), including polarisation and review of flux densities.Ambrocio-Cruz *et al.* 2014, RMxAA, 50, 323. [SII] spectroscopy.Su *et al.* 2017, Apj, 836, 211. CO observations.**G213.0–0.6****RA:** 06^h50^m50^s
Dec: –00°30′**1-GHz flux/Jy:** 21
Spectral index: 0.4**Size/arcmin:** 160×140?
Type: S

Has also been called G213.3–0.4.

Radio: Large, faint shell.**Optical:** Filamentary shell.**Point sources:** Central X-ray source.**Distance:** Association with molecular cloud suggests about 1 kpc.**References:**Reich *et al.* 2003, A&A, 408, 961. Effelsberg 100-m at 863 MHz (14′.5 : S = 22.0±3.7 Jy).Stupar & Parker 2012, MNRAS, 419, 1413. H α , radio survey and other observations.Su *et al.* 2017, Apj, 836, 211. CO observations.

G260.4–3.4

Puppis A, MSH 08–44

RA: 08^h22^m10^s**1-GHz flux/Jy:** 130**Size/arcmin:** 60×50**Dec:** –43°00′**Spectral index:** 0.5**Type:** S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Angular shell, brightest to the E, poorly defined to the W.

Optical: Nebulosity and wisps.

X-ray: Brightest to the E.

Point sources: Central possible pulsating X-ray source.

Distance: Association with H_I implies 1.3 kpc.

References:

Green 1971, *AJPh*, 24, 773. Molonglo at 408 MHz (3′:S=198±20 Jy).

Goudis & Meaburn 1978, *A&A*, 62, 283. H α + [NII] optical image.

Petre *et al.* 1982, *ApJ*, 258, 22. Einstein observations.

Milne *et al.* 1983, *MNRAS*, 204, 237. FIRST at 1415 MHz (50″), large scale emission missing.

Teske & Petre 1987, *ApJ*, 318, 370. Coronal optical line emission.

Dubner & Arnal 1988, *A&AS*, 75, 363. H_I and CO observations of surroundings.

Arendt *et al.* 1990, *ApJ*, 350, 266. MOST at 843 MHz (44″×65″), with large scale emission added, plus IR, optical and X-ray observations.

Dubner *et al.* 1991, *AJ*, 101, 1466. VLA at 327 MHz (43″×73″) and 1.5 GHz (43″×77″).

Arendt *et al.* 1991, *ApJ*, 368, 474. IR observations.

Milne *et al.* 1993, *MNRAS*, 261, 366. Parkes 64-m at 4.75 (4′.5 :S=59±5 Jy) and 8.4 GHz (3′ :S=38±4 Jy), plus polarisation.

Berthiaume *et al.* 1994, *ApJ*, 425, 132. X-ray spectroscopy.

Sutherland & Dopita 1995, *ApJ*, 439, 365. Spectrophotometry.

Reynoso *et al.* 1995, *AJ*, 110, 318. VLA at 1.4 GHz (90″) including neutral hydrogen.

Blair *et al.* 1995, *ApJ*, 454, L35. Far UV spectroscopy.

Petre *et al.* 1996, *ApJ*, 465, L43. ROSAT of central source.

Bock *et al.* 1998, *AJ*, 116, 1886. MOST at 843 MHz (43″×60″).

Pavlov *et al.* 1999, *ApJ*, 511, L45. Possible pulsation detection from central X-ray source.

Zavlin *et al.* 1999, *ApJ*, 525, 959. X-ray observations of central source.

Bocchino *et al.* 2000, *A&A*, 359, 316. Optical studies of selected filaments in N.

Woermann *et al.* 2000, *MNRAS*, 317, 421. OH observations.

Gaensler *et al.* 2000, *ApJ*, 537, L35. Radio limit for nebula around possible pulsar.

Reynoso *et al.* 2003, *MNRAS*, 345, 671. ATCA at 1.4 GHz (90″) for H_I near central X-ray source.

Hwang *et al.* 2005, *ApJ*, 635, 355. Chandra observations of E edge.

Hui & Becker 2006, *A&A*, 454, 543. XMM-Newton and Chandra observations.

Hui & Becker 2006, *A&A*, 457, L33. Chandra proper motion study of central source.

Castelletti *et al.* 2006, *A&A*, 459, 535. VLA at 1.4 GHz (16″×34″ :S=114±8 Jy) and 327 MHz (45″×90″ :S=263±20 Jy).

Winkler & Petre 2007, *ApJ*, 670, 635. Chandra proper motion study of central source.

Paron *et al.* 2008, *A&A*, 480, 439. CO observations of E.

Hwang *et al.* 2008, *ApJ*, 676, 378. Suzaku observations.

Katsuda *et al.* 2008, *ApJ*, 678, 297. XMM-Newton observations.

Mignani *et al.* 2009, *A&A*, 500, 1211. Optical limits for compact X-ray source.

Katsuda *et al.* 2010, *ApJ*, 714, 1725. Chandra and XMM-Newton observations.

Arendt *et al.* 2010, *ApJ*, 725, 585. Spitzer observations.

Becker *et al.* 2012, *ApJ*, 755, 141. Chandra proper motion study of central X-ray source.

Katsuda *et al.* 2012, *ApJ*, 756, 49. XMM-Newton spectroscopy.

Hewitt *et al.* 2012, *ApJ*, 759, 89. Fermi and WMAP 23 to 90 GHz observations.

Dubner *et al.* 2013, *A&A*, 555, A9. XMM-Newton and Chandra observations.

Katsuda *et al.* 2013, *ApJ*, 768, 182. XMM-Newton observations.

Abramowski *et al.* 2015, A&A, 575, A81. HESS limit.
 Reynoso & Walsh 2015, MNRAS, 451, 3044. ATCA at 1.4 GHz (51'' \times 82'') and 1.7 GHz for spectral index study.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 4 frequencies between 30 and 100 GHz.
 Luna *et al.* 2016, A&A, 590, A70. XMM-Newton and Chandra observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Reynoso *et al.* 2017, MNRAS, 464, 3029. ATCA at 1.4 GHz including H α .
 Garca *et al.* 2017, A&A, 604, L5. XMM-Newton observations of SW.
 Slane *et al.* 2018, ApJ, 865, 86. XMM-Newton and Chandra observations of Vela X.
 Reynoso *et al.* 2018, MNRAS, 477, 2087. ATCA at 1.3 to 2.6 GHz, including polarisation.

G261.9+5.5

RA: 09^h04^m20^s
Dec: -38°42'

1-GHz flux/Jy: 10?
Spectral index: 0.4?

Size/arcmin: 40 \times 30
Type: S

Radio: Faint shell with little limb brightening.

References:

Hill 1967, AujPh, 20, 297. Parkes 64-m at 2650 MHz (7'5 : S = 7 Jy) also $S_{1410 \text{ MHz}} = 8 \text{ Jy}$, $S_{81.5 \text{ MHz}} = 25 \text{ Jy}$.
 Colomb & Dubner 1980, A&A, 82, 244. Argentine 30-m dish at 1.4 GHz, for H α possibly associated with remnant.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 71'').
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G263.9-3.3

RA: 08^h34^m00^s
Dec: -45°50'

1-GHz flux/Jy: 1750
Spectral index: varies

Vela (XYZ)
Size/arcmin: 255
Type: C

This refers to the whole Vela XYZ complex, of which X has at times been classified as a separate (filled-centre) remnant. This remnant is overlapped by G260.4-3.4 and G266.2-1.2.

Radio: Large shell, with flatter spectrum component (Vela X), and pulsar nebula.

Optical: Filaments.

X-ray: Patchy shell, with extensions, central nebula and pulsar.

Point sources: Pulsar within Vela X, with one-sided 'jet'.

Distance: Vela pulsar parallax gives 0.3 kpc, optical spectra and H α studies suggest 0.25 kpc.

References:

Milne 1968, AujPh, 21, 201. Parkes 64-m at various frequencies, including 408 MHz (48' : S = 2300 \pm 300 Jy), 635 MHz (31' : S = 2360 \pm 300 Jy), 1410 MHz (14' : S = 1640 \pm 300 Jy) and 2650 MHz (7'5 : S = 1400 \pm 250 Jy), plus discussion of the distance.
 Milne 1980, A&A, 81, 293. Maps of Vela X with Parkes 64-m at 1.66, 2.7 and 5 GHz (12', 8'.4 and 4'.4).

- Weiler & Panagia 1980, A&A, 90, 269. Clarification of notation of this region and review previous observations.
- Bignami & Caraveo 1988, ApJ, 325, L5. Pulsar proper motion from optical observations.
- Dwarakanath 1991, JApA, 12, 199. Gauribidanur 'T' array at 34.5 MHz ($26' \times 84'$; $S = 1800$ Jy for Vela X and $S = 3900$ Jy for Vela YZ), plus review of flux densities.
- Bietenholz *et al.* 1991, ApJ, 376, L41. VLA at 5 GHz in vicinity of pulsar.
- Willmore *et al.* 1992, MNRAS, 254, 139. Hard X-ray observations.
- Dubner *et al.* 1992, A&AS, 96, 505. Argentine 30-m at 1.4 GHz ($30'$) of surrounding Hi.
- Strom *et al.* 1995, Nature, 373, 590. Radio of X-ray extensions.
- Markwardt & Ögelman 1995, Nature, 375, 40. X-ray jet from pulsar.
- Jenkins & Wallerstein 1995, ApJ, 440, 227. Optical absorption of associated neutral carbon cloud.
- Milne 1995, MNRAS, 277, 1435. Parkes 64-m at 8.4 GHz ($3'$), including polarisation, of Vela X.
- Danks & Sembach 1995, AJ, 109, 2627. Optical spectroscopy of background stars.
- Blair *et al.* 1995, AJ, 110, 312. UV spectroscopy.
- Duncan *et al.* 1996, MNRAS, 280, 252. Parkes 64-m at 2.4 GHz ($8'9$).
- Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 62''$) of part.
- Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
- Frail *et al.* 1997, ApJ, 475, 224. VLA at 330 MHz ($1' \times 1'8$) and comparison with ROSAT observations of Vela X.
- Markwardt & Ögelman 1997, ApJ, 480, L13. ASCA observations of pulsar 'jet'.
- Raymond *et al.* 1997, ApJ, 482, 881. UV spectroscopy.
- Yoshikoshi *et al.* 1997, ApJ, 487, L65. γ -ray observations of pulsar.
- Jenkins *et al.* 1998, ApJ, 492, L147. UV absorption to background star.
- Dubner *et al.* 1998, AJ, 116, 813. Parkes 64-m at 1.4 GHz ($15'$) for Hi studies.
- Bock *et al.* 1998, AJ, 116, 1886. MOST at 843 MHz ($43'' \times 60''$).
- Cha *et al.* 1999, ApJ, 515, L25. Optical spectra, for distance.
- Lu & Aschenbach 2000, A&A, 362, 1083. ROSAT spatially resolved spectroscopy.
- Chadwick *et al.* 2000, ApJ, 537, 414. Limit on high energy γ -rays from pulsar.
- Sankrit *et al.* 2001, ApJ, 549, 416. Far-UV observations of selected region.
- Pavlov *et al.* 2001, ApJ, 554, L189. Chandra two epoch observations of pulsar nebula.
- Helfand *et al.* 2001, ApJ, 556, 380. Chandra observations of pulsar and its nebula.
- Miyata *et al.* 2001, ApJ, 559, L45. Chandra of X-ray extension.
- Alvarez *et al.* 2001, A&A, 372, 636. Radio spectral index studies.
see also: Alvarez *et al.* 2001, A&A, 379, 323. Erratum.
- Caraveo *et al.* 2001, ApJ, 561, 930. HST parallax observations of pulsar.
- Moriguchi *et al.* 2001, PASJ, 53, 1025. CO observations.
- Dodson *et al.* 2003, MNRAS, 343, 116. ATCA at 1.4, 2.4, 5.2 and 8.5 GHz ($6''2 \times 8''1$, $26'' \times 36''$, $10''5 \times 12''1$ and $10''6 \times 11''2$) of pulsar nebula.
- Sankrit *et al.* 2003, ApJ, 589, 242. Optical nebulosity to NE of G266.2–1.2.
- Nichols & Slavin 2004, ApJ, 610, 285. UV absorption toward background sources.
- Hales *et al.* 2004, ApJ, 613, 977. Vela X at 31 GHz ($4'1$).
- Mongano *et al.* 2005, A&A, 436, 917. XMM-Newton and other X-ray observations of pulsar nebula.
- Miceli *et al.* 2005, A&A, 442, 513. XMM-Newton observations of N rim.
- Katsuda & Tsunemi 2005, PASJ, 57, 621. XMM-Newton observations of E.
- Aharonian *et al.* 2006, A&A, 448, L43. HESS observations.
- McConnell *et al.* 2006, AJ, 131, 648. ATCA at 4.9 GHz ($12'$) including polarisation.
- Katsuda & Tsunemi 2006, ApJ, 642, 917. XMM-Newton observations of NE.
- Nishikida *et al.* 2006, ApJ, 644, L171. Far UV observations.
- Miceli *et al.* 2008, ApJ, 676, 1064. XMM-Newton observations.
- LaMassa *et al.* 2008, ApJ, 689, L121. XMM-Newton observations of Vela X.
- Grondin *et al.* 2013, ApJ, 774, 110. Fermi observations of Vela X.
- Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30, 44 and 70 GHz.
- Rao *et al.* 2016, MNRAS, 455, 2529. Time variation of optical line absorption.
- Abdalla *et al.* 2018, A&A, 612, A1. HESS observations of Vela X.

G266.2–1.2

RX J0852.0–4622

RA: 08^h52^m00^s**1-GHz flux/Jy:** 50?**Size/arcmin:** 120**Dec:** –46°20′**Spectral index:** 0.3?**Type:** S

This remnant overlaps the Vela SNR (G263.9–3.3).

Radio: Incomplete shell, confused by the Vela SNR.

Optical: Nebulosity offset to NE.

X-ray: Non-thermal shell, confused by the Vela SNR, with central source, and possible associated pulsar.

Point sources: Central X-ray source, with optical nebula, and possible associated pulsar.

Distance: X-ray data suggest an upper limit of 1 kpc.

References:

- Aschenbach 1998, *Nature*, 396, 141. ROSAT identification.
- Iyudin *et al.* 1998, *Nature*, 396, 142. γ -ray observations.
- Combi *et al.* 1999, *ApJ*, 519, L177. Radio observations.
- Duncan & Green 2000, *A&A*, 364, 732. Parkes 64-m at 1.4 GHz (14'0), and comparison with other observations.
- Redman *et al.* 2000, *ApJ*, 543, L153. Optical of nearly nebulosity.
- Tsunemi *et al.* 2000, *PASJ*, 52, 887. ASCA spectral observations.
- Slane *et al.* 2001, *ApJ*, 548, 814. ASCA observations.
- Mereghetti *et al.* 2001, *ApJ*, 548, L213. BeppoSAX observations of central sources.
- Pavlov *et al.* 2001, *ApJ*, 559, L131. Chandra of central X-ray source.
- Moriguchi *et al.* 2001, *PASJ*, 53, 1025. CO observations.
- Pellizzoni *et al.* 2002, *A&A*, 393, L65. Optical observations of central source.
- Redman *et al.* 2002, *MNRAS*, 336, 1093. Optical nebulosity to NE.
- Kargaltsev *et al.* 2002, *ApJ*, 580, 1060. Chandra observations of central source.
- Sankrit *et al.* 2003, *ApJ*, 589, 242. Optical nebulosity to NE.
- Redman & Meaburn 2005, *MNRAS*, 356, 969. Possible pulsar association.
- Iyudin *et al.* 2005, *A&A*, 429, 225. XMM-Newton observations.
- Aharonian *et al.* 2005, *A&A*, 437, L7. HESS observations.
- Katagiri *et al.* 2005, *ApJ*, 619, L163. γ -ray observations.
- Bamba *et al.* 2005, *ApJ*, 632, 294. Chandra of NW rim.
- Reynoso *et al.* 2006, *A&A*, 449, 243. ATCA at 1.38 GHz (32'' \times 37'').
- Enomoto *et al.* 2006, *ApJ*, 652, 1268. γ -ray observations.
- Mignani *et al.* 2007, *A&A*, 473, 883. Deep optical observations of compact X-ray source.
- Aharonian *et al.* 2007, *ApJ*, 661, 236. HESS observations.
- Katsuda *et al.* 2008, *ApJ*, 678, L35. XMM-Newton proper motion study.
- Pannuti *et al.* 2010, *ApJ*, 721, 1492. Chandra observations of NW.
- Allen *et al.* 2015, *ApJ*, 798, 82. Two epoch Chandra observations for expansion.
- Acerio *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
- Takeda *et al.* 2016, *PASJ*, 68, S10. Suzaku observations.
- Fukui *et al.* 2017, *ApJ*, 850, 71. CO and H α observations.
- Maxted *et al.* 2018, *ApJ*, 866, 76. ATCA at 1.4 to 2.9 GHz, plus other observations.
- Abdalla *et al.* 2018, *A&A*, 612, A7. HESS observations.

G272.2–3.2**RA:** 09^h06^m50^s
Dec: –52°07′**1-GHz flux/Jy:** 0.4
Spectral index: 0.6**Size/arcmin:** 15?
Type: S?**Radio:** Diffuse shell.**Optical:** Detected.**X-ray:** Centrally brightened.**References:**Greiner *et al.* 1994, A&A, 286, L35. ROSAT observations, plus optical observations.Duncan *et al.* 1997, MNRAS, 289, 97. Parkes 64-m at 1.4 GHz (18′: $S = 0.38 \pm 0.09$ Jy) 2.4 GHz (10′.6: $S = 0.25 \pm 0.04$ Jy) and 4.8 GHz (5′.7: $S = 0.17 \pm 0.02$ Jy), MOST at 843 MHz (45″×70″: $S = 0.45 \pm 0.10$ Jy), and ATCA at 2.4 GHz (37″×52″), plus ROSAT observations.Harrus *et al.* 2001, ApJ, 552, 614. ASCA and ROSAT observations, plus review of earlier observations.Kamitsukasa *et al.* 2016, PASJ, 68, S7. Suzaku observations.

G279.0+1.1**RA:** 09^h57^m40^s
Dec: –53°15′**1-GHz flux/Jy:** 30?
Spectral index: 0.6?**Size/arcmin:** 95
Type: S**Radio:** Faint, incomplete shell.**Optical:** Detected.**Point sources:** Pulsar nearby.**References:**Woermann & Jonas 1988, MNRAS, 234, 971. Hartesbeesthoek 26-m at 1.6 (30′: $S = 25.2 \pm 4$ Jy) and 2.3 GHz (20′: $S = 20.7 \pm 3$ Jy).Duncan *et al.* 1995, MNRAS, 277, 319. Parkes 64-m at 1.4 (18′: $S = 28 \pm 3$ Jy) and 2.4 GHz (11′: $S = 20 \pm 2$ Jy), including polarisation.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×53″) of part.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G284.3–1.8**RA:** 10^h18^m15^s
Dec: –59°00′**1-GHz flux/Jy:** 11?
Spectral index: 0.3?MSH 10–53
Size/arcmin: 24?
Type: S

Has been called G284.2–1.8.

Radio: Incomplete, poorly defined shell.**Optical:** Faint filament near edge.**X-ray:** Diffuse emission, brighter in N and W.**Point sources:** Central X-/ γ -ray source, not thought to be related.**References:**

Ruiz & May 1986, ApJ, 309, 667. CO and optical observations.

Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43″×50″) and Parkes 64-m at 8.4 GHz (3′: $S = 5.4 \pm 0.8$ Jy) including polarisation, plus earlier flux densities.Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.

Camilo *et al.* 2001, ApJ, 557, L51. Observations of nearby pulsar.
 Camilo *et al.* 2004, ApJ, 616, 1118. Chandra observations of pulsar and nebula.
 Abramowski *et al.* 2012, A&A, 541, A5. HESS observations.
 Williams *et al.* 2015, ApJ, 808, L19. Chandra and XMM-Newton observations.
 Marcote *et al.* 2018, A&A, 619, A76. Observations of central source.

G286.5–1.2

RA: 10^h35^m40^s
Dec: –59°42′

1-GHz flux/Jy: 1.4?
Spectral index: ?

Size/arcmin: 26×6
Type: S?

Radio: Double, elongated arc.

Optical: Detected.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×50″: $S=1.6$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G289.7–0.3

RA: 11^h01^m15^s
Dec: –60°18′

1-GHz flux/Jy: 6.2
Spectral index: 0.2?

Size/arcmin: 18×14
Type: S

Radio: Incomplete shell.

Point sources: Compact radio source near centre.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×50″: $S=6.4\pm 0.5$ Jy), plus Parkes 64-m at 4.5 GHz ($S=7.5\pm 2.5$ Jy) and 8.55 GHz ($S=3.6\pm 0.9$ Jy).

G290.1–0.8

RA: 11^h03^m05^s
Dec: –60°56′

1-GHz flux/Jy: 42
Spectral index: 0.4

Size/arcmin: 19×14
Type: S
 MSH 11–61A

Radio: Elongated, clumpy shell.

Optical: Filaments detected.

X-ray: Centrally brightened.

Point sources: Pulsar nearby, with PWN and extended ‘jet’ in X-rays.

Distance: H α absorption indicates 7 ± 1 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4′.4: $S=20.2$ Jy).
 Elliott & Malin 1979, MNRAS, 186, 45P. Optical image and spectra.
 Kirshner & Winkler 1979, ApJ, 227, 853. Optical observations.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″×50″).
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43″×49″: $S=45\pm 11$ Jy), and Parkes 64-m at 8.4 GHz (3′: $S=19.5\pm 1.0$ Jy), including polarisation.
 Seward 1990, ApJS, 73, 781. Einstein observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×49″: $S=43$ Jy).

Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Kaspi *et al.* 1997, ApJ, 485, 820. Pulsar detection.
 Gotthelf & Kaspi 1998, ApJ, 497, L29. ASCA observations of pulsar.
 Slane *et al.* 2002, ApJ, 564, 284. ASCA observations.
 Filipović *et al.* 2005, SerAJ, 170, 47. ATCA at 1.4 GHz (21''), plus other observations, including CO of surroundings.
 Reynoso *et al.* 2006, MNRAS, 369, 416. ATCA at 1.4 GHz (22'5×25'0) including H α .
 Pavan *et al.* 2014, A&A, 562, A122. Chandra and ATCA observations at 2 GHz (4'':1×5'':2) of PWN.
 Halpern *et al.* 2014, ApJ, 795, L27. XMM-Newton observations of PWN.
 Auchetti *et al.* 2015, ApJ, 810, 43. Fermi and Suzaku observations.
 Kamitsukasa *et al.* 2015, PASJ, 67, 16. Suzaku observations.
 Pavan *et al.* 2016, A&A, 591, A91. Chandra observations, including PWN and 'jet'.

G291.0–0.1

(MSH 11–62)

RA: 11^h11^m54^s**1-GHz flux/Jy:** 16**Size/arcmin:** 15×13**Dec:** –60°38'**Spectral index:** 0.29**Type:** C**Radio:** Centrally brightened core, with surrounding arcs.**Optical:** Detected.**X-ray:** Centrally brightened.**Point sources:** Central compact X-ray source.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Roger *et al.* 1986, MNRAS, 219, 815. MOST at 843 MHz (43''×50'' : S=17.2±1.0 Jy), and Parkes 64-m at 5 and 8.4 GHz (4'6 and 3' : S=10.4±0.4 Jy and 9'1±0.2), with polarisation.
 Wilson 1986, ApJ, 302, 718. Einstein observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'' : S=12.7 Jy).
 Harrus *et al.* 1998, ApJ, 499, 273. ASCA observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Slane *et al.* 2012, ApJ, 749, 131. Chandra, XMM-Newton and Fermi observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G292.0+1.8

MSH 11–54

RA: 11^h24^m36^s**1-GHz flux/Jy:** 15**Size/arcmin:** 12×8**Dec:** –59°16'**Spectral index:** 0.4**Type:** C**Radio:** Centrally brightened source surrounded by a plateau of faint emission.**Optical:** Oxygen rich.**X-ray:** Ring of emission, with diffuse central nebula and pulsar.**Point sources:** Central pulsar.**Distance:** H α absorption implies 6.0 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Lockhart *et al.* 1977, MNRAS, 179, 147. Fleurs at 1415 MHz (50'' : S=13.0 Jy).
 Goss *et al.* 1979, MNRAS, 188, 357. Optical spectra.
 Clark *et al.* 1980, MNRAS, 193, 129. X-ray spectrum.
 Tuohy *et al.* 1982, ApJ, 260, L65. Einstein observations.
 Dopita & Tuohy 1984, ApJ, 282, 135. Optical spectra.

Braun *et al.* 1986, A&A, 162, 259. MOST at 843 MHz (0'8) and IRAS.
 Hughes & Singh 1994, Apj, 422, 126. EXOSAT spectrum.
 Sutherland & Dopita 1995, Apj, 439, 365. Spectrophotometry.
 Hughes *et al.* 2001, Apj, 559, L153. Chandra observations, including central nebula.
 Park *et al.* 2002, Apj, 564, L39. Chandra observations.
 Camilo *et al.* 2002, Apj, 567, L71. Pulsar detection.
 Gonzalez & Safi-Harb 2003, Apj, 583, L91. Chandra observations.
 Hughes *et al.* 2003, Apj, 591, L139. Chandra observations of pulsar.
 Gaensler & Wallace 2003, Apj, 594, 326. ATCA at 1.4, 2.3 and 5.2 GHz (8''0×9''6, 6''2×7''2 and 4''8×5''5 : S = 11.9±0.1, 11.4±0.1 and 8.8±0.1), plus H_i observations.
 Park *et al.* 2004, Apj, 602, L33. Chandra observations.
 Vink *et al.* 2004, NuPhS, 132, 62. XMM-Newton observations.
 Ghavamian *et al.* 2005, Apj, 635, 365. Optical imaging spectroscopy.
 Winkler & Long 2006, Aj, 132, 360. Optical observations.
 Park *et al.* 2007, Apj, 670, L121. Chandra observations.
 Zhariikov *et al.* 2008, A&A, 492, 805. Possible optical counterpart to pulsar and nebula.
 Winkler *et al.* 2009, Apj, 692, 1489. Optical proper motion studies.
 Ghavamian *et al.* 2009, Apj, 696, 1307. Spitzer spectroscopy.
 Lee *et al.* 2009, Apj, 706, 441. IR observations.
 Lee *et al.* 2010, Apj, 711, 861. Chandra observations.
 Ghavamian *et al.* 2012, Apj, 750, 39. Spitzer observations.
 Zhariikov *et al.* 2013, A&A, 554, A120. IR of pulsar wind nebula.
 Yamaguchi *et al.* 2014, Apj, 785, L27. Suzaku observations.
 Kamatsukasa *et al.* 2014, PASJ, 66, 64. Suzaku observations.
 Bhalerao *et al.* 2015, Apj, 800, 65. Chandra observations.
 Koo *et al.* 2016, Apj, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Ghavamian & Williams 2016, Apj, 831, 188. Spitzer observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G292.2–0.5

RA: 11^h19^m20^s
Dec: –61°28'

1-GHz flux/Jy: 7
Spectral index: 0.5

Size/arcmin: 20×15
Type: S

Radio: Shell.

X-ray: Shell, brighter to W, with central nebula.

Point sources: Central, young pulsar.

Distance: H_i absorption indicates 8.4 kpc.

References:

Camilo *et al.* 2000, Apj, 541, 367. Pulsar detection.
 Crawford *et al.* 2001, Apj, 554, 152. ATCA at 1.4 GHz (25''×29'' : S = 5.6±0.3 Jy) and 2.5 GHz (20''×21'').
 Pivovarov *et al.* 2001, Apj, 554, 161. ROSAT and ASCA observations.
 Gonzalez & Safi-Harb 2003, Apj, 591, L143. Chandra observations of pulsar.
 Caswell *et al.* 2004, MNRAS, 352, 1405. ATCA at 5 GHz (1' : S = 2.8 Jy), including polarisation, and 1.4 GHz for H_i absorption.
 Gonzalez & Safi-Harb 2005, Apj, 619, 856. Chandra observations.
 Gonzalez *et al.* 2005, Apj, 630, 489. XMM-Newton observations of pulsar.
 Safi-Harb *et al.* 2008, Apj, 684, 532. Chandra observations of pulsar and nebula.
 Kumar *et al.* 2012, Apj, 754, 96. Chandra and XMM-Newton observations.
 Ng *et al.* 2012, Apj, 761, 65. XMM-Newton observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G293.8+0.6**RA:** 11^h35^m00^s
Dec: -60°54'**1-GHz flux/Jy:** 5?
Spectral index: 0.6?**Size/arcmin:** 20
Type: C**Radio:** Central source, with faint extended plateau.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3': $S=9.0$ Jy) and Parkes 64-m at 5 GHz (4': $S=2.1$ Jy).Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 51'').Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 49'': $S=2.6$ Jy).

G294.1-0.0**RA:** 11^h36^m10^s
Dec: -61°38'**1-GHz flux/Jy:** >2?
Spectral index: ?**Size/arcmin:** 40
Type: S**Radio:** Faint shell.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 49'': $S>2$).

G296.1-0.5**RA:** 11^h51^m10^s
Dec: -62°34'**1-GHz flux/Jy:** 8?
Spectral index: 0.6?**Size/arcmin:** 37 \times 25
Type: S

Incorporates the previously catalogued remnant G296.1-0.7. Has been called G296.05-0.50.

Radio: Irregular shell, with nearby H_{II} regions.**Optical:** Detected.**X-ray:** Irregular, incomplete shell.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3': $S=6.9$ Jy) and Parkes 64-m at 5 GHz (4': $S>0.74$).Longmore *et al.* 1977, MNRAS, 181, 541. Optical spectra.

van den Bergh 1978, ApJS, 38, 119. Optical observations.

Markert *et al.* 1981, ApJ, 248, L17. Einstein observations.Caswell & Barnes 1983, ApJ, 271, L55. Molonglo at 408 MHz (3': $S=12.4$ Jy).Bignami *et al.* 1986, ApJ, 302, 606. EXOSAT and Einstein observations.

Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'': $S>2.4$).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Castro *et al.* 2011, ApJ, 734, 86. XMM-Newton observations.

Gök & Sezer 2012, MNRAS, 419, 1603. Suzaku observations.

G296.5 + 10.0

PKS 1209–51/52

RA: 12^h09^m40^s**1-GHz flux/Jy:** 48**Size/arcmin:** 90×65**Dec:** –52°25′**Spectral index:** 0.5**Type:** S

Has been called G296.5+9.7, and erroneously G295.5+9.7.

Radio: Shell with two bright limbs.

Optical: Detected.

X-ray: Incomplete shell, with central pulsar.

Point sources: Central pulsar.

References:

- Irvine & Irvine 1974, *Apj*, 192, L111. Optical observations.
 Danziger & Dennefeld 1976, *PASP*, 88, 44. Optical spectra.
 Dickel & Milne 1976, *AujPh*, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8′.4) and 5 GHz (4′.4).
 Tuohy *et al.* 1979, *Apj*, 230, L27. X-ray detection with HEAO-1 A2 experiment.
 Ruiz 1983, *Aj*, 88, 1210. Optical spectra.
 Dubner *et al.* 1986, *Aj*, 91, 343. Argentine 30-m dish at 1.4 GHz (34′), plus H_i.
 Kellett *et al.* 1987, *MNRAS*, 225, 199. EXOSAT of the west of the remnant, including the compact source.
 Kesteven & Caswell 1987, *A&A*, 183, 118. MOST at 843 MHz (44″×56″).
 Matsui *et al.* 1988, *Apj*, 329, 838. Einstein observations, including compact source.
 Roger *et al.* 1988, *Apj*, 332, 940. MOST at 843 MHz (44″×56″).
 Bignami *et al.* 1992, *Apj*, 389, L67. Optical in vicinity of X-ray source.
 Milne & Haynes 1994, *MNRAS*, 270, 106. Parkes 64-m at 2.4 GHz (8′.3 : S = 33±3 Jy), 4.8 GHz (4′.5 : S = 23.3±3 Jy) and 8.4 GHz (3′.0 : 18.8±3), including polarisation and review of flux densities.
 Mereghetti *et al.* 1996, *Apj*, 464, 842. Radio, optical and X-ray observations of central source.
 Green *et al.* 1997, *Aj*, 114, 2058. Parkes 64-m and ATCA OH observations.
 Vasisht *et al.* 1997, *Apj*, 476, L43. ASCA observations of neutron star.
 Zavlin *et al.* 1998, *A&A*, 331, 821. ROSAT and ASCA observations of neutron star.
 Giacani *et al.* 2000, *Aj*, 119, 281. ATCA at 1.4 GHz (2′.7×4′.0) for H_i studies.
 Zavlin *et al.* 2000, *Apj*, 540, L25. Chandra observations of central pulsar.
 Gotthelf & Halpern 2007, *Apj*, 664, L35. X-ray timing observations of pulsar.
 Harvey-Smith *et al.* 2010, *Apj*, 712, 1157. ATCA at 1.4 GHz (1′.8×3′.3), including polarisation.
 Araya 2013, *MNRAS*, 434, 2202. Fermi observations.
 Halpern & Gotthelf 2015, *Apj*, 812, 61. Two epoch Chandra observations for pulsar proper motion.
 Arnaud *et al.* 2016, *A&A*, 586, A134. Planck flux densities at 4 frequencies between 30 and 100 GHz.
 Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
 Ackermann *et al.* 2018, *ApJS*, 237, 32. Fermi observations.

G296.7 – 0.9**RA:** 11^h55^m30^s**1-GHz flux/Jy:** 3**Size/arcmin:** 15×8**Dec:** –63°08′**Spectral index:** 0.5**Type:** S

Radio: Bilateral shell.

X-ray: Brighter to SE.

References:

- Schaudel *et al.* 2002, *ASPC*, 271, 391. ROSAT observations.
 Robbins *et al.* 2012, *MNRAS*, 419, 2623. ATCA at 1.4 GHz (33″×45″ : S = 2.5±0.2 Jy), plus MOST at 843 MHz (43″×49″), plus other observations.

Prinz & Becker 2013, A&A, 550, A33. XMM-Newton observations.
Green *et al.* 2014, PASA, 31, 42. Radio observations at 843 MHz (45'' \times 50'').

G296.8–0.3

1156–62

RA: 11^h58^m30^s
Dec: –62°35'

1-GHz flux/Jy: 9
Spectral index: 0.6

Size/arcmin: 20 \times 14
Type: S

Radio: Shell, brighter to the NW.

X-ray: Detected.

Distance: H α absorption gives 9.6 kpc.

References:

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3' : S=15.0 Jy) and Parkes 64-m at 5 GHz (4' : S=3.2 Jy).
Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
Hwang & Markert 1994, ApJ, 431, 819. ROSAT observations.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'' : S=9.2 Jy).
Gaensler *et al.* 1998, MNRAS, 296, 813. ATCA at 1.3 GHz (22'' \times 24'' : S = 7.0 \pm 0.3 Jy), including polarisation and H α observations, plus review of flux densities.
Sánchez-Ayaso *et al.* 2012, Ap&SS, 337, 573. XMM-Newton and IR observations.

G298.5–0.3

RA: 12^h12^m40^s
Dec: –62°52'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 5?
Type: ?

Radio: Not well resolved, may be part of a larger ring?

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'' : S=1.8 Jy).
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G298.6–0.0

RA: 12^h13^m41^s
Dec: –62°37'

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 12 \times 9
Type: S

Has been called G298.6–0.1.

Radio: Incomplete shell, in complex region.

X-ray: Centrally brightened.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 50'').
Hwang & Markert 1994, ApJ, 431, 819. ROSAT upper limit.
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'' : S=7.4 Jy).
Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

Bamba *et al.* 2016, PASJ, 68, S5. Suzaku observations.

G299.2–2.9

RA: 12^h15^m13^s
Dec: –65°30′

1-GHz flux/Jy: 0.5?
Spectral index: ?

Size/arcmin: 18×11
Type: S

Radio: Faint source.

Optical: Filaments in W.

X-ray: Centrally brightened with shell at higher energies.

References:

Busser *et al.* 1996, A&A, 310, L1. ROSAT detection, plus optical studies.
 Slane *et al.* 1996, ApJ, 465, 840. Einstein, IRAS and radio observations.
 Bai & Wang 2000, ApJ, 539, 760. ASCA observations.
 Park *et al.* 2007, ApJ, 665, 1173. Chandra observations.
 Post *et al.* 2014, ApJ, 792, L20. Chandra observations.

G299.6–0.5

RA: 12^h21^m45^s
Dec: –63°09′

1-GHz flux/Jy: 1.0?
Spectral index: ?

Size/arcmin: 13
Type: S

Radio: Faint shell, brightest to E.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×48″:S=1.1 Jy).

G301.4–1.0

RA: 12^h37^m55^s
Dec: –63°49′

1-GHz flux/Jy: 2.1?
Spectral index: ?

Size/arcmin: 37×23
Type: S

Radio: Faint, incomplete shell, with possible extension to southwest.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×48″:S=2.3 Jy).

G302.3+0.7

RA: 12^h45^m55^s
Dec: –62°08′

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 17
Type: S

Radio: Distorted shell, in complex region, with possibly associated filament.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′:S=7.5 Jy) and Parkes 64-m at 5 GHz (4′:S=3.0 Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×49″:S=3.2 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

G304.6 + 0.1

Kes 17

RA: 13^h05^m59^s**1-GHz flux/Jy:** 14**Size/arcmin:** 8**Dec:** -62°42'**Spectral index:** 0.5**Type:** S**Radio:** Incomplete shell.**X-ray:** Detected.**Distance:** Possible limit of > 9.7 kpc from H_I absorption.**References:**

Shaver & Goss 1970, *AJPA*, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Milne & Dickel 1975, *AJPh*, 28, 209. Parkes 64-m at 5 GHz (4'.4 : S=6.9 Jy).
 Caswell *et al.* 1975, *A&A*, 45, 239. Parkes H_I absorption.
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×48'' : S=18 Jy).
 Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2009, *ApJ*, 694, 1266. Spitzer spectroscopy.
 Combi *et al.* 2010, *A&A*, 523, A76. XMM-Newton observations.
 Lee *et al.* 2011, *ApJ*, 740, 31. Akari and Spitzer observations.
 Wu *et al.* 2011, *ApJ*, 740, L12. Fermi observations.
 Andersen *et al.* 2011, *ApJ*, 742, 7. Spitzer observations.
 Gök & Sezer 2012, *MNRAS*, 423, 1215. Suzaku observations.
 Gelfand *et al.* 2013, *ApJ*, 777, 148. ATCA at 1.4 GHz (8''×23'' : S=10.9±0.4 Jy), plus other observations.
 Pannuti *et al.* 2014, *AJ*, 147, 55. ASCA and XMM-Newton observations.
 Koo *et al.* 2016, *ApJ*, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.
 Washino *et al.* 2016, *PASJ*, 68, S4. Suzaku observations.

G306.3 – 0.9**RA:** 13^h21^m50^s**1-GHz flux/Jy:** 0.16?**Size/arcmin:** 4**Dec:** -63°34'**Spectral index:** 0.5?**Type:** S?**Radio:** Diffuse emission.**X-ray:** Partial shell.**References:**

Reynolds *et al.* 2013, *ApJ*, 766, 112. Chandra observations, and ATCA at 5.5 GHz (23''×26''), and 9 GHz.
 Combi *et al.* 2016, *A&A*, 592, A125. XMM-Newton and Chandra observations.
 Sezar *et al.* 2017, *MNRAS*, 466, 3434. Suzaku and Fermi observations.

G308.1 – 0.7**RA:** 13^h37^m37^s**1-GHz flux/Jy:** 1.2?**Size/arcmin:** 13**Dec:** -63°04'**Spectral index:** ?**Type:** S**Radio:** Faint shell.**References:**

Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43''×48'' : S=1.3 Jy).

G308.4–1.4**RA:** 13^h41^m30^s
Dec: –63°44′**1-GHz flux/Jy:** 0.4?
Spectral index: ?**Size/arcmin:** 12×6?
Type: S?

W part has been called G308.3–1.4.

Radio: Complex structure, with multiple arcs.**X-ray:** Limb brightened partial shell in W.**References:**Schaudel *et al.* 2002, ASPC, 271, 391. ROSAT observations.Prinz & Becker 2012, A&A, 544, A7. ATCA at 1.4 GHz (53″×64″: $S=0.33$ Jy) and 2.5 GHz (29″×35″: $S=0.24$ Jy), plus Chandra and other observations.Hui *et al.* 2012, ApJ, 750, 7. XMM-Newton and other observations.De Horta *et al.* 2013, MNRAS, 428, 1980. ATCA at 1.4 GHz (54″×65″) and 2.5 GHz (29″×35″), plus other observations.Green *et al.* 2014, PASA, 31, 42. Radio observations at 843 MHz (45″×50″).

G308.8–0.1**RA:** 13^h42^m30^s
Dec: –62°23′**1-GHz flux/Jy:** 15?
Spectral index: 0.4?**Size/arcmin:** 30×20?
Type: C?

Incorporates previous catalogued remnant G308.7+0.0.

Radio: Bright ridge in north, and arc to south.**Point sources:** Pulsar near centre of remnant.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=16.7$ Jy) and Parkes 64-m at 5 GHz (4′: $S=7.0$ Jy).Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4′.4: $S=6.5$ Jy).Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50″).

Wilson 1986, ApJ, 302, 718. Einstein upper limit.

Caswell *et al.* 1992, ApJ, 399, L151. MOST at 843 MHz (43″×49″).Kaspi *et al.* 1992, ApJ, 399, L155. Pulsar observations.Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G309.2–0.6**RA:** 13^h46^m31^s
Dec: –62°54′**1-GHz flux/Jy:** 7?
Spectral index: 0.4?**Size/arcmin:** 15×12
Type: S

Has been called G309.2–0.7.

Radio: Distorted shell.**X-ray:** Extended emission, with unrelated central source.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=10.0$ Jy) and Parkes 64-m at 5 GHz (4′: $S=3.9$ Jy).Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (1′.5).

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 50'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'': $S=6$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Gaensler *et al.* 1998, MNRAS, 299, 812. ATCA at 1.3 GHz (23'' \times 24'': $S=5.2\pm 0.2$ Jy).
 Rakowski *et al.* 2001, ApJ, 548, 258. ASCA and ROSAT observations.
 Rakowski *et al.* 2006, ApJ, 649, L111. Observations of central source.
 Safi-Harb *et al.* 2007, ApJ, 659, 407. Observations of central source.

G309.8 + 0.0

RA: 13^h50^m30^s
Dec: -62°05'

1-GHz flux/Jy: 17
Spectral index: 0.5

Size/arcmin: 25 \times 19
Type: S

Radio: Distorted shell.

Point sources: Steep radio spectrum source near the centre of the remnant.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : $S=26.4$ Jy) and Parkes 64-m at 5 GHz (4' : $S=7.4$ Jy).
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (1').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 49'': $S>8.8$).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G310.6 - 1.6

RA: 14^h00^m45^s
Dec: -63°26'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 2.5
Type: C?

Radio: Not detected.

X-ray: Bright central nebula, with faint shell.

Point sources: X-ray pulsar, and extended emission.

References:

Tomsick *et al.* 2009, ApJ, 701, 811. Chandra observations.
 Renaud *et al.* 2010, ApJ, 716, 663. Chandra and other observations.

G310.6 - 0.3

RA: 13^h58^m00^s
Dec: -62°09'

1-GHz flux/Jy: 5?
Spectral index: ?

Kes 20B
Size/arcmin: 8
Type: S

Has been called G310.65-0.29.

Radio: Asymmetric shell.

References:

Whiteoak *et al.* 1994, MNRAS, 269, 294. MOST at 843 MHz.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 49'': $S=5.4$ Jy).
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G310.8–0.4

Kes 20A

RA: 14^h00^m00^s**Dec:** –62°17′**1-GHz flux/Jy:** 6?**Spectral index:** ?**Size/arcmin:** 12**Type:** S

Has been called G310.80–0.41.

Radio: Arc in E, in complex region.

References:

Whiteoak *et al.* 1994, MNRAS, 269, 294. MOST at 843 MHz.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'' : S=6.9 Jy).

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.

G311.5–0.3**RA:** 14^h05^m38^s**Dec:** –61°58′**1-GHz flux/Jy:** 3?**Spectral index:** 0.5**Size/arcmin:** 5**Type:** S

Has been called G311.52–0.37.

Radio: Shell, not well resolved.

X-ray: Detected.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3').

Whiteoak *et al.* 1994, MNRAS, 269, 294. MOST at 843 MHz.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'' : S=2.9 Jy).

Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.

Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.

Pannuti *et al.* 2014, AJ, 147, 55. ASCA observations.

G312.4–0.4**RA:** 14^h13^m00^s**Dec:** –61°44′**1-GHz flux/Jy:** 45**Spectral index:** 0.36**Size/arcmin:** 38**Type:** S

Has been called G312.44–0.36.

Radio: Irregular, incomplete shell.

X-ray: Weak emission in W.

Point sources: Nearby γ -ray sources and pulsars.

Distance: H α absorption suggests > 6 kpc and possibly > 14 kpc.

References:

Caswell & Barnes 1985, MNRAS, 216, 753. Molonglo at 408 MHz (3' : S=56 Jy).

Whiteoak *et al.* 1994, MNRAS, 269, 294. MOST at 843 MHz.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×49'' : S > 19), plus Parkes 64-m at 4.5 GHz (S=30±2 Jy) and 8.55 GHz (S=17±4 Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Roberts *et al.* 1999, ApJ, 515, 712. MOST at 843 MHz (43''×49'').

Case & Bhattacharya 1999, ApJ, 521, 246. Nearby γ -ray sources.

Doherty *et al.* 2003, MNRAS, 339, 1048. ATCA at 1.4 GHz (25'') plus H α absorption, and Chandra observations.

G312.5–3.0

RA: 14^h21^m00^s
Dec: –64°12'

1-GHz flux/Jy: 3.5?
Spectral index: ?

Size/arcmin: 20×18
Type: S

Radio: Distorted shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10'.4)

Kane & Vaughan 2003, MNRAS, 344, 625. ATCA at 1.4 GHz (116''×129'') and 2.4 GHz (67''×75'').

G315.1+2.7

RA: 14^h24^m30^s
Dec: –57°50'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 190×150
Type: S

Radio: Poorly defined shell.

Optical: Filaments, brighter in NE.

References:

Duncan *et al.* 1997, MNRAS, 287, 722. Parkes 64-m at 2.4 GHz (11':S=19±3 Jy).

Combi *et al.* 1998, A&A, 333, 298. Radio survey observations.

Stupar *et al.* 2007, MNRAS, 374, 1441. Optical and various radio observations.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G315.4–2.3

RA: 14^h43^m00^s
Dec: –62°30'

1-GHz flux/Jy: 49
Spectral index: 0.6

RCW 86, MSH 14–63

Size/arcmin: 42
Type: S

Possibly the remnant of the SN of AD185?

Radio: Shell, brightest to the SW.

Optical: Bright, radiative filaments, with some faint Balmer dominated filaments.

X-ray: Partial shell, with thermal and non-thermal emission.

Point sources: Several X-ray sources.

Distance: Optical observations imply 2.3 kpc.

References:

van den Bergh *et al.* 1973, ApJS, 26, 19. Optical observations.

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3':S=86 Jy) and Parkes 64-m at 5 GHz (4':S=18.2 Jy).

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).

Winkler 1978, ApJ, 221, 220. X-ray detection.

Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.

Pisarski *et al.* 1984, ApJ, 277, 710. Einstein observations

Nugent *et al.* 1984, ApJ, 284, 612. X-ray spectrum.

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44'' \times 50'').
 Long & Blair 1990, ApJ, 358, L13. Balmer dominated optical filaments.
 Greidanus & Strom 1990, A&A, 240, 385. IRAS observations.
 Kaastra *et al.* 1992, A&A, 264, 654. Ginga X-ray spectra.
 Strom 1994, MNRAS, 268, L5. Historical association
 Chin & Huang 1994, Nature, 371, 398. Questioning of historical association.
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 48'': $S > 22$).
 Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Smith *et al.* 1997, AJ, 114, 2664. Observations of Balmer dominated filaments.
 Vink *et al.* 1997, A&A, 328, 628. ASCA spectroscopy.
 Bocchino *et al.* 2000, A&A, 360, 671. BeppoSAX observations of N and SW.
 Bamba *et al.* 2000, PASJ, 52, 1157. ASCA observations.
 Dickel *et al.* 2001, ApJ, 546, 447. ATCA at 1.34 GHz (8''), including polarisation.
 Ghavamian *et al.* 2001, ApJ, 547, 995. Optical spectroscopy.
 Borkowski *et al.* 2001, ApJ, 550, 334. ASCA observations.
 Rho *et al.* 2002, ApJ, 581, 1116. Chandra observations.
 Gvaramadze & Vikhlinin 2003, A&A, 401, 625. Chandra point source search.
 Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
 Kaplan *et al.* 2004, ApJS, 153, 269. Chandra limits for any compact source.
 Bamba *et al.* 2005, ApJ, 621, 793. Chandra observations of rim.
 Vink *et al.* 2006, ApJ, 648, L33. Chandra and XMM-Newton observations.
 Ueno *et al.* 2007, PASJ, 59, S171. Suzaku observations.
 Aharonian *et al.* 2009, ApJ, 692, 1500. HESS observations.
 Yamaguchi *et al.* 2011, PASJ, 63, S837. Suzaku observations.
 Helder *et al.* 2011, ApJ, 737, 85. XMM-Newton and optical observations.
 Williams *et al.* 2011, ApJ, 741, 96. Spitzer observations.
 Mignani *et al.* 2012, MNRAS, 425, 2309. X-ray and optical observations of compact X-ray sources.
 Lemoine-Goumard *et al.* 2012, A&A, 545, A28. Fermi observations.
 Helder *et al.* 2013, MNRAS, 435, 910. Optical proper motion studies.
 Castro *et al.* 2013, ApJ, 779, 49. Chandra of NW.
 Yuan *et al.* 2014, ApJ, 785, L22. Fermi observations.
 Tsubone *et al.* 2014, in EFXU, p72. Suzaku observations.
 Broersen *et al.* 2014, MNRAS, 441, 3040. XMM-Newton observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 30, 44 and 70 GHz.
 Ajello *et al.* 2016, ApJ, 819, 98. Fermi observations, and ATCA H α of surroundings.
 Yamaguchi *et al.* 2016, ApJ, 820, L3. Chandra observations of NE.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Tsubone *et al.* 2017, ApJ, 835, 34. Suzaku observations.
 Sano *et al.* 2017, JHEAp, 15, 1. CO and H α observations.
 Abramowski *et al.* 2018, A&A, 612, A4. HESS observations.

G315.4–0.3

RA: 14^h35^m55^s

Dec: –60°36'

1-GHz flux/Jy: 8

Spectral index: 0.4

Size/arcmin: 24 \times 13

Type: ?

Radio: Irregular non-thermal emission, with H α region superposed in E.

Optical: Detected.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : $S = 15.9$ Jy) and Parkes 64-m at 5 GHz (4' : $S = 4.9$ Jy).

Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz ($50'' : S = 6.25$ Jy), re-assessment of earlier flux densities.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 49'' : S = 3.1$ Jy).

Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G315.9–0.0

RA: $14^{\text{h}}38^{\text{m}}25^{\text{s}}$
Dec: $-60^{\circ}11'$

1-GHz flux/Jy: 0.8?
Spectral index: ?

Size/arcmin: 25×14
Type: S

Has been called G315.8–0.0.

Radio: Faint, distorted shell, with elongated trail to pulsar.

Point sources: Pulsar at end of radio trail.

References:

Kesteven *et al.* 1987, AujPh, 40, 855. MOST at 843 MHz ($44'' \times 50''$).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 0.9$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.

Camilo *et al.* 2009, ApJ, 703, L55. Pulsar detection.

Ng *et al.* 2012, ApJ, 746, 105. ATCA at 1.4, 2.5, 5 and 9 GHz, including polarisation, of pulsar trail.

G316.3–0.0

(MSH 14–57)

RA: $14^{\text{h}}41^{\text{m}}30^{\text{s}}$
Dec: $-60^{\circ}00'$

1-GHz flux/Jy: 20?
Spectral index: 0.4

Size/arcmin: 29×14
Type: S

Radio: Distorted shell, with possible 'blowout'.

X-ray: Detected.

Distance: H α absorption data suggests > 7.2 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz ($4'.4 : S = 16.7$ Jy).

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 51''$).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 20$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Acerio *et al.* 2016, ApJS, 224, 8. Fermi observations.

G317.3–0.2

RA: $14^{\text{h}}49^{\text{m}}40^{\text{s}}$
Dec: $-59^{\circ}46'$

1-GHz flux/Jy: 4.7?
Spectral index: ?

Size/arcmin: 11
Type: S

Radio: Incomplete shell.

Optical: Detected.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 50'' : S = 5.2$ Jy).

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G318.2 + 0.1

RA: 14^h54^m50^s
Dec: -59°04'

1-GHz flux/Jy: >3.9?
Spectral index: ?

Size/arcmin: 40×35
Type: S

Radio: Faint shell, with central H II region.

X-ray: Sources within remnant.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'' : S > 4.3).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Bocchino *et al.* 2001, A&A, 367, 629. BeppoSAX and ROSAT observations.

Onako *et al.* 2016, ApJ, 829, 106. Akari observations.

G318.9 + 0.4

RA: 14^h58^m30^s
Dec: -58°29'

1-GHz flux/Jy: 4?
Spectral index: 0.2?

Size/arcmin: 30×14
Type: C

May not be a SNR?

Radio: Complex arcs, with off-centre core.

References:

Whiteoak 1990, Nature, 347, 157. MOST at 843 MHz (43''×51'').

Whiteoak 1993, ApJ, 415, 701. MOST at 843 MHz (43''×51'' : S = 4.8±0.6 Jy), Parkes 64-m at 4.5 GHz (4'.7 : S = 3.7±0.2 Jy) and 8.4 GHz (2'.8 : S = 3.0±0.4 Jy) including polarisation, and ATCA at 1.4 GHz and 4.8 GHz (11''×13'') of core.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'' : S = 4.8 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G320.4 – 1.2

RA: 15^h14^m30^s
Dec: -59°08'

1-GHz flux/Jy: 60?
Spectral index: 0.4

MSH 15–52, RCW 89

Size/arcmin: 35
Type: C

Has been suggested as the remnant of the SN of AD185?

Radio: Ragged shell.

Optical: RCW 89 is the H α emitting region to the NW.

X-ray: Partial shell, central nebula and pulsar and 'jet'.

Point sources: Radio and X-ray pulsar, with wind nebula.

Distance: H I absorption indicates 5.2 kpc.

References:

Caswell *et al.* 1975, A&A, 45, 239. Parkes H I absorption.

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).

Dopita *et al.* 1977, ApJ, 214, 179. Optical spectra.

Caswell *et al.* 1981, MNRAS, 195, 89. FIRST at 1415 MHz (50'').

Seward & Harnden 1982, ApJ, 256, L45. X-ray observations of pulsar.

Manchester *et al.* 1982, ApJ, 262, L31. Radio observations of pulsar.
 Seward *et al.* 1983, ApJ, 267, 698. X-ray, Optical and IR.
 van den Bergh & Kamper 1984, ApJ, 280, L51. Optical expansion.
 Seward *et al.* 1984, ApJ, 281, 650. X-ray observations of pulsar and nebulosity.
 Lortet *et al.* 1987, A&A, 180, 65. Optical observations.
 Trussoni *et al.* 1990, A&A, 234, 403. EXOSAT observations.
 Asaoka & Koyama 1990, PASJ, 42, 625. Ginga X-ray spectrum.
 Arendt 1991, AJ, 101, 2160. IRAS observations, including compact source.
 Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz (4'5 : S = 37±7 Jy) and 8.4 GHz (3'0 : S = 24±4 Jy), including polarisation and review of flux densities.
 Strom 1994, MNRAS, 268, L5. Historical association.
 Chin & Huang 1994, Nature, 371, 398. Questioning of historical association.
 Matz *et al.* 1994, ApJ, 434, 288. X-ray observations of pulsar.
 Schaefer 1995, AJ, 110, 1793. Questioning of historical association.
 Du Plessis *et al.* 1995, ApJ, 453, 746. Hartesbeesthoek 26-m at 2.3, 5 and 8.5 GHz (S = 42, 35.6 and 14.5 Jy).
 Greiveldinger *et al.* 1995, ApJ, 454, 855. ROSAT observations.
 Trussoni *et al.* 1996, A&A, 306, 581. ROSAT observations.
 Tamura *et al.* 1996, PASJ, 48, L33. ASCA observations.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'' : S = 62 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Marsden *et al.* 1997, ApJ, 491, L39. X-ray spectroscopy.
 Brazier & Becker 1997, MNRAS, 284, 335. ROSAT observations.
 Gaensler *et al.* 1999, MNRAS, 305, 724. ATCA at 1.4 GHz (21''×24''), plus H_i observations, and 5.3 GHz (10''×15'').
 Sako *et al.* 2000, ApJ, 537, 422. Possible γ -ray detection of pulsar.
 Mineo *et al.* 2001, A&A, 380, 695. BeppoSAX observations.
 Gaensler *et al.* 2002, ApJ, 569, 878. Chandra observations of pulsar and nebula.
 Dubner *et al.* 2002, AJ, 123, 337. ATCA at 1.4 GHz (2'7×4'0), plus H_i observations.
 Aharonian *et al.* 2005, A&A, 435, L17. HESS detection.
 Yatsu *et al.* 2005, ApJ, 631, 312. Chandra observations of pulsar and jet.
 DeLaney *et al.* 2006, ApJ, 640, 929. Chandra and ROSAT multi-epoch observations of pulsar wind nebula.
 Kaplan & Moon 2006, ApJ, 644, 1056. Possible IR detection of pulsar.
 Forot *et al.* 2006, ApJ, 651, L45. X-ray observations of pulsar and nebula.
 Nakamori *et al.* 2008, ApJ, 677, 297. γ -ray observations.
 Yatsu *et al.* 2009, PASJ, 61, 129. Chandra observations of pulsar wind nebula.
 Koo *et al.* 2011, ApJ, 732, 6. Akari and Spitzer observations.
 An *et al.* 2014, ApJ, 793, 90. NuSTAR observations of PWN.
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.

G320.6–1.6

RA: 15^h17^m50^s
Dec: –59°16'

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 60×30
Type: S

Radio: Faint shell, overlapping G320.4–1.2 in W.

Optical: Detected.

References:

Milne *et al.* 1993, MNRAS, 264, 853. Parkes 64-m at 4.8 GHz (4'5) and 8.4 GHz (3'0), including polarisation.
 Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz (10'4).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×50'' : S > 9.3).
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G321.9–1.1**RA:** 15^h23^m45^s
Dec: –58°13′**1-GHz flux/Jy:** >3.4?
Spectral index: ?**Size/arcmin:** 28
Type: S**Radio:** Faint shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×51'' : S > 3.8).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G321.9–0.3**RA:** 15^h20^m40^s
Dec: –57°34′**1-GHz flux/Jy:** 13
Spectral index: 0.3**Size/arcmin:** 31×23
Type: S**Radio:** Shell brighter to the W, with Cir X-1 to N.**Point sources:** Pulsar near E edge.**References:**

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : S = 18.3 Jy) and Parkes 64-m at 5 GHz (4' : S = 7.8 Jy).
Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×52'').
Stewart *et al.* 1993, MNRAS, 261, 593. ATCA at 1.5 GHz (21'').
Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×51'' : S > 8.3).
Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
Mignani *et al.* 2002, A&A, 386, 487. HST proper motion study of Cir X-1.
Tudose *et al.* 2006, MNRAS, 372, 417. MOST at 843 MHz.
Pletsch *et al.* 2013, ApJ, 779, L11. Pulsar detection.
Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G322.1+0.0**RA:** 15^h20^m49^s
Dec: –57°10′**1-GHz flux/Jy:** ?
Spectral index: ?**Size/arcmin:** 8×4.5?
Type: S?**Radio:** Circular shell, with extension to S.**X-ray:** Diffuse emission.**Point sources:** Cir X-1 HMXB at centre.**References:**

Heinz *et al.* 2013, ApJ, 779, 171. Chandra observations and ATCA at 1.1--3.1 GHz (4''0×4''9).

G322.5–0.1

RA: 15^h23^m23^s
Dec: –57°06′

1-GHz flux/Jy: 1.5
Spectral index: 0.4

Size/arcmin: 15
Type: C

Radio: Shell with central extended source.

Point sources: PN Pe 2-8 within boundary.

References:

Whiteoak 1992, MNRAS, 256, 121. MOST at 843 MHz (43″×51″: $S=2.0\pm0.3$ Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×51″: $S=1.7$ Jy), plus Parkes 64-m at 4.5 GHz ($=0.89\pm0.13$).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G323.5+0.1

RA: 15^h28^m42^s
Dec: –56°21′

1-GHz flux/Jy: 3?
Spectral index: 0.4?

Size/arcmin: 13
Type: S

Radio: Distorted shell, confused with thermal emission.

Point sources: Compact, probably thermal source near centre.

References:

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3′: $S=4.2$ Jy) and Parkes 64-m at 5 GHz (4′: $S=1.5$ Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×52″: $S=4.2$ Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G323.7–1.0

RA: 15^h34^m30^s
Dec: –57°12′

1-GHz flux/Jy: ?
Spectral index: ?

Size/arcmin: 51×38
Type: S

Radio: Faint shell.

X-ray: Faint diffuse emission.

Distance: CO and H_I observations suggest 3.5 kpc.

References:

Green *et al.* 2014, PASA, 31, 42. MGPS observations at 843 MHz (43″×51″).

Araya *et al.* 2017, ApJ, 843, 12. Fermi observations.

Abdalla *et al.* 2018, A&A, 612, A8. HESS observations.

Maxted *et al.* 2018, MNRAS, 480, 134. CO and H_I observations.

Saji *et al.* 2018, PASJ, 70, 23. Suzaku observations.

G326.3–1.8

MSH 15–56

RA: 15^h53^m00^s**1-GHz flux/Jy:** 145**Size/arcmin:** 38**Dec:** –56°10′**Spectral index:** varies**Type:** C

Has been called G326.2–1.7.

Radio: Shell, with elongated, flat-spectrum core.

Optical: Emission around the shell.

X-ray: Shell, with central extended emission.

Point sources: Compact X-ray source.

References:

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′: $S=180$ Jy).

Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.

Milne *et al.* 1979, MNRAS, 188, 437. FIRST at 1415 MHz (0′.8: $S>95$) and Parkes 64-m at 14.7 GHz (2′.2: $S=69\pm 8$ Jy).

van den Bergh 1979, ApJ, 227, 497. Optical observations.

Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.

Dennefeld 1980, PASP, 92, 603. Optical spectra.

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″ \times 53″).

Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43″ \times 52″: $S=153\pm 40$ Jy), and Parkes 64-m at 8.4 GHz (3′: $S=68\pm 5$ Jy), including polarisation.

Seward 1990, ApJS, 73, 781. Einstein observations.

Kassim *et al.* 1993, ApJ, 419, 733. ROSAT image.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ \times 52″: $S>130$).

Rosado *et al.* 1996, A&A, 315, 243. Optical kinematics.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Dickel *et al.* 2000, ApJ, 543, 840. ATCA at 1.34 GHz (6′.4 \times 8′.5: $S>60$ Jy), 4.80 GHz (3′.8: $S=25$ Jy for core only) and 8.64 GHz (3′.2: $S=15$ Jy for core only).

Temim *et al.* 2013, ApJ, 768, 61. XMM-Newton and Chandra observations.

Yatsu *et al.* 2013, ApJ, 773, 25. XMM-Newton and Chandra observations.

Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 5 frequencies between 30 and 143 GHz.

Acerro *et al.* 2016, ApJS, 224, 8. Fermi observations.

Temim *et al.* 2017, ApJ, 851, 128. Chandra proper motion study of central source.

Devin *et al.* 2018, A&A, 617, A5. Fermi observations.

G327.1–1.1**RA:** 15^h54^m25^s**1-GHz flux/Jy:** 7?**Size/arcmin:** 18**Dec:** –55°09′**Spectral index:** ?**Type:** C

Radio: Shell, with off-centre core.

X-ray: Diffuse, with core.

References:

Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (8′.4: $S=10\pm 15\%$ Jy).

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=10.6$ Jy) and Parkes 64-m at 5 GHz (4′: $S=4.3$ Jy).

Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ \times 52″: $S=7.6$ Jy).

Seward *et al.* 1996, ApJ, 471, 887. ROSAT observations.

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

Sun *et al.* 1999, ApJ, 511, 274. ASCA and ROSAT observations.

Bocchino & Bandiera 2003, A&A, 398, 195. BeppoSAX observations.
 Temim *et al.* 2009, ApJ, 691, 895. Chandra and XMM-Newton observations.
 Temim *et al.* 2015, ApJ, 808, 100. Chandra observations.
 Ma *et al.* 2016, ApJ, 820, 100. ATCA at 1.4 (19''×23''), 2.3 (11''×16''), 4.8 (13''×15'') and 8.6 GHz (8''×10'') of core, including polarisation.
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.

G327.2–0.1

RA: 15^h50^m55^s
Dec: –54°18'

1-GHz flux/Jy: 0.4
Spectral index: ?

Size/arcmin: 5
Type: S

Has been called G327.24–0.13.

Radio: Shell, possibly with central emission.

Point sources: Central pulsar (magnetar).

References:

Camilo *et al.* 2007, ApJ, 666, L93. Pulsar observations.
 Gelfand & Gaensler 2007, ApJ, 667, 1111. Chandra and XMM-Newton observations of magnetar, and 843 GHz (43''×53'': $S=0.5\pm0.1$ Jy) and SGPS at 1.4 GHz ($S=0.3\pm0.1$ Jy).

G327.4+0.4

RA: 15^h48^m20^s
Dec: –53°49'

1-GHz flux/Jy: 30?
Spectral index: 0.6

Size/arcmin: 21
Type: S

Kes 27

Has been called G327.3+0.4 and G327.3+0.5.

Radio: Incomplete, multi-arc shell, brightest to the SE.

X-ray: Diffuse, brighter in E.

Distance: H_I absorption indicates 4.3 to 5.4 kpc.

References:

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3': $S=58$ Jy) and Parkes 64-m at 5 GHz (4': $S=12.4$ Jy).
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'.4) and 5 GHz (4'.4).
 Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×55'').
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz (43''×53'': $S=32.2\pm6$ Jy), and Parkes 64-m at 8.4 GHz (3': $S=9.4\pm0.8$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×53'': $S=25$ Jy).
 Seward *et al.* 1996, ApJ, 471, 887. ROSAT observations.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m and ATCA OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1'.8×2'.0), plus H_I.
 Enoguchi *et al.* 2002, PASJ, 54, 229. ASCA observations.
 Kawasaki *et al.* 2005, ApJ, 631, 935. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Chen *et al.* 2008, ApJ, 676, 1040. Chandra observations.
 Minami *et al.* 2014, in EFXU, p48. Suzaku observations.
 Xing *et al.* 2015, ApJ, 805, 19. Fermi observations.

G327.4 + 1.0**RA:** 15^h46^m48^s**Dec:** -53°20′**1-GHz flux/Jy:** 1.9?**Spectral index:** ?**Size/arcmin:** 14**Type:** S**Radio:** Asymmetric shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×54'' : S=2.1 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1'8×2'0), plus H_r.**G327.6 + 14.6****RA:** 15^h02^m50^s**Dec:** -41°56′**1-GHz flux/Jy:** 19**Spectral index:** 0.6

SN1006, PKS 1459-41

Size/arcmin: 30**Type:** S

This is the remnant of the SN of AD1006.

Radio: Shell, with two bright arcs.**Optical:** Filaments to the NW, with broad H α component.**X-ray:** Thermal shell, with non-thermal limb-brightened arcs.**Point sources:** The background Schweizer--Middleditch star is near the middle of the remnant.**Distance:** Optical spectra and proper motion indicate 1.85 kpc.**References:**

van den Bergh 1976, ApJ, 208, L17. Optical observations.

Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz (8'4 and 5 GHz (4'4).

Schweizer & Middleditch 1980, ApJ, 241, 1039. Possible stellar remnant.

Caswell *et al.* 1983, MNRAS, 204, 921. FIRST at 1415 MHz (77'').

Reynolds & Gilmore 1986, AJ, 92, 1138. VLA at 1.37 and 1.67 GHz (16''×20'').

Kirshner *et al.* 1987, ApJ, 315, L135. Broad H α optical component.

Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44''×66'').

Fesen *et al.* 1988, ApJ, 327, 164. UV absorption spectra of the Schweizer--Middleditch star.Roger *et al.* 1988, ApJ, 332, 940. MOST at 843 MHz (44''×66'' : S=17.5±1.5 Jy).Long *et al.* 1988, ApJ, 333, 749. Optical proper motion for distance.Wu *et al.* 1993, ApJ, 416, 247. UV spectra.

Reynolds & Gilmore 1993, AJ, 106, 272. VLA at 1.37 and 1.67 GHz (24''), including polarisation.

Moffett *et al.* 1993, AJ, 106, 1566. VLA at 1.37 and 1.67 GHz (10''×14''), for proper motion studies.Ozaki *et al.* 1994, PASJ, 46, 367. X-ray observations.Raymond *et al.* 1995, ApJ, 454, L31. Far UV spectroscopy.Koyama *et al.* 1995, Nature, 378, 255. ASCA observations.Willingale *et al.* 1996, MNRAS, 278, 749. ROSAT observations.Blair *et al.* 1996, ApJ, 468, 871. Optical absorption studies.Laming *et al.* 1996, ApJ, 472, 267. Modelling of optical spectra, including distance.Wu *et al.* 1997, ApJ, 477, L53. Far UV absorption spectra of the Schweizer--Middleditch star.

Winkler & Long 1997, ApJ, 486, L137. UV absorption spectra of background quasar.

Winkler & Long 1997, ApJ, 491, 829. ROSAT and optical images.

Vink *et al.* 2000, A&A, 354, 931. X-ray spectroscopy.Burleigh *et al.* 2000, A&A, 356, 585. Optical spectroscopy of the Schweizer--Middleditch star.Allen *et al.* 2001, ApJ, 558, 739. ASCA, ROSAT and other X-ray observations.Dubner *et al.* 2002, A&A, 387, 1047. ATCA at 1.4 GHz (3'0×4'7) for H_r, plus CO observations.

Ghavamian *et al.* 2002, ApJ, 572, 888. Optical of filaments in NW.
 Sollerman *et al.* 2003, A&A, 407, 249. Optical spectroscopy.
 Winkler *et al.* 2003, ApJ, 585, 324. Optical proper motion studies.
 Vink *et al.* 2003, ApJ, 587, L31. XMM-Newton observations.
 Bamba *et al.* 2003, ApJ, 589, 827. Chandra observations of NE.
 Winkler *et al.* 2005, ApJ, 624, 189. HST absorption towards background sources.
 Kalemci *et al.* 2006, ApJ, 644, 274. INTEGRAL observations.
 Acero *et al.* 2007, A&A, 475, 883. XMM-Newton observations.
 Raymond *et al.* 2007, ApJ, 659, 1257. HST H α observations.
 Hamilton *et al.* 2007, MNRAS, 381, 771. UV spectroscopy of the Schweizer--Middleditch star.
 Bamba *et al.* 2008, AdSpR, 41, 411. Suzaku observations.
 Cassam-Chenaï *et al.* 2008, ApJ, 680, 1180. ATCA and VLA at 1.5 GHz (6'' \times 9'') plus Chandra observations.
 Yamaguchi *et al.* 2008, PASJ, 60, S141. Suzaku observations.
 Dyer *et al.* 2009, AJ, 137, 2956. GBT and VLA observations at 1.4 GHz.
 Winkler *et al.* 2011, ApJ, 742, 80. Multi-epoch UV spectroscopy of the Schweizer--Middleditch star.
 Miceli *et al.* 2013, A&A, 550, A80. XMM-Newton observations.
 Broersen *et al.* 2013, A&A, 552, A9. XMM-Newton observations of NW knot.
 Katsuda *et al.* 2013, ApJ, 763, 85. Chandra proper motion study of NW.
 Winkler *et al.* 2013, ApJ, 764, 156. Spitzer observations.
 Uchida *et al.* 2013, ApJ, 771, 56. Suzaku observations.
 Reynoso *et al.* 2013, AJ, 145, 104. VLA and ATCA at 1.4 GHz (10''), including polarisation.
 Nikolić *et al.* 2013, Science, 340, 45. Optical spectroscopy.
 Winkler *et al.* 2014, ApJ, 781, 65. Chandra observations and H α spectroscopy.
 Miceli *et al.* 2014, ApJ, 782, L33. XMM-Newton observations of SW.
 Acero *et al.* 2015, A&A, 580, A74. Fermi limit.
 Sparks *et al.* 2015, ApJ, 815, L9. Polarised optical line observations.
 Li *et al.* 2015, MNRAS, 453, 3953. XMM-Newton observations.
 Arnaud *et al.* 2016, A&A, 586, A134. Planck flux densities at 4 frequencies between 30 and 100 GHz.
 Xing *et al.* 2016, ApJ, 823, 44. Fermi observations.
 Li *et al.* 2016, MNRAS, 462, 158. XMM-Newton observations.
 Raymond *et al.* 2017, ApJ, 851, 12. UV and optical spectra, for proper motions and distance.
 Condon *et al.* 2017, ApJ, 851, 100. Fermi observations.
 Kerzendorf *et al.* 2018, MNRAS, 479, 192. Optical search for progenitor companion.
 Li *et al.* 2018, ApJ, 864, 85. NuSTAR observations of NE and SW limbs.

G328.4 + 0.2

(MSH 15–57)

RA: 15^h55^m30^s**1-GHz flux/Jy:** 15**Size/arcmin:** 5**Dec:** –53°17'**Spectral index:** 0.0**Type:** F**Radio:** Amorphous emission, with central bar.**X-ray:** Detected at high energies.**Distance:** H α absorption indicates > 17.4 kpc.**References:**

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50'').
 Wilson 1986, ApJ, 302, 718. Lack of detection with Einstein.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43'' \times 54'' : $S = 15$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Gaensler *et al.* 2000, ApJ, 542, 380. ATCA at 1.4 GHz (16''5 \times 19''5 : $S = 14.3 \pm 0.1$ Jy) and 4.5 GHz (1''5 \times 2''0 : $S = 12.5 \pm 0.2$ Jy).
 Hughes *et al.* 2000, ApJ, 542, 386. ASCA observations.

McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1'8×2'0), plus H_i. Johnston *et al.* 2004, MNRAS, 348, L19. ATCA at 19 GHz (6''1×7''7), including polarisation. Gelfand *et al.* 2007, ApJ, 663, 468. ATCA at 1.4 GHz (5''8×7''0: S=13.8±0.4 Jy), plus XMM-Newton observations.

G329.7 + 0.4

RA: 16^h01^m20^s
Dec: -52°18'

1-GHz flux/Jy: >34?
Spectral index: ?

Size/arcmin: 40×33
Type: S

Radio: Diffuse shell, in complex region.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×53'' : S > 38).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz (1'8×2'0), plus H_i.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.

G330.0 + 15.0

RA: 15^h10^m00^s
Dec: -40°00'

1-GHz flux/Jy: 350?
Spectral index: 0.5?

Size/arcmin: 180?
Type: S

Lupus Loop

Radio: Low surface brightness loop with H_i shell.

X-ray: Detected, with central source.

Point sources: Central, possibly pulsating, X-ray source.

References:

Milne 1971, AujPh, 24, 757. Parkes 64-m at 408 MHz (48'), 635 MHz (31') and 1410 MHz (15').
 Milne & Dickel 1974, AujPh, 27, 549. Parkes 64-m at 2.7 GHz (8'4 : S=120±30% Jy).
 Toor 1980, A&A, 85, 184. X-ray image and spectrum.
 Colomb & Dubner 1982, A&A, 112, 141. Argentine 30-m dish at 1.42 GHz (30'), H_i observations.
 Leahy *et al.* 1991, ApJ, 374, 218. HEAO-1 X-ray spectra.
 Ozaki *et al.* 1994, PASJ, 46, 367. X-ray observations.
 Kaplan *et al.* 2006, ApJS, 163, 344. X-ray upper limit on compact sources.
 Shinn *et al.* 2006, ApJ, 644, L189. Far UV observations.

G330.2 + 1.0

RA: 16^h01^m06^s
Dec: -51°34'

1-GHz flux/Jy: 5?
Spectral index: 0.3

Size/arcmin: 11
Type: S?

Radio: Clumpy non-thermal emission, possibly a distorted shell.

X-ray: Shell.

Point sources: Central compact X-ray source.

Distance: H_i absorption indicates > 4.9 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : S=8.6 Jy) and Parkes 64-m at 5 GHz (4' : S=4.0 Jy).
 Caswell *et al.* 1983, MNRAS, 204, 915. FIRST at 1415 MHz (47''×52''), and MOST at 843 MHz (43''×55'').

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 55''$): $S = 4.7$ Jy.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ($1'.8 \times 2'.0$), plus H_I.
 Torii *et al.* 2006, PASJ, 58, L11. ASCA detection.
 Park *et al.* 2009, ApJ, 695, 431. XMM-Newton and Chandra observations.
 Abramowski *et al.* 2014, MNRAS, 441, 790. HESS upper limit.
 Doroshenko *et al.* 2018, A&A, 619, A26. XMM-Newton observations of central source.
 Williams *et al.* 2018, ApJ, 855, 118. XMM-Newton observations.
 Borkowski *et al.* 2018, ApJ, 868, L21. Chandra observations for expansion study.

G332.0+0.2

RA: $16^{\text{h}}13^{\text{m}}17^{\text{s}}$
Dec: $-50^{\circ}53'$

1-GHz flux/Jy: 8?
Spectral index: 0.5

Size/arcmin: 12
Type: S

Radio: Incomplete shell.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 57''$).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 55''$): $S = 8.9$ Jy.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 McClure-Griffiths *et al.* 2001, ApJ, 551, 394. ATCA and Parkes 64-m at 1.4 GHz ($1'.8 \times 2'.0$), plus H_I.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.

G332.4-0.4

RA: $16^{\text{h}}17^{\text{m}}33^{\text{s}}$
Dec: $-51^{\circ}02'$

1-GHz flux/Jy: 28
Spectral index: 0.5

RCW 103
Size/arcmin: 10
Type: S

Radio: Shell, brightest to the S.

Optical: Filaments correspond well to the radio shell, brightest in SE.

X-ray: Brightest to NW, with point source near centre.

Point sources: Central, variable X-ray source, and nearby pulsar.

Distance: H_I absorption indicates 3.1 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.
 Tuohy & Garmire 1980, ApJ, 239, L107. Einstein detection of X-ray point source.
 Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz ($50''$).
 Lamb & Markert 1981, ApJ, 244, 94. Einstein observations.
 Ruiz 1983, AJ, 88, 1210. Optical spectra.
 Tuohy *et al.* 1983, ApJ, 268, 778. X-ray observations.
 Leibowitz & Danziger 1983, MNRAS, 204, 273. Optical spectra.
 Nugent *et al.* 1984, ApJ, 284, 612. X-ray spectrum (and Einstein image from Tuohy, private communication).
 Meaburn & Allen 1986, MNRAS, 222, 593. Optical spectra.
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 57''$).
 Oliva *et al.* 1990, A&A, 240, 453. IR spectroscopy.
 Dickel *et al.* 1996, AJ, 111, 340. ATCA at 1.36 (8'') and 2.37 GHz ($4''.5''$), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 55''$): $S = 34$ Jy.
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

Gotthelf *et al.* 1997, ApJ, 487, L175. ASCA observations of compact X-ray source.
 Carter *et al.* 1997, PASP, 109, 990. Optical expansion.
 Torii *et al.* 1998, ApJ, 494, L207. ASCA detection of nearby pulsar.
 Kaspi *et al.* 1998, ApJ, 503, L161. Pulsar observations.
 Gotthelf *et al.* 1999, ApJ, 514, L107. X-ray variability of central source.
 Oliva *et al.* 1999, A&A, 343, 943. ISO spectroscopy.
 Torii *et al.* 2000, ApJ, 534, L71. X-ray timing on pulsar, including glitch.
 Reynoso *et al.* 2004, PASA, 21, 82. ATCA at 1.4 GHz (50''), including H α absorption to central source.
 Russeil *et al.* 2005, A&A, 429, 497. H α observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Paron 2006, PASA, 23, 69. CO and HCO⁺ observations of surroundings.
 De Luca 2006, Science, 313, 814. XMM-Newton observations of central source.
 De Luca 2007, Ap&SS, 308, 231. XMM-Newton observation of periodicity of central source.
 Matsumoto *et al.* 2007, PASJ, 59, S199. Suzaku observations.
 De Luca 2008, ApJ, 682, 1185. IR observations of central source.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Xing *et al.* 2014, ApJ, 781, 64. Fermi observations.
 Frank *et al.* 2015, ApJ, 810, 113. Chandra observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Rea *et al.* 2016, ApJ, 828, L13. Chandra, NuSTAR and Swift observations.
 Borghese *et al.* 2018, MNRAS, 478, 741. NuSTAR and Swift observations of central source.

G332.4 + 0.1

MSH 16–51, Kes 32

RA: 16^h15^m20^s**1-GHz flux/Jy:** 26**Size/arcmin:** 15**Dec:** –50°42'**Spectral index:** 0.5**Type:** S

Has been called G332.4+0.2.

Radio: Distorted shell, with thermal jet and plume adjacent.

Optical: Detected.

X-ray: Shell, brightest to NW.

Point sources: Pulsar nearby.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Roger *et al.* 1985, Nature, 316, 44. MOST at 843 MHz (44''×57'').
 Caraveo 1993, ApJ, 415, L111. Nearby pulsar.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×56'' : S=29 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.
 Brinkmann *et al.* 1999, A&A, 346, 599. ROSAT image of remnant and nearby pulsar.
 Vink *et al.* 2004, ApJ, 604, 693. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G332.5 – 5.6

RA: 16^h43^m20^s
Dec: –54°30′

1-GHz flux/Jy: 2?
Spectral index: 0.7?

Size/arcmin: 35
Type: S

Radio: Bipolar shell, with central emission also.

Optical: Patchy filaments.

X-ray: Emission from centre.

Distance: Optical extinction suggests 3 kpc.

References:

Reynoso & Green 2007, MNRAS, 375, 92. ATCA at 1.4 GHz (40″ : $S = 1.90 \pm 0.15$ Jy) and 2.4 GHz (90″ : $S = 1.3 \pm 0.2$ Jy) including polarisation, and 843 MHz (43″ × 53″) plus ROSAT observations. Stupar *et al.* 2007, MNRAS, 381, 377. Optical observations, plus ATCA at 1.4 GHz (58″) and 2.4 GHz (95″) including polarisation, and other observations. Stupar & Parker 2011, MNRAS, 414, 2282. H α observations. Zhu *et al.* 2015, MNRAS, 452, 3470. Suzaku observations, and distance estimate. Suárez *et al.* 2015, A&A, 583, A84. XMM-Newton observations of central region. Ackermann *et al.* 2018, ApJS, 237, 32. Fermi observations.

G335.2 + 0.1

RA: 16^h27^m45^s
Dec: –48°47′

1-GHz flux/Jy: 16
Spectral index: 0.5

Size/arcmin: 21
Type: S

Radio: Well defined shell.

Point sources: Old pulsar within remnant boundary.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : $S = 27.1$ Jy) and Parkes 64-m at 5 GHz (4′ : $S = 8.6$ Jy). Kaspi *et al.* 1996, AJ, 111, 2028. Pulsar observations. Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ × 57″ : $S = 16$ Jy). Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations. Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection. de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.

G336.7 + 0.5

RA: 16^h32^m11^s
Dec: –47°19′

1-GHz flux/Jy: 6
Spectral index: 0.5

Size/arcmin: 14 × 10
Type: S

Radio: Irregular shell.

Optical: Detected.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′). Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz (44″ × 60″). Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ × 58″ : $S = 6.1$ Jy). Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations. Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G337.0–0.1

(CTB 33)

RA: 16^h35^m57^s**Dec:** –47°36′**1-GHz flux/Jy:** 1.5**Spectral index:** 0.6?**Size/arcmin:** 1.5**Type:** S

This entry refers to a small (1′5) SNR, not the larger previously catalogued G337.0–0.1. Has mistakenly been called G337.7–0.1.

Radio: Shell, in a complex region.

Point sources: Associated with a soft gamma repeater.

Distance: Association with CTB 33 gives 11 kpc.

References:

Shaver & Goss 1970, *AujPA*, 14, 133. Molonglo at 408 MHz (3′) and Parkes 64-m at 5 GHz (4′).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43″×58″:S=21 Jy).
 Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant, including masers.
 Sarma *et al.* 1997, *ApJ*, 483, 335. ATCA at 1.4 GHz (12″) including H_I, and 8.9 GHz (13″×15″) for recombination lines, clarifying extent of the remnant.
 Woods *et al.* 1999, *ApJ*, 519, L139. Soft gamma repeater observations.
 Hurley *et al.* 2000, *ApJ*, 528, L21. ASCA observations of soft gamma repeater.
 Brogan *et al.* 2000, *ApJ*, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Castro *et al.* 2013, *ApJ*, 774, 36. Fermi observations.

G337.2–0.7**RA:** 16^h39^m28^s**Dec:** –47°51′**1-GHz flux/Jy:** 1.5**Spectral index:** 0.4**Size/arcmin:** 6**Type:** S

Radio: Shell, brighter in S.

X-ray: Extended emission.

Distance: H_I absorption suggests 2.0 to 9.3 kpc.

References:

Clark *et al.* 1975, *AujPA*, 37, 1. Molonglo at 408 MHz (3′:S=3.8 Jy) and Parkes 64-m at 5 GHz (4′:S=0.70 Jy).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43″×58″:S=2.0 Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Rakowski *et al.* 2001, *ApJ*, 548, 258. ASCA and ROSAT observations.
 Rakowski *et al.* 2006, *ApJ*, 646, 982. ATCA observations at 1.3 GHz (15″:S=1.55±0.05 Jy) and 5 GHz (15″:S=0.93±0.02 Jy), plus H_I and Chandra observations.
 Yamaguchi *et al.* 2014, *ApJ*, 785, L27. Suzaku observations.
 Takata *et al.* 2016, *PASJ*, 68, S3. Suzaku observations.

G337.2+0.1**RA:** 16^h35^m55^s**Dec:** –47°20′**1-GHz flux/Jy:** 1.5?**Spectral index:** ?**Size/arcmin:** 3×2**Type:** ?

Radio: Not well defined.

X-ray: Detected.

Distance: Association with H_I hole gives 14 kpc.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59''$: $S = 1.6$ Jy)
 Combi *et al.* 2005, A&A, 431, L9. ASCA and other observations.
 Aharonian *et al.* 2006, ApJ, 636, 777. HESS observations of nearby source.
 Combi *et al.* 2006, ApJ, 653, L41. XMM-Newton observations.

G337.3 + 1.0

Kes 40

RA: $16^{\text{h}}32^{\text{m}}39^{\text{s}}$
Dec: $-46^{\circ}36'$

1-GHz flux/Jy: 16
Spectral index: 0.55

Size/arcmin: 15×12
Type: S

Radio: Nearly complete shell.

References:

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz ($3'$: $S = 24.6$ Jy) and Parkes 64-m at 5 GHz ($4'$: $S = 7.2$ Jy).
 Dickel & Milne 1976, AujPh, 29, 435. Comparison of earlier Parkes 64-m maps at 2.7 GHz ($8'.4$) and 5 GHz ($4'.4$).
 Kesteven & Caswell 1987, A&A, 183, 118. MOST at 843 MHz ($44'' \times 61''$).
 Milne *et al.* 1989, PASA, 8, 187. MOST at 843 MHz ($43'' \times 59''$: $S = 14.8 \pm 3.0$ Jy), and Parkes 64-m at 8.4 GHz ($3'$: $S = 5.1 \pm 0.6$ Jy), including polarisation.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59''$: $S = 20$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G337.8 – 0.1

Kes 41

RA: $16^{\text{h}}39^{\text{m}}01^{\text{s}}$
Dec: $-46^{\circ}59'$

1-GHz flux/Jy: 15
Spectral index: 0.5

Size/arcmin: 9×6
Type: S

Radio: Distorted shell.

X-ray: Centrally brightened.

Distance: H α absorption suggests 11 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz ($3'$) and Parkes 64-m at 5 GHz ($4'$).
 Caswell *et al.* 1975, A&A, 45, 239. Parkes H α absorption.
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 59''$: $S = 18$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA detection of compact OH emission.
 Caswell 2004, MNRAS, 349, 99. ATCA at 1.7 GHz, for associated OH masers.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including H α .
 Combi *et al.* 2008, A&A, 488, L25. XMM-Newton observations.
 Zhang *et al.* 2015, ApJ, 799, 103. XMM-Newton, Chandra and CO observations.
 Liu *et al.* 2015, ApJ, 809, 102. Fermi observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Qiao *et al.* 2016, ApJS, 227, 26. OH maser observations.
 Supan *et al.* 2018, A&A, 619, A108. CO, H α and Spitzer observations.
 Supan *et al.* 2018, A&A, 619, A109. Fermi observations and review of radio flux densities.

G338.1 + 0.4**RA:** 16^h37^m59^s**Dec:** -46°24'**1-GHz flux/Jy:** 4?**Spectral index:** 0.4**Size/arcmin:** 15?**Type:** S**Radio:** Arc in NE, merging with thermal emission in S.**Optical:** Detected.**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Zealey *et al.* 1979, A&AS, 38, 39. Optical detection.

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×59'':S=3.8 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G338.3 – 0.0****RA:** 16^h41^m00^s**Dec:** -46°34'**1-GHz flux/Jy:** 7?**Spectral index:** ?**Size/arcmin:** 8**Type:** C?**Radio:** Irregular shell, in complex region.**X-ray:** Central X-ray source and nebula.**Point sources:** Central pulsar.**Distance:** H_I observations suggest 8 to 13 kpc.**References:**

Shaver & Goss 1970, AuJPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×59'':S=7.4 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.Aharonian *et al.* 2005, Science, 307, 1938. HESS detection.Aharonian *et al.* 2006, ApJ, 636, 777. HESS observations.Landi *et al.* 2006, ApJ, 651, 190. X-ray observations.Funk *et al.* 2007, ApJ, 662, 517. XMM-Newton observations.Lemiere *et al.* 2009, ApJ, 706, 1269. Chandra observations of central source and nebula.Slane *et al.* 2010, ApJ, 720, 266. Fermi observations.Castelletti *et al.* 2011, A&A, 536, A98. GMRT at 235 MHz (10''×26''), 610 MHz (5''0×12''6) and 1280 MHz (5''0×6''4), plus ATCA at 2.3 GHz (4''3×4''8).Gotthelf *et al.* 2014, ApJ, 788, 155. NuSTAR observations.Lemoine-Goumard *et al.* 2014, ApJ, 794, L16. Fermi observations.Abramowski *et al.* 2014, MNRAS, 439, 2828. HESS observations.*see also:* Abramowski *et al.* 2014, MNRAS, 441, 3640. Erratum.Supan *et al.* 2016, A&A, 589, A51. SGPS H_I and other observations.Lau *et al.* 2017, MNRAS, 464, 3757. CO and other molecular line observations, plus H_I observations.de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.Xin *et al.* 2018, ApJ, 867, 55. Fermi observations.

G338.5 + 0.1

RA: 16^h41^m09^s
Dec: -46°19'

1-GHz flux/Jy: 12?
Spectral index: ?

Size/arcmin: 9
Type: ?

Radio: Circle of non-thermal emission in complex region, not well defined.

Distance: H_I absorption suggests 11 kpc.

References:

Shaver & Goss 1970, AujPA, 14, 133. Molonglo at 408 MHz (3') and Parkes 64-m at 5 GHz (4').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×59'' : S=13 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including H_I.
 Abramowski *et al.* 2014, ApJ, 794, L1. HESS observations.
 Lau *et al.* 2017, MNRAS, 464, 3757. CO and other molecular line observations, plus H_I observations.
 de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.

G340.4 + 0.4

RA: 16^h46^m31^s
Dec: -44°39'

1-GHz flux/Jy: 5
Spectral index: 0.4

Size/arcmin: 10×7
Type: S

Radio: Distorted shell, elongated east--west.

Optical: Detected.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : S=8.2 Jy) and Parkes 64-m at 5 GHz (4' : S=2.9 Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50'').
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (48''×75'' : S=9.8±0.9 Jy) and 1.4 GHz (9''×27'' : S=3.6±0.1 Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×61'' : S=5.9 Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

G340.6 + 0.3

RA: 16^h47^m41^s
Dec: -44°34'

1-GHz flux/Jy: 5?
Spectral index: 0.4?

Size/arcmin: 6
Type: S

Radio: Incomplete shell.

Optical: Possible associated filaments.

Distance: H_I absorption suggests 15 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3' : S=7.0 Jy) and Parkes 64-m at 5 GHz (4' : S=2.8 Jy).
 Zealey *et al.* 1979, A&AS, 38, 39. Optical observations.
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50'').
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 330 MHz (48''×75'' : S=9.2±0.9 Jy) and 1.4 GHz (9''×27'' : S=5.8±0.1 Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×61'' : S=4.5 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer possible detection.
 Kothes & Dougherty 2007, A&A, 468, 993. SGPS at 1.4 GHz including Hi.

G341.2 + 0.9

RA: 16^h47^m35^s
Dec: -43°47'

1-GHz flux/Jy: 1.5?
Spectral index: 0.6?

Size/arcmin: 22×16
Type: C

Radio: Incomplete shell, with extension to SW.

Point sources: Pulsar in W, with wind nebula.

References:

Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (54''×116'': $S=3.0\pm 0.1$ Jy) and 1.4 GHz (21''×25'': $S=12.5\pm 0.05$ Jy).
 Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4 and 4.9 GHz (25'') of pulsar wind nebula.

G341.9 – 0.3

RA: 16^h55^m01^s
Dec: -44°01'

1-GHz flux/Jy: 2.5
Spectral index: 0.5

Size/arcmin: 7
Type: S

Radio: Incomplete shell, brightest to NE.

References:

Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3': $S=7.4$ Jy) and Parkes 64-m at 5 GHz (4': $S=1.7$ Jy).
 Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50''), revision of previous flux densities.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (11''×35'': $S=2.2\pm 0.1$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×62'': $S=2.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.0 – 0.2

RA: 16^h54^m50^s
Dec: -43°53'

1-GHz flux/Jy: 3.5?
Spectral index: 0.4?

Size/arcmin: 12×9
Type: S

Radio: Distorted shell.

References:

Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (50''), estimate $S_{408\text{ MHz}}=5$ Jy, $S_{5\text{ GHz}}=2$ Jy from previous maps.
 Dubner *et al.* 1996, AJ, 111, 1304. VLA at 1.4 GHz (11''×35'').
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×62'': $S=3.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G342.1 + 0.9**RA:** 16^h50^m43^s**Dec:** -43°04′**1-GHz flux/Jy:** 0.5?**Spectral index:** ?**Size/arcmin:** 10×9**Type:** S**Radio:** Incomplete shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×63″: S=0.6 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G343.0 – 6.0****RA:** 17^h25^m00^s**Dec:** -46°30′**1-GHz flux/Jy:** ?**Spectral index:** ?

RCW 114

Size/arcmin: 250**Type:** S**Radio:** Faint, poorly defined.**Optical:** Filamentary shell.**References:**

Walker & Zealey 2001, MNRAS, 325, 287. Optical observations, and review of earlier observations.

Welsh *et al.* 2003, A&A, 403, 605. Optical spectroscopy.Casandjian & Grenier 2008, A&A, 489, 849. γ -ray observations.Kim *et al.* 2010, ApJ, 709, 823. UV observations.**G343.1 – 2.3****RA:** 17^h08^m00^s**Dec:** -44°16′**1-GHz flux/Jy:** 8?**Spectral index:** 0.5?**Size/arcmin:** 32?**Type:** C?**Radio:** Incomplete shell?**X-ray:** Pulsar wind nebula.**Point sources:** Pulsar near edge, with wind nebula.**References:**McAdam *et al.* 1993, Nature, 361, 516. MOST at 843 MHz (smoothed to 2′).Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (56″×111″: S=10.6 Jy) and 1.4 GHz (22″×27″) near pulsar.Becker *et al.* 1995, A&A, 298, 528. ROSAT of pulsar, and limit for remnant.Giacani *et al.* 2001, AJ, 121, 3133. VLA at 1.4, 4.9 and 8.5 GHz (25″) of pulsar wind nebula.Gotthelf *et al.* 2002, ApJ, 567, L125. Pulsar x-ray detection.

Dodson & Golap 2002, MNRAS, 334, L1. ATCA at 1.4 GHz (47″×70″) including polarisation, and Chandra observations of pulsar wind nebula.

Aharonian *et al.* 2005, A&A, 432, L9. HESS limit.Abramowski *et al.* 2011, A&A, 528, A143. HESS observations.

G343.1–0.7**RA:** 17^h00^m25^s**Dec:** –43°14′**1-GHz flux/Jy:** 7.8**Spectral index:** 0.55**Size/arcmin:** 27×21**Type:** S**Radio:** Shell, with smaller thermal shell adjacent.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×63'': $S=58.5\pm 0.6$), plus Parkes 64-m at 4.5 GHz ($S=3.9\pm 0.6$ Jy) and 8.55 GHz ($S=2.4\pm 0.5$ Jy)
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.

G344.7–0.1**RA:** 17^h03^m51^s**Dec:** –41°42′**1-GHz flux/Jy:** 2.5?**Spectral index:** 0.3?**Size/arcmin:** 8**Type:** C?**Radio:** Asymmetric shell, with possible core.**X-ray:** Detected.**Distance:** H_I absorption and association with features suggests 6.3 kpc.**References:**

Clark *et al.* 1975, AuJPA, 37, 1. Molonglo at 408 MHz (3': $S=4.7$ Jy) and Parkes 64-m at 5 GHz (4': $S=1.3$ Jy).
 Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (30''×43'': $S=1.7\pm 0.1$ Jy).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×65'': $S=2.5$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Yamauchi *et al.* 2005, PASJ, 57, 459. ASCA observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Giacani *et al.* 2011, A&A, 531, A138. VLA and ATCA at 1.4 GHz (5''6×8''), ATCA at 5 GHz (10''×13'') and 8.4 GHz (5''×10''), plus XMM-Newton observations and H_I from SGPS.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Yamaguchi *et al.* 2012, ApJ, 749, 137. Suzaku observations.
 Yang *et al.* 2013, ApJ, 766, 44. Suzaku spectroscopy.

G345.7–0.2**RA:** 17^h07^m20^s**Dec:** –40°53′**1-GHz flux/Jy:** 0.6?**Spectral index:** ?**Size/arcmin:** 6**Type:** S**Radio:** Poorly defined diffuse shell.**Point sources:** Old pulsar nearby.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×66'': $S=0.7$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.

G346.6–0.2**RA:** 17^h10^m19^s**Dec:** –40°11′**1-GHz flux/Jy:** 8?**Spectral index:** 0.5?**Size/arcmin:** 8**Type:** S**Radio:** Irregular shell.**X-ray:** Centrally brightened, clumpy.**References:**

- Clark *et al.* 1975, *AJPA*, 37, 1. Molonglo at 408 MHz (3′ : S = 14.9 Jy) and Parkes 64-m at 5 GHz (4′ : S = 4.3 Jy).
 Dubner *et al.* 1993, *AJ*, 105, 2251. VLA at 1.47 GHz (31″ × 43″ : S = 8.1 ± 0.9 Jy).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz (43″ × 67″ : S = 8.7 Jy).
 Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, *AJ*, 116, 1323. VLA detection of compact OH emission.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2009, *ApJ*, 694, 1266. Spitzer spectroscopy.
 Sezer *et al.* 2011, *MNRAS*, 415, 301. Suzaku observations.
 Andersen *et al.* 2011, *ApJ*, 742, 7. Spitzer observations.
 Yamauchi *et al.* 2013, *PASJ*, 65, 6. Suzaku observations.
 Pihlström *et al.* 2014, *AJ*, 147, 73. VLA search for methanol masers.
 Pannuti *et al.* 2014, *AJ*, 147, 55. ASCA observations.
 Auchettl *et al.* 2017, *ApJ*, 847, 121. XMM-Newton observations.

G347.3–0.5**RA:** 17^h13^m50^s**Dec:** –39°45′**1-GHz flux/Jy:** 30?**Spectral index:** ?

RX J1713.7–3946

Size/arcmin: 65 × 55**Type:** S?**Radio:** Faint emission.**X-ray:** Non-thermal, limb-brightened to W, with central source.**Point sources:** Central X-ray source.**Distance:** Association with molecular clouds and X-ray observations imply 1.3 kpc.**References:**

- Koyama *et al.* 1997, *PASJ*, 49, L7. ASCA of NW.
 Slane *et al.* 1999, *ApJ*, 525, 357. ASCA and other observations.
 Muraishi *et al.* 2000, *A&A*, 354, L57. γ -ray detection.
 Butt *et al.* 2001, *ApJ*, 562, L167. Associated γ -ray emission.
 Ellison *et al.* 2001, *ApJ*, 563, 191. ATCA at 1.4 GHz (36″ × 46″), and ASCA observations.
 Enomoto *et al.* 2002, *Nature*, 416, 823. γ -ray observations.
 Uchiyama *et al.* 2002, *PASJ*, 54, L73. ASCA observations.
 Uchiyama *et al.* 2003, *A&A*, 400, 567. Chandra spectroscopy.
 Pannuti *et al.* 2003, *ApJ*, 593, 377. ROSAT, ASCA and RXTE observations.
 Lazendic *et al.* 2003, *ApJ*, 593, L27. Chandra, XMM-Newton and other X-ray observations of central source.
 Fukui *et al.* 2003, *PASJ*, 55, L61. CO observations of surroundings.
 Cassam-Chenāi *et al.* 2004, *A&A*, 427, 199. XMM-Newton and other observations.
 Aharonian *et al.* 2004, *Nature*, 432, 75. HESS detection.
 Lazendic *et al.* 2004, *ApJ*, 602, 271. Chandra observations of parts, and ATCA at 1.4 GHz (36″ × 46″).
 Hiraga *et al.* 2005, *A&A*, 431, 953. XMM-Newton observations,
 Moriguchi *et al.* 2005, *ApJ*, 631, 947. CO observations of surroundings.

Aharonian *et al.* 2007, A&A, 464, 235. HESS observations.
see also: Aharonian *et al.* 2011, A&A, 531, C1. Erratum.
 Uchiyama *et al.* 2007, Nature, 449, 576. Chandra multi-epoch observations for study of small-scale variability.
 Tanaka *et al.* 2008, ApJ, 685, 988. Suzaku observations.
 Takahashi *et al.* 2008, PASJ, 60, S131. Suzaku observations of SW.
 Mignani *et al.* 2008, A&A, 484, 457. Optical and IR observations of central source.
 Acero *et al.* 2009, A&A, 505, 157. XMM-Newton observations, plus other radio, IR and HESS observations.
 Sano *et al.* 2010, ApJ, 724, 59. CO observations of region.
 Maxted *et al.* 2012, MNRAS, 422, 2230. Molecular line observations.
see also: Maxted *et al.* 2013, MNRAS, 430, 2511. Erratum.
 Sano *et al.* 2013, ApJ, 778, 59. CO, SGPS H_I and Suzaku observations.
 Federici *et al.* 2015, A&A, 577, A12. Fermi detection.
 Sano *et al.* 2015, ApJ, 799, 175. Suzaku observations.
 Katsuda *et al.* 2015, ApJ, 814, 29. XMM-Newton observations.
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Tsuji & Uchiyama 2016, PASJ, 68, 108. Chandra observations of NW.
 Acero *et al.* 2017, A&A, 597, A106. XMM-Newton observations for proper motion studies.
 Okuno *et al.* 2018, PASJ, 70, 77. Chandra observations.
 Abdalla *et al.* 2018, A&A, 612, A6. HESS observations.

G348.5–0.0

RA: 17^h15^m26^s
Dec: –38°28′

1-GHz flux/Jy: 10?
Spectral index: 0.4?

Size/arcmin: 10?
Type: S?

Radio: Arc, overlapping G348.5+0.1.

References:

Kassim *et al.* 1991, ApJ, 374, 212. VLA at 333 MHz (46″×53″), 1.4 GHz (18″×33″) and part at 5 GHz (2″5×3″9).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×69″:S=10.2 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Pannuti *et al.* 2014, AJ, 147, 55. XMM-Newton upper limit.

G348.5+0.1

RA: 17^h14^m06^s
Dec: –38°32′

1-GHz flux/Jy: 72
Spectral index: 0.3

Size/arcmin: 15
Type: S

CTB 37A

Radio: Shell, poorly define to S and W, overlapping G348.5–0.0 in E.

X-ray: Brighter to W.

Distance: H_I absorption indicates 8.0 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 75. Molonglo at 408 MHz (3′:S=97 Jy).

Milne & Dickel 1975, *AJPh*, 28, 209. Parkes 64-m at 5 GHz ($4'.4:S=43$ Jy).
 Caswell *et al.* 1975, *A&A*, 45, 239. Parkes H α absorption.
 Milne *et al.* 1979, *MNRAS*, 188, 437. FIRST at 1415 MHz ($0'.8:S>50$) and Parkes 64-m at 14.7 GHz ($2'.2:S=18\pm 5$ Jy).
 Downes 1984, *MNRAS*, 210, 845. VLA at 1465 MHz ($20''\times 45''$).
 Kassim *et al.* 1991, *ApJ*, 374, 212. VLA at 333 MHz ($46''\times 53''$), 1.4 GHz ($18''\times 33''$) and part at 5 GHz ($2''5\times 3''9$).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz ($43''\times 69'' : S=71$ Jy).
 Brogan *et al.* 2000, *ApJ*, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Reynoso & Mangum 2000, *ApJ*, 545, 874. CO observations.
 Reach *et al.* 2006, *AJ*, 131, 1479. Spitzer observations.
 Aharonian *et al.* 2008, *A&A*, 490, 685. HESS observations.
 Castro & Slane 2010, *ApJ*, 717, 372. Fermi observations.
 Andersen *et al.* 2011, *ApJ*, 742, 7. Spitzer observations.
 Sezer *et al.* 2011, *MNRAS*, 417, 1387. Suzaku observations.
 Tian & Leahy 2012, *MNRAS*, 421, 2593. SGPS H α absorption observations.
 Maxted *et al.* 2013, *MNRAS*, 434, 2188. Molecular line observations of region.
 Pannuti *et al.* 2014, *AJ*, 147, 55. XMM-Newton and Chandra observations.
 Yamauchi *et al.* 2014, *PASJ*, 66, 2. Suzaku observations.
 Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.

G348.7 + 0.3

CTB 37B

RA: $17^{\text{h}}13^{\text{m}}55^{\text{s}}$
Dec: $-38^{\circ}11'$

1-GHz flux/Jy: 26
Spectral index: 0.3

Size/arcmin: 17?
Type: S

Radio: Incomplete shell with faint eastern extensions.

X-ray: Diffuse emission.

Point sources: X-ray pulsar.

Distance: H α absorption suggests 13 kpc.

References:

Clark *et al.* 1975, *AJPA*, 37, 75. Molonglo at 408 MHz ($3':S=34$ Jy).
 Milne & Dickel 1975, *AJPh*, 28, 209. Parkes 64-m at 5 GHz ($4'.4:S=32$ Jy).
 Caswell *et al.* 1975, *A&A*, 45, 239. Parkes H α absorption.
 Milne *et al.* 1979, *MNRAS*, 188, 437. FIRST at 1415 MHz ($0'.8:S>20$) and Parkes 64-m at 14.7 GHz ($2'.2:S=8\pm 3$ Jy).
 Downes 1984, *MNRAS*, 210, 845. VLA at 1465 MHz ($20''\times 45''$).
 Kassim *et al.* 1991, *ApJ*, 374, 212. VLA at 333 MHz ($46''\times 53''$).
 Whiteoak & Green 1996, *A&AS*, 118, 329. MOST at 843 MHz ($43''\times 69'' : S=33$ Jy).
 Frail *et al.* 1996, *AJ*, 111, 1651. OH emission near remnant.
 Aharonian *et al.* 2006, *ApJ*, 636, 777. HESS detection.
 Aharonian *et al.* 2008, *A&A*, 486, 829. HESS and Chandra observations.
 Nakamura *et al.* 2009, *PASJ*, 61, S197. Suzaku and Chandra observations.
 Halpern & Gotthelf 2010, *ApJ*, 710, 941. Chandra detection of pulsar.
 Halpern & Gotthelf 2010, *ApJ*, 725, 1384. Chandra observations of pulsar.
 Tian & Leahy 2012, *MNRAS*, 421, 2593. SGPS H α observations.
 Xin *et al.* 2016, *ApJ*, 817, 64. Fermi observations.

G349.2–0.1

RA: 17^h17^m15^s
Dec: –38°04′

1-GHz flux/Jy: 1.4?
Spectral index: ?

Size/arcmin: 9×6
Type: S

Radio: Elongated shell, adjacent to bright H_{II} region.

References:

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×70″:S=1.6 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant.

G349.7+0.2

RA: 17^h17^m59^s
Dec: –37°26′

1-GHz flux/Jy: 20
Spectral index: 0.5

Size/arcmin: 2.5×2
Type: S

Radio: Incomplete clumpy shell, with enhancement to the S.

X-ray: Irregular shell, brighter to S and E.

Distance: H_I absorption indicates 11.5 kpc.

References:

Becker & Kundu 1975, AJ, 80, 679. NRAO 140-ft at 10.6 GHz (3′).
 Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′:S=31.0 Jy) and Parkes 64-m at 5 GHz (4′:S=9.1 Jy), no maps.
 Caswell *et al.* 1975, A&A, 45, 239. Parkes H_I absorption.
 Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz (3′4×14′5).
 Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×71″:S=22 Jy).
 Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
 Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
 Reynoso & Mangum 2001, AJ, 121, 347. CO observations of the vicinity.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Slane *et al.* 2002, ApJ, 580, 904. ASCA observations.
 Dubner *et al.* 2004, A&A, 426, 201. CO observations of surroundings.
 Lazendic *et al.* 2005, ApJ, 618, 733. Chandra observations.
 Reach *et al.* 2006, AJ, 131, 1479. Spitzer observations.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Hewitt *et al.* 2009, ApJ, 694, 1266. Spitzer spectroscopy.
 Castro & Slane 2010, ApJ, 717, 372. Fermi observations.
 Lazendic *et al.* 2010, MNRAS, 409, 371. OH, CO and other molecular line observations of region.
 Andersen *et al.* 2011, ApJ, 742, 7. Spitzer observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Tian & Leahy 2014, ApJ, 783, L2. SGPS H_I observations.
 Yasumi *et al.* 2014, PASJ, 66, 68. Suzaku observations.
 Abramowski *et al.* 2015, A&A, 574, A100. HESS observations.
see also: Abramowski *et al.* 2015, A&A, 580, C1. Corrigendum.
 Ergin *et al.* 2015, ApJ, 804, 124. Fermi and Suzaku observations.
 Rho *et al.* 2015, ApJ, 812, 44. Herschel observations.
 Koo *et al.* 2016, ApJ, 821, 20. Spitzer and Herschel flux densities (and comparison with X-ray properties).
 Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.

G350.0–2.0**RA:** 17^h27^m50^s**Dec:** –38°32′**1-GHz flux/Jy:** 26**Spectral index:** 0.4**Size/arcmin:** 45**Type:** S

Incorporates the previously catalogued G350.0–1.8 in the NW.

Radio: Shell, brightest in NW.

Optical: Detected.

X-ray: Brighter to NW.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : S = 49.5 Jy) and Parkes 64-m at 5 GHz (4′ : S = 13.6 Jy).

Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4′.4).

Gaensler 1998, ApJ, 493, 781. VLA and Parkes 64-m at 1.4 GHz (18″ × 21″ : S = 22.3 ± 0.3 Jy), clarifying extent of remnant.

Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.

Karpova *et al.* 2016, MNRAS, 462, 3845. XMM-Newton observations.

G350.1–0.3**RA:** 17^h21^m05^s**Dec:** –37°27′**1-GHz flux/Jy:** 6?**Spectral index:** 0.8?**Size/arcmin:** 4?**Type:** ?

Radio: Several clumps of emission.

X-ray: Diffuse emission, with compact source.

Point sources: X-ray source.

Distance: H α absorption indicates 4.5 to 10.7 kpc, possible interaction with molecular cloud indicates 4.5 kpc.

References:

Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′ : S = 10.7) and Parkes 64-m at 5 GHz (4′ : S = 1.7).

Salter *et al.* 1986, A&A, 162, 217. VLA at 1.5 (4″ × 10″.4) and 4.8 GHz (15″ × 35″) and 15 GHz (2″ × 5″).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″ × 71″)..

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Gaensler *et al.* 2008, ApJ, 680, L37. VLA at 4.8 GHz (5″.6 × 11″.4) plus XMM-Newton observations.

Lovchinsky *et al.* 2011, ApJ, 731, 70. Chandra and Spitzer observations.

Yasumi *et al.* 2014, PASJ, 66, 68. Suzaku observations.

G351.0–5.4**RA:** 17^h46^m00^s**Dec:** –39°25′**1-GHz flux/Jy:** ?**Spectral index:** ?**Size/arcmin:** 30**Type:** S

Radio: Faint shell, brighter to E and W.

References:

de Gasperin *et al.* 2014, A&A, 568, A107. GMRT at 325 MHz (89″ × 141″), plus other observations.

G351.2 + 0.1**RA:** 17^h22^m27^s**Dec:** −36°11′**1-GHz flux/Jy:** 5?**Spectral index:** 0.4**Size/arcmin:** 7**Type:** C?

Has been called G351.3+0.2.

Radio: Distorted shell, with possible flat-spectrum core.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=8.1$ Jy) and Parkes 64-m at 5 GHz (4′: $S=3.1$ Jy).

Becker & Helfand 1988, AJ, 95, 883. VLA at 5 GHz (15″), and at 15 GHz of core.

Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (32″×36″: $S=4.8\pm0.2$ Jy).Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×73″: $S=5.5$ Jy).Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.**G351.7 + 0.8****RA:** 17^h21^m00^s**Dec:** −35°27′**1-GHz flux/Jy:** 10**Spectral index:** 0.5?**Size/arcmin:** 18×14**Type:** S**Radio:** Elongated shell, adjacent to bright H_{II} region.**Point sources:** Pulsar nearby.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×74″: $S=11$ Jy).Tian *et al.* 2007, MNRAS, 378, 1283. SGPS at 1.4 GHz (100″: $S=8.4\pm0.7$ Jy) including H_I.**G351.9 − 0.9****RA:** 17^h28^m52^s**Dec:** −36°16′**1-GHz flux/Jy:** 1.8?**Spectral index:** ?**Size/arcmin:** 12×9**Type:** S**Radio:** Asymmetric shell.**References:**Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43″×73″: $S=2.0$ Jy).**G352.7 − 0.1****RA:** 17^h27^m40^s**Dec:** −35°07′**1-GHz flux/Jy:** 4**Spectral index:** 0.6**Size/arcmin:** 8×6**Type:** S**Radio:** Distorted shell.**X-ray:** Detected.**Distance:** H_I absorption indicates 6.8 to 8.4 kpc.**References:**Clark *et al.* 1975, AujPA, 37, 1. Molonglo at 408 MHz (3′: $S=9.6$ Jy) and Parkes 64-m at 5 GHz (4′: $S=2.3$ Jy).Caswell *et al.* 1983, MNRAS, 203, 595. FIRST at 1415 MHz (1′.1).Dubner *et al.* 1993, AJ, 105, 2251. VLA at 1.47 GHz (34″: $S=3.4\pm0.4$ Jy).

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz ($43'' \times 75''$; $S = 4.4$ Jy).
 Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.
 Kinugasa *et al.* 1998, PASJ, 50, 249. ASCA observations.
 Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.
 Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.
 Giacani *et al.* 2009, A&A, 507, 841. VLA at 4.8 GHz ($9'' \times 12''$), plus H α and XMM-Newton observations.
 Pannuti *et al.* 2014, ApJ, 782, 102. XMM-Newton and Chandra observations.
 Sezer & Gök 2014, ApJ, 790, 81. Suzaku observations.

G353.6–0.7

RA: $17^{\text{h}}32^{\text{m}}00^{\text{s}}$
Dec: $-34^{\circ}44'$

1-GHz flux/Jy: 2.5?
Spectral index: ?

Size/arcmin: 30
Type: S

Has erroneously been called G353.6–0.37.

Radio: Shell, brighter to S.

X-ray: Patchy shell, brighter to E.

Point sources: Central X-ray source.

Distance: Various observations suggest 3.2 kpc.

References:

Tian *et al.* 2008, ApJ, 679, L85. SGPS at 1.4 GHz ($100''$), plus 843 MHz ($43''$) and X-ray observations.
 Halpern & Gotthelf 2010, ApJ, 710, 941. XMM-Newton observations.
 Tian *et al.* 2010, ApJ, 712, 790. XMM-Newton, Suzaku and CO observations.
 Abramowski *et al.* 2011, A&A, 531, A81. HESS observations.
 Bamba *et al.* 2012, ApJ, 756, 149. Suzaku observations.
see also: Bamba *et al.* 2012, ApJ, 761, 80. Erratum.
 Klochkov *et al.* 2013, A&A, 556, A41. XMM-Newton and other X-ray observations.
 Fukuda *et al.* 2014, ApJ, 788, 94. HESS, SGPS H α and CO observations.
 Nayana *et al.* 2017, MNRAS, 467, 155. GMRT at 325 MHz ($97'' \times 135''$) and 610 MHz ($105'' \times 150''$), plus other observations.
 de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.
 Doroshenko *et al.* 2017, A&A, 608, A23. XMM-Newton observations.
 Condon *et al.* 2017, ApJ, 851, 100. Fermi observations.
 Guo *et al.* 2018, ApJ, 853, 2. Fermi observations.
 Abdalla *et al.* 2018, A&A, 612, A1. HESS observations.
 Maxted *et al.* 2018, MNRAS, 474, 662. CO and H α observations.

G353.9–2.0

RA: $17^{\text{h}}38^{\text{m}}55^{\text{s}}$
Dec: $-35^{\circ}11'$

1-GHz flux/Jy: 1?
Spectral index: 0.5?

Size/arcmin: 13
Type: S

Radio: Shell, with central double source.

References:

Green 2001, MNRAS, 326, 283. VLA at 327 MHz ($2'.7 \times 3'.0$) and 1.4 GHz ($36'' \times 42''$), plus 8.4 GHz ($6'' \times 8''.4$) of central source only.

G354.1 + 0.1**RA:** 17^h30^m28^s**Dec:** −33°46′**1-GHz flux/Jy:** ?
Spectral index: varies**Size/arcmin:** 15×3?**Type:** C?

Is this a SNR?

Radio: Elongated N--S.**Point sources:** Pulsar at S tip.**References:**Frail *et al.* 1994, ApJ, 437, 781. VLA at 330 MHz (47''×99'') and 1.4 GHz (8''×21'').Ajello *et al.* 2016, ApJ, 819, 44. Fermi observations.**G354.8 − 0.8****RA:** 17^h36^m00^s**Dec:** −33°42′**1-GHz flux/Jy:** 2.8?
Spectral index: ?**Size/arcmin:** 19**Type:** S**Radio:** Distorted shell.**References:**

Whiteoak & Green 1996, A&AS, 118, 329. MOST at 843 MHz (43''×78'':S=3.1 Jy).

Green *et al.* 1997, AJ, 114, 2058. Parkes 64-m OH observations.Koralesky *et al.* 1998, AJ, 116, 1323. VLA search for OH emission.**G355.4 + 0.7****RA:** 17^h31^m20^s**Dec:** −32°26′**1-GHz flux/Jy:** 5?
Spectral index: ?**Size/arcmin:** 25**Type:** S**Radio:** Faint, incomplete shell.**References:**

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43''×77'').

Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1':S=8.9±1.3 Jy).

Ajello *et al.* 2016, ApJ, 819, 44. Fermi observations.Acero *et al.* 2016, ApJS, 224, 8. Fermi observations.**G355.6 − 0.0****RA:** 17^h35^m16^s**Dec:** −32°38′**1-GHz flux/Jy:** 3?
Spectral index: ?**Size/arcmin:** 8×6**Type:** S**Radio:** Well defined shell.**X-ray:** Centrally brightened.**References:**

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×80'':S=2.6 Jy).

Sugizaki *et al.* 2001, ApJS, 134, 77. ASCA survey observations.

Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1':S=3.3±0.5 Jy).

Marquez-Lopez & Phillips 2010, MNRAS, 407, 94. Mid-IR observations.

Minami *et al.* 2013, PASJ, 65, 99. Suzaku observations.

G355.9–2.5**RA:** 17^h45^m53^s
Dec: –33°43′**1-GHz flux/Jy:** 8
Spectral index: 0.5**Size/arcmin:** 13
Type: S**Radio:** Distorted shell, brightest to SE.**References:**Clark *et al.* 1975, *AujPA*, 37, 1. Molonglo at 408 MHz (3′ : $S=12.3$ Jy) and Parkes 64-m at 5 GHz (4′ : $S=3.4$ Jy).Dubner *et al.* 1993, *AJ*, 105, 2251. VLA at 1.47 GHz (32″×34″ : $S=5.0\pm0.3$ Jy).Gray 1994, *MNRAS*, 270, 835. MOST at 843 MHz (43″×77″).Green *et al.* 1997, *AJ*, 114, 2058. Parkes 64-m OH observations.Marquez-Lopez & Phillips 2010, *MNRAS*, 407, 94. Mid-IR observations.

G356.2+4.5**RA:** 17^h19^m00^s
Dec: –29°40′**1-GHz flux/Jy:** 4
Spectral index: 0.7**Size/arcmin:** 25
Type: S

Has been called G356.2+4.4.

Radio: Faint shell.**References:**Duncan *et al.* 1995, *MNRAS*, 277, 36. Parkes 64-m at 2.4-GHz (10′.4).Bhatnagar 2000, *MNRAS*, 317, 453. GMRT at 327 MHz (1′.5×3′ : $S=8.1\pm1.7$ Jy), and NVSS at 1.4 GHz.

G356.3–1.5**RA:** 17^h42^m35^s
Dec: –32°52′**1-GHz flux/Jy:** 3?
Spectral index: ?**Size/arcmin:** 20×15
Type: S**Radio:** Double arc.**References:**Gray 1994, *MNRAS*, 270, 847. MOST at 843 MHz (43″×79″ : $S=2.8$ Jy).Bhatnagar 2002, *MNRAS*, 332, 1. GMRT at 327 MHz (0′.8×1′.7 : $S=5.7\pm0.2$).

G356.3–0.3**RA:** 17^h37^m56^s
Dec: –32°16′**1-GHz flux/Jy:** 3?
Spectral index: ?**Size/arcmin:** 11×7
Type: S

Has been suggested this part of a larger SNR.

Radio: Diffuse emission.**References:**Gray 1994, *MNRAS*, 270, 847. MOST at 843 MHz (43″×81″ : $S=2.6$ Jy).Roy & Pramesh Rao 2002, *MNRAS*, 329, 775. GMRT at 330 MHz (2′.7×4′.8).Acero *et al.* 2016, *ApJS*, 224, 8. Fermi observations.

G357.7–0.1

MSH 17–39

RA: 17^h40^m29^s
Dec: –30°58′**1-GHz flux/Jy:** 37
Spectral index: 0.4**Size/arcmin:** 8×3?
Type: ?

Has been suggested that this is not a SNR.

Radio: Multiple arcs and filaments, brighter to NW ‘head’.

X-ray: Detected from NW ‘head’, and SW ‘tail’.

Distance: H_i absorption suggests beyond Galactic Centre.

References:

- Caswell *et al.* 1975, AujPA, 37, 39. Molonglo at 408 MHz (3′:S=54.2 Jy) and Parkes 64-m at 5 GHz (4′:S=18.5 Jy).
- Milne & Dickel 1975, AujPh, 28, 209. Parkes 64-m at 5 GHz (4.4′:S=14.6 Jy).
- Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2.6′).
- Caswell *et al.* 1980, MNRAS, 190, 881. FIRST at 1415 MHz (50′′).
- Weiler & Panagia 1980, A&A, 90, 269. Effelsberg 100-m at 9 GHz (1.5′) (private communication from Baker).
- Shaver *et al.* 1985, Nature, 313, 113. VLA at 1.4 GHz (3′′8×10′′9) and 5 GHz (12′′×26′′).
- Becker & Helfand 1985, Nature, 313, 115. VLA at 1.4 GHz and 5 GHz.
- Helfand & Becker 1985, Nature, 313, 118. Suggest it is not a SNR.
- Shaver *et al.* 1985, A&A, 147, L23. Observations of peripheral compact source.
- Caswell *et al.* 1989, PASA, 8, 184. MOST at 843 MHz (43′′×83′′).
- Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43′′×84′′).
- Stewart *et al.* 1994, ApJ, 432, L39. ATCA at 4.79 and 5.84 GHz (12′′×22′′) and Effelsberg 100-m at 10.6 GHz (1′), including polarisation.
- Frail *et al.* 1996, AJ, 111, 1651. OH emission near remnant, including masers.
- Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
- LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24′′×43′′).
- see also:* LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
- Brogan *et al.* 2000, ApJ, 537, 875. VLA at 1.7 GHz for OH Zeeman splitting.
- Brogan & Goss 2003, AJ, 125, 272. VLA at 1.4 GHz (11′′4×13′′6), including H_i, and 8.3 GHz (6′′8×14′′3) recombination line observation of H_{II} region.
- Gaensler *et al.* 2003, ApJ, 594, L35. Chandra detection.
- Lazendic *et al.* 2003, AN, 324 (No S1), 157. Molecular line observations.
- Burton *et al.* 2004, MNRAS, 348, 638. IR and radio observations of H_{II} region.
- Lazendic *et al.* 2004, MNRAS, 354, 393. IR and molecular line observations.
- Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
- Phillips *et al.* 2009, MNRAS, 397, 1215. Observations of interactions with surroundings.
- Castro *et al.* 2013, ApJ, 774, 36. Fermi observations.
- Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
- Acerro *et al.* 2016, ApJS, 224, 8. Fermi observations.
- Qiao *et al.* 2018, ApJS, 239, 15. OH maser observations.

G357.7+0.3**RA:** 17^h38^m35^s
Dec: –30°44′**1-GHz flux/Jy:** 10
Spectral index: 0.4?**Size/arcmin:** 24
Type: S

Radio: Non-thermal shell in complex region.

References:

Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz ($4'3'' : S = 7 \pm 1.5$ Jy), $S_{5 \text{ GHz}} = 5.5 \pm 1.5$ Jy from surveys.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 84''$).
 Yusef-Zadeh *et al.* 1999, ApJ, 527, 172. VLA of nearby OH masers.
 Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Phillips & Marquez-Lugo 2010, MNRAS, 409, 701. Spitzer observations of region.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 Rho *et al.* 2017, ApJ, 834, 12. Molecular line observations.

G358.0 + 3.8

RA: 17^h26^m00^s
Dec: -28°36'

1-GHz flux/Jy: 1.5?
Spectral index: ?

Size/arcmin: 38
Type: S

Radio: Faint shell.

References:

Duncan *et al.* 1995, MNRAS, 277, 36. Parkes 64-m at 2.4-GHz ($10'4$).
 Bhatnagar 2000, MNRAS, 317, 453. GMRT at 327 MHz ($1'3'' \times 2'2'' : S = 2.5 \pm 1.3$ Jy), and NVSS at 1.4 GHz.

G358.1 + 1.0

RA: 17^h37^m00^s
Dec: -29°59'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 20
Type: S

Was erroneously called G358.1+0.1.

Radio: Faint shell.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz ($43'' \times 77''$).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1' : S = 6.0 \pm 2.5$ Jy).

G358.5 - 0.9

RA: 17^h46^m10^s
Dec: -30°40'

1-GHz flux/Jy: 4?
Spectral index: ?

Size/arcmin: 17
Type: S

Radio: Shell, brighter to NE.

References:

Gray 1994, MNRAS, 270, 835. MOST at 843 MHz ($43'' \times 77''$).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz ($1' : S = 8.0 \pm 2.5$ Jy).

G359.0–0.9

RA: 17^h46^m50^s
Dec: –30°16′

1-GHz flux/Jy: 23
Spectral index: 0.5

Size/arcmin: 23
Type: S

Radio: Incomplete shell.

Optical: Detected.

X-ray: Partial shell.

References:

Reich *et al.* 1988, IAUCom, 101, 293. Summary of parameters.
 Reich *et al.* 1990, A&AS, 85, 633. Effelsberg 100-m at 2.7 GHz (4′3).
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43″×86″).
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24″×43″).
see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8″4×12″8) of part.
 Bamba *et al.* 2009, ApJ, 691, 1854. Suzaku observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H₂ IR observations.
 de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.

G359.1–0.5

RA: 17^h45^m30^s
Dec: –29°57′

1-GHz flux/Jy: 14
Spectral index: 0.4?

Size/arcmin: 24
Type: S

Has been called G359.10–0.5.

Radio: Non-thermal shell in complex region, crossed by the ‘snake’.

Optical: Detected.

X-ray: Centrally brightened.

Point sources: Several compact radio sources near centre, OH masers around edge.

References:

Downes *et al.* 1979, A&AS, 35, 1. From observations by Altenhoff *et al.* 1979, A&AS, 35, 23. Effelsberg 100-m at 4.9 GHz (2′6:S=13 Jy).
 Reich & Fürst 1984, A&AS, 57, 165. Effelsberg 100-m at 2.7 GHz (4′3:S=10±1.5 Jy) and 4.8 GHz (2′4:S=8.1±0.5 Jy).
 Uchida *et al.* 1992, ApJ, 398, 128. VLA at 1.5 GHz (10″×11″), and observations of nearby molecular material.
 Uchida *et al.* 1992, AJ, 104, 1533. VLA at 1.4 GHz.
 Gray 1994, MNRAS, 270, 835. MOST at 843 MHz (43″×85″).
 Yusef-Zadeh *et al.* 1995, Science, 270, 1801. VLA at 1.4 GHz (31″×33″), and 1.7 GHz for OH survey.
 LaRosa *et al.* 2000, AJ, 119, 207. VLA at 333 MHz (24″×43″).
see also: LaRosa *et al.* 2000, AJ, 119, 3145. Erratum.
 Bamba *et al.* 2000, PASJ, 52, 259. ASCA observations.
 Lazendic *et al.* 2002, MNRAS, 331, 537. Observations of shocked molecular gas where the ‘snake’ crosses the remnant.
 Yusef-Zadeh *et al.* 2004, ApJS, 155, 421. VLA at 1.4 GHz (8″4×12″8).
 Aharonian *et al.* 2008, A&A, 483, 509. XMM-Newton and HESS observations.

Hewitt *et al.* 2008, ApJ, 683, 189. GBT at 1.6 and 1.7 GHz for OH masers.
 Bamba *et al.* 2009, ApJ, 691, 1854. Suzaku observations.
 Stupar & Parker 2011, MNRAS, 414, 2282. H α observations.
 Ohnishi *et al.* 2011, PASJ, 63, 527. Suzaku observations.
 Pihlström *et al.* 2014, AJ, 147, 73. VLA search for methanol masers.
 Ponti *et al.* 2015, MNRAS, 453, 172. XMM-Newton observations.
 Froebrich *et al.* 2015, MNRAS, 454, 2586. H $_2$ IR observations.
 Hui *et al.* 2016, MNRAS, 457, 4262. Fermi observations of region.
 de Wilt *et al.* 2017, MNRAS, 468, 2093. Molecular line observations of region.
 Qiao *et al.* 2018, ApJS, 239, 15. OH maser observations.

G359.1 + 0.9

RA: 17^h39^m36^s
Dec: -29°11'

1-GHz flux/Jy: 2?
Spectral index: ?

Size/arcmin: 12×11
Type: S

Radio: Shell, brightest in E.

References:

Gray 1994, MNRAS, 270, 847. MOST at 843 MHz (43''×88'': S=4.3 Jy).
 Roy & Bhatnagar 2006, JPhCS, 54, 152. GMRT at 330 MHz (1': S=4.3±1.0 Jy).
 Law *et al.* 2008, ApJS, 177, 515. VLA at 1.4 GHz (10''9×15''9: S=1.3±0.5 Jy).